Published in final edited form as:

Biol Psychol. 2017 February ; 123: 269–277. doi:10.1016/j.biopsycho.2016.10.009.

Neural Markers of Attention to Aversive Pictures Predict Response to Cognitive Behavioral Therapy in Anxiety and Depression

Jonathan P. Stange^a, Annmarie MacNamara^b, Olga Barnas^a, Amy E. Kennedy^{a,c}, Greg Hajcak^d, K Luan Phan^{a,c,e,f}, and Heide Klumpp^{a,e}

^aDepartment of Psychiatry, University of Illinois at Chicago, 1747 W. Roosevelt Rd., Chicago, IL 60608, USA

^bDepartment of Psychology, Texas A&M University, 4235 TAMU, College Station, TX 77840, USA

^cMental Health Service Line, Jesse Brown VA Medical Center, 820 S. Damen Ave., Chicago, IL 60612, USA

^dDepartment of Psychology, Stony Brook University, Psychology B Building, Stony Brook, NY 11794, USA

^eDepartment of Psychology, University of Illinois at Chicago, 1007 W. Harrison St., Chicago, IL 60607, USA

^fDepartment of Anatomy and Cell Biology, and the Graduate Program in Neuroscience, University of Illinois at Chicago, 808 S. Wood St., Chicago, IL 60612, USA

Abstract

Excessive attention toward aversive information may be a core mechanism underlying emotional disorders, but little is known about whether this is predictive of response to treatments. We evaluated whether enhanced attention toward aversive stimuli, as indexed by an event-related potential component, the late positive potential (LPP), would predict response to cognitive behavioral therapy (CBT) in patients with social anxiety disorder and/or major depressive disorder. Thirty-two patients receiving 12 weeks of CBT responded to briefly-presented pairs of aversive and neutral pictures that served as targets or distracters while electroencephalography was recorded. Patients with larger pre-treatment LPPs to aversive relative to neutral distracters (when targets were aversive) were more likely to respond to CBT, and demonstrated larger reductions in symptoms of depression and anxiety following treatment. Increased attention toward irrelevant aversive stimuli may signal attenuated top-down control, so treatments like CBT that improve this control could be beneficial for these individuals.

Correspondence to: Heide Klumpp, PhD, Department of Psychiatry, University of Illinois at Chicago, 1746 West Roosevelt Rd., Chicago, IL 60608, Phone: (312) 996-0416, hklumpp@psych.uic.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

event-related potentials; transdiagnostic; internalizing disorders; CBT; treatment prediction

Introduction

Social anxiety disorder (SAD) and major depressive disorder (MDD) are prevalent, frequently comorbid and highly impairing (Kessler et al., 2003, 2005, 2006; Mineka et al., 1998; Kaufman & Charney, 2000; Stein et al., 2001; Beesdo et al., 2007). Excessive attention toward aversive information has been proposed as a core mechanism underlying these emotional disorders (Mathews & MacLeod, 2005; Bar-Haim et al., 2007). Cognitive-behavioral therapy (CBT) is a gold-standard psychosocial treatment for anxiety and depressive disorders that targets emotional disorders by facilitating coping with aversive emotions, and that may change how threat is processed (Beck et al., 1979; Hofmann et al., 2012). Although CBT has demonstrated moderate effectiveness for emotional disorders, not all patients benefit equally from CBT, and many patients remain symptomatic after an initial intervention (Hofmann & Smits, 2008; Kemp et al., 2008). Identifying patient characteristics associated with response to CBT may lead to more personalized treatment decision-making (Paulus, 2015).

Given the extremely high rates of comorbidity between SAD and MDD, it is likely that common factors, such as heightened negative affectivity and increased attention toward negative environmental information, may underlie both disorders (Gibb et al., 2015; Mathews & MacLeod, 2002; Pessoa et al., 2002). As a result, CBT involves similar treatment strategies for SAD and MDD, including cognitive restructuring about real or potential negative situations, and encouraging exposure to environmental situations that are perceived as negative or undesirable (Beck & Bredemeier, 2016; Rodebaugh et al., 2004). Reducing the salience of negative emotional or environmental stimuli that interfere with situational goals thus is one aim of treatment, and the degree to which such stimuli are salient could serve as an indicator of which patients are most likely to benefit from CBT.

Given the high degree of comorbidity and the possibility of partially overlapping mechanisms of illness, it will be important to identify common factors predicting treatment response across SAD and MDD. One potential predictor of treatment response is increased attention toward aversive stimuli (Eysenck, 1992; Mathews & MacLeod, 2002; Peckham et al., 2010; Pessoa et al., 2002; Gibb et al., 2015). Elevated salience of aversive stimuli may lead to excessive bottom-up processing of goal-irrelevant, sensory-driven stimuli, at the expense of top-down control to attend to the goal at hand (Pessoa et al., 2002). Increased attention to aversive stimuli has been found across multiple emotional disorders (Bar-Haim et al., 2007), including SAD (Kircansky et al., 2015) and MDD (Peckham et al., 2010; but see Weinberg et al., 2016; Kircansky et al., 2015). Neural measures, such as functional magnetic resonance imaging (fMRI), provide promising tools for assessing attention to aversive stimuli (Doehrmann et al., 2013; Klumpp et al., 2013; Fu et al., 2008; Whalen et al., 2008). In comparison to fMRI, event-related potentials (ERPs) such as the late positive potential (LPP), assessed by EEG, provide a less expensive and clinically practical means of

elucidating the processing of negatively-valenced information in emotional disorders (Bar-Haim et al., 2005; Hajcak et al., 2012; MacNamara et al., 2011, 2013). Prior work has demonstrated that the LPP to aversive stimuli is elevated among individuals with anxiety disorders (MacNamara & Hajcak, 2010; Li et al., 2007), with or without the presence of comorbid depression (MacNamara & Proudfit, 2014; Dillon et al., 2013; Brown, 2007; Desseilles et al., 2009, 2011). A more limited literature in depression has suggested that depression without anxiety may be characterized by attenuated LPPs to motivationally-salient stimuli (Proudfit et al., 2015). Thus, measurement of the attentional processing of aversive stimuli using the LPP may help to elucidate the neural mechanisms underlying SAD and MDD (e.g., Gibb et al., 2015), and could serve as useful measures of propensity to benefit from treatment.

Despite evidence that shared mechanisms may underlie SAD and MDD (Dillon et al., 2013; Mineka et al., 1998) and increasing interest in understanding neural predictors of treatment response (Andreescu & Aizenstein, 2016; Paulus, 2015) across traditional diagnostic groups (Cuthbert, 2014), limited work has evaluated whether brain-based measures of attention to aversive stimuli prior to treatment could predict response to CBT for these disorders. Prior work has suggested that response to treatment for anxiety and depression is associated with reductions in attention toward aversive stimuli (Etkin & Schatzberg, 2011; Pishyar et al., 2008); therefore, neural activity associated with attention toward aversive stimuli might be helpful in predicting who is most likely to benefit from such treatments (e.g., Doehrmann et al., 2013; Klumpp et al., 2013, 2014). In line with this hypothesis, prior neuroimaging work has demonstrated that greater higher-order visual cortex activation for negative stimuli prior to treatment predicted better response to CBT for social anxiety (Doehrmann et al., 2013; Klumpp et al., 2013). Other studies have found that response to CBT was predicted by greater pre-treatment reactivity in prefrontal cortical areas in youth with anxiety disorders (Kujawa et al., 2016), in rostral anterior cingulate cortex (ACC) to fearful faces among adults with generalized anxiety disorder (Whalen et al., 2008), and in dorsal ACC activity to sad faces among depressed adults (Fu et al., 2008). Furthermore, simultaneous fMRI/EEG studies have suggested that activation in these regions (particularly the visual cortices) may represent a key neural source contributing to the LPP elicited by aversive stimuli (Liu et al., 2012; Sabatinelli et al., 2007); however, few studies have evaluated the LPP – a relatively cost-effective and well-tolerated neural measure – as a predictor of treatment outcome in the anxiety and depressive disorders. One extant study of individuals with spider phobia (Leutgeb et al., 2009) found treatment-related increases in the LPP for aversive stimuli, suggesting that a higher LPP to aversive stimuli may be indicative of less avoidance (e.g., a willingness to engage with aversive stimuli; Weinberg & Hajcak, 2011) and better outcomes. Few studies of depression have evaluated the LPP as a predictor of treatment outcome. However, neuroimaging studies have found evidence that greater pre-treatment activation in the amygdala (Canli et al., 2005; Siegle et al., 2006) and the temporal cortex (Ritchey et al., 2011) is associated with an improved course of depressive symptoms and greater response to CBT. Together, these results suggest that individuals who show neural correlates of enhanced attention toward aversive, goal-irrelevant stimuli may be particularly likely to benefit from CBT.

In the present study, we evaluated whether individual differences in attention to aversive stimuli (as indexed by the LPP) presented in attended or unattended locations would be associated with reduced illness severity following CBT for SAD or MDD, using a task previously shown to differentiate individuals with anxiety from those without (MacNamara & Hajcak, 2009, 2010). As in prior work (MacNamara & Hajcak, 2009, 2010), we expected that LPPs would be greater for aversive stimuli than for neutral stimuli when presented in attended locations, but not when stimuli were presented in unattended locations. Given prior work demonstrating that greater LPPs to aversive stimuli are associated with anxiety (MacNamara & Hajcak, 2009, 2010), and fMRI results suggesting that greater attention to aversive stimuli is associated with improved CBT outcomes (Doehrmann et al., 2013; Klumpp et al., 2013; Fu et al., 2008; Whalen et al., 2008; Kujawa et al., 2016; Canli et al., 2005; Siegle et al., 2006; MacNamara & Hajcak, 2010), we hypothesized that individuals with larger LPPs to aversive stimuli would be more likely to respond to CBT, and would show larger decreases in symptoms of anxiety and depression, relative to individuals with smaller LPPs to aversive stimuli. Given that prior work demonstrated associations between anxiety and attention to aversive targets (MacNamara & Hajcak, 2009, 2010), we expected that greater attention to aversive targets would predict better treatment outcome; we did not have a priori hypotheses about whether treatment outcome would be associated with LPPs to aversive distracters. Distracters were included in prior studies using this task and in the current study to explore whether attention to aversive stimuli would predict treatment response differentially as a function of the relevance of the stimuli to the current goal (i.e., attending to targets, not distracters).

Method

Participants

All participants met Diagnostic and Statistical Manual of Mental Disorders (DSM-IV) criteria for a current diagnosis of social anxiety disorder or major depressive disorder (see Table 1). All participants were free of psychotropic medication for at least 8 weeks prior to, and throughout, the study. Exclusion criteria were as follows: a) substance abuse or dependence in the prior six months, b) history of bipolar disorder or schizophrenia, or the presence of an organic mental syndrome, intellectual disability, or pervasive developmental disorder, c) ongoing psychotherapy and/or current treatment with any psychotropic medication, and d) clinically significant medical or neurologic condition. Participants were between 18 and 55 years of age and right-handed. The study protocol was approved by the Institutional Review Boards of the University of Michigan Medical School and the University of Illinois at Chicago, and all participants provided written informed consent.

Materials and Measures

Diagnostic Interview—Participants were interviewed by Master's- or Doctoral-level clinicians using the Structured Clinical Interview for DSM-IV (SCID-IV) (First et al., 1996) to assess Axis I disorders (see Table 1).

Treatment Outcome Measures—To assess illness severity and response to CBT, clinicians completed the Clinical Global Impression (CGI) Severity and Improvement scales

(Busner & Targum, 2007). Both measures use 7-point scales, ranging from 1 (normal, not at all ill) to 7 (extremely ill) for CGI Severity and from 1 (very much improved) to 7 (very much worse) for CGI Improvement. As in prior work (Barlow et al., 2000), we used the CGI-Severity and CGI-Improvement scales in conjunction to determine degree of treatment response. Participants with scores < 3 on both scales were determined to have achieved clinically-significant treatment response and were categorized as “Responders,” while those with scores ≥ 3 were classified as “Non-Responders.”

The 17-item Hamilton Depression Rating Scale (HAM-D; Hamilton, 1960), a widely-used interview-based measure of depression symptom severity and the Hamilton Anxiety Rating Scale (HAM-A; Hamilton, 1959), a 14-item clinician-administered measure of severity of anxious symptomatology, were administered by trained, independent evaluators at pre- and post-treatment to assess changes in symptoms of depression and anxiety, respectively.

Affective pictures—Forty-eight aversive (e.g., attack scenes, mutilated bodies) and 48 neutral pictures (e.g., household objects, neutral faces) were selected from the International Affective Picture System (Lang et al., 2005). In prior studies validating this task, aversive pictures were rated as less pleasant and higher in arousal than neutral pictures (see MacNamara & Hajcak, 2009, for details). Stimuli were presented on a Dell Optiplex 750 computer, using Presentation software (Neurobehavioral Systems, Inc.). Participants were seated approximately 60 cm from the screen.

Procedures

CBT—Patients received 12 weeks of manualized, individual CBT conducted by doctoral-level clinical psychologists (Beck et al., 1979; Craske et al., 1992; Hope et al., 2006; Martel et al., 2010). A licensed clinical psychologist with expertise in CBT and in clinical trial investigations involving CBT provided supervision to ensure adherence to treatment. CBT included psychoeducation, cognitive restructuring, in vivo exposures, behavioral activation, and relapse prevention. The specific type of CBT provided was targeted toward each patient's primary diagnosis.

Affective picture task—Pre-treatment, participants completed a computerized task while EEG was recorded. In brief, four pictures – two to the left and right, and two above and below the center of the screen – were presented simultaneously on each trial; participants were asked to indicate whether two of the pictures (either the vertical or horizontal picture pairs) were the same or different. Picture valence (aversive or neutral) was always the same in both the horizontal and vertical pairs. From here on, stimuli presented in task-relevant spatial locations will be referred to as “targets,” and those presented in task-irrelevant locations will be referred to as “distracters.”

There were four trial types: neutral targets paired with neutral distracters, neutral targets paired with aversive distracters, aversive targets paired with neutral distracters, and aversive targets paired with aversive distracters. Participants used the left and right mouse buttons (counterbalanced across participants) to indicate if targets were identical (“same”) or different (“different”); participants were encouraged to respond as quickly and as accurately as possible. Before each trial, two white rectangles appeared on a black background for

1,000 ms to indicate which picture pair (horizontal or vertical) would be the targets for the same/different decision in the upcoming trial; pictures were displayed in color for 250 ms. Participants completed 10 practice trials and 320 experimental trials. Pictures presented during practice trials were not repeated during experimental trials. Trial order and pictures were presented pseudo-randomly, with each picture repeated 10 times across the task (for more details see MacNamara & Hajcak, 2009).

Electroencephalographic Recording and Behavioral Responses

An elastic cap and the ActiveTwo BioSemi system (Amsterdam, Netherlands) were used to record the continuous EEG. Thirty-four electrode sites (standard 32 channel setup plus Iz and FCz) based on the 10/20 system, were used, with one additional electrode on each of the left and right mastoids. Four facial electrodes recorded the electrooculogram generated from eye blinks and eye movements: vertical eye movements and blinks were measured with two electrodes placed approximately 1 cm above and below the right eye; horizontal eye movements were measured with two electrodes placed approximately 1 cm beyond the outer edge of each eye. Online data were referenced according to BioSemi's design using two separate electrodes for grounding (the Common Mode Sense active electrode and the Driven Right Leg passive electrode) and data were digitized at 1024 Hz.

Off-line analyses were performed using Brain Vision Analyzer (Brain Products, Gilching, Germany); data were re-referenced to the average of the two mastoids and were band-pass filtered with low and high cutoffs of 0.01 and 30Hz, respectively. The baseline for each trial was defined as the 200ms before picture onset. ERPs were segmented for each trial beginning 200ms before picture onset until 1200ms (1000ms beyond picture onset). Eye blink and ocular corrections were made using the algorithm developed by Miller et al., (1988). Artifact analysis identified a voltage step of more than 50 μ V between sample points, a voltage difference of 300 μ V within a trial, and a maximum voltage difference of less than 0.50 μ V within 100 ms intervals. Trials were also inspected visually for any remaining artifacts. Intervals containing artifacts were rejected from individual channels in each trial. As in prior work, the LPP was scored by averaging activity from 400 to 1000ms at four centro-parietal sites where the LPP was maximal: CP1, CP2, Cz, and Pz (e.g., Weinberg & Hajcak, 2010; Hajcak et al., 2007). Averages of LPPs for each trial type (80 trials in each of the four conditions noted above) were created for each participant. Only trials associated with a correct response made within 1,800 ms following picture offset were included in the ERP analyses. The average reaction time (RT) per condition was determined as the average time taken to respond following picture onset on correct trials and accuracy was assessed as the percentage of correct responses per condition.

Participants generally performed well on the task ($M = 89.44\%$ correct, $SD = 8.84\%$). One (female) participant was removed from analyses because of excessive EEG artifacts (> 50% of trials excluded), and two participants (one male, one female) were excluded because of poor task performance (less than 50% accuracy), yielding a final sample of $n = 32$ for analyses.

Statistical Analyses

Task effects on the LPP, reaction time, and accuracy were evaluated with 2 (target type: neutral, aversive) $\times$ 2 (distracter type: neutral, aversive) repeated-measures analyses of variance (ANOVAs). To evaluate whether attention to aversive stimuli at pre-treatment predicted treatment outcome, we performed three analyses of covariance (ANCOVA). First a 2 (target type: neutral, aversive) $\times$ 2 (distracter type: neutral, aversive) ANCOVA was conducted, with pre-treatment CGI-Severity entered as a covariate of no interest, and post-treatment CGI Responder status as a covariate of interest. Next, we conducted the same 2 $\times$ 2 ANCOVA, but with pre-treatment HAM-D scores (instead of CGI-Severity) entered as a covariate of no interest, and post-treatment HAM-D (instead of CGI Responder status) as a continuous covariate of interest. Finally, we conducted the same 2 $\times$ 2 ANCOVA, this time controlling for pre-treatment HAM-A scores (as a covariate of no interest), and with post-treatment HAM-A as a continuous covariate of interest.

To follow up on significant ANCOVAs, post-hoc tests involved regressions for each of the four possible pairwise comparisons between trial types, predicting post-treatment symptoms (linear regressions) or recovery status (logistic regression): (1) aversive minus neutral distracter when targets were neutral; (2) aversive minus neutral distracter when targets were negative; (3) aversive minus neutral target when distracters were neutral; and (4) aversive minus neutral target when distracters were aversive.

Results

Sample demographics and clinical characteristics are detailed in Table 1.

Changes in Clinical Measures Across Treatment

CGI-Severity scores decreased from pre-treatment to post-treatment ($t(31) = 6.80, p < .001, d = 1.50$; Table 1). Based on the conservative approach of combining CGI-Severity and CGI-Improvement indices to determine treatment response (per Barlow et al., 2000), 36% of the sample (14 of 39 patients) were classified as Responders (i.e., rated “normal, not at all ill” or “borderline mentally ill” at post-treatment on CGI-Severity and “very much improved” or “much improved” at post-treatment). Patients' primary diagnosis was not significantly associated with treatment responder status (SAD primary: $n = 12$ responders (57.1%); MDD primary: $n = 3$ responders (33.3%); $\chi^2(1) = 0.75, p = .39$). There were also significant decreases in HAM-D ($t(31) = 5.20, p < .001, d = 1.05$) and in HAM-A ($t(31) = 7.25, p < .001, d = 1.35$) from pre-treatment to post-treatment. Symptom scores as a function of CGI responder status are displayed in Table 2. At baseline, CGI responders had greater severity on the CGI ($t(20) = 2.86, p = .01, d = 0.89$).

Task Effects

Figure 1a depicts grand-average waveforms for each of the four trial types at centro-parietal sites where the LPP was scored (i.e., the average of CP1, CP2, Cz, and Pz); Figure 1b displays scalp distributions for each trial type, from 400-1,000 ms post-picture onset. As expected based on prior work (MacNamara & Hajcak, 2009, 2010), there was a significant effect of target type on the LPP ($F(1,31) = 29.86, p < .001, \eta_p^2 = .49$), such that larger LPPs

were elicited for aversive ($M = 4.64 \mu\text{V}$, $SD = 4.27 \mu\text{V}$) relative to neutral targets ($M = 2.53 \mu\text{V}$, $SD = 4.15 \mu\text{V}$). However, there was no main effect of distracter type ($F(1,31) = 0.04$, $p = .95$, $\eta_p^2 = .0001$; aversive $M = 3.57 \mu\text{V}$, $SD = 4.19 \mu\text{V}$; neutral $M = 3.60 \mu\text{V}$, $SD = 4.20 \mu\text{V}$), nor was there an interaction between target and distracter type ($F(1,31) = 0.05$, $p = .82$, $\eta_p^2 = .002$).

In terms of RT for accurate trials, there was a marginal main effect of target type ($F(1,31) = 3.30$, $p = .08$, $\eta_p^2 = .10$) such that aversive targets ($M = 687.03$ ms, $SD = 104.95$ ms) were associated with longer RTs than neutral targets ($M = 679.53$ ms, $SD = 104.81$ ms). The main effect of distracter type was not significant ($F(1,31) = 0.24$, $p = .63$, $\eta_p^2 = .008$), but there was a significant interaction between target and distracter type ($F(1,31) = 8.91$, $p < .01$, $\eta_p^2 = .22$; neutral targets with neutral distracters: $M = 668.36$ ms, $SD = 110.15$ ms; neutral targets with aversive distracters: $M = 672.70$ ms, $SD = 102.46$ ms; aversive targets with neutral distracters: $M = 682.21$ ms, $SD = 105.43$ ms; aversive targets with aversive distracters: $M = 691.85$ ms, $SD = 106.16$ ms). Post-hoc comparisons revealed that when targets were negative, negative distracters were associated with longer RTs ($p = .05$), whereas when targets were neutral, negative distracters were associated with shorter RTs ($p = .04$). When distracters were negative, negative targets were associated with longer RTs ($p < .01$), whereas when distracters were neutral, distracter valence was not associated with RT ($p = .39$). No significant effects were observed for accuracy (all $ps > .24$).

Treatment Response Prediction

Figure 2 depicts LPP amplitudes for all conditions and spatial distributions for aversive relative to neutral distracters (under conditions of aversive targets) for Non-Responders (Figure 2a) and Responders (Figure 2b).¹ There was a significant three-way interaction between target type, distracter type, and CGI responder status for the LPP ($F(1,29) = 4.87$, $p = .04$, $\eta_p^2 = .14$), after controlling for pre-treatment CGI-Severity. Post-hoc tests indicated that when targets were aversive, larger LPPs for aversive (relative to neutral) distracters were associated with a greater likelihood of recovery (Wald $\chi^2(1) = 4.35$, $p = .04$, $OR = 1.44$, $CI = 1.02-2.03$, Nagelkerke $R^2 = .21$) (Figure 2). None of the other trial-type differences (i.e., aversive minus neutral distracter when targets were neutral; aversive minus neutral target when distracters were neutral/aversive) were associated with recovery status ($ps > .13$).

Similarly, there was a significant three-way interaction between target type, distracter type, and post-treatment HAM-D ($F(1,29) = 6.23$, $p = .02$, $\eta_p^2 = .18$), after controlling for pre-treatment HAM-D scores. Larger LPPs for aversive (relative to neutral) distracters when targets were aversive were associated with lower levels of depression at post-treatment ($\beta = -0.36$, $t = -2.28$, $p = .03$, $R^2 = .13$) (Figure 3a). None of the other pairwise trial-type differences were associated with post-treatment HAM-D ($ps > .05$).

There also was a significant three-way interaction between target type, distracter type, and post-treatment HAM-A ($F(1,29) = 5.29$, $p = .03$, $\eta_p^2 = .15$), after controlling for pre-

¹Per a reviewer's suggestion, we also examined whether responders and non-responders differed in the N2, an ERP that is thought to represent cognitive control functioning. The N2 was maximal between 250-300ms at Fc1, Fc2, Cz, and Fcz, for which a pooled variable was created for each of the four trial types. Responders and non-responders did not differ on N2 amplitudes as a function of target valence, distracter valence, or their interaction.

treatment HAM-A. Post-hoc tests indicated that when targets were aversive, larger LPPs for aversive (relative to neutral) distracters were associated at a trend level with lower levels of anxiety symptoms at post-treatment ($\beta = -0.31$, $t = -1.96$, $p = .06$, $R^2 = .09$) (Figure 3b). None of the other three pairwise trial-type differences were associated with post-treatment HAM-A ($ps > .15$).

All treatment prediction results reported above maintained significance when accounting for patients' primary diagnosis (SAD vs. MDD; ANCOVA 3-way interaction terms $ps < .05$). There were no significant interactions between outcome variables and target type, distracter type, or the target $\times$ distracter interaction for response time ($ps > .14$) or accuracy ($ps > .16$).

Discussion

We evaluated neural markers of attention to aversive stimuli (as indexed by the LPP) as a predictor of response to CBT for SAD or MDD. Results showed that patients with larger LPPs for aversive relative to neutral distracters (when targets were aversive), showed greater response to CBT. These results were evident when using responder status based on CGI ratings, as well as when using continuous changes in depression and anxiety on the HAM-D and HAM-A, respectively. Thus, the LPP at pre-treatment predicted reductions in symptoms evident across a number of clinical measures. These findings support the utility of assessing negative valence systems transdiagnostically using neurobiological measures, to evaluate predictors of treatment outcome, in search of a personalized approach to the treatment of anxiety and depressive disorders (Sanislow et al., 2010; Simpson, 2012; Gibb et al., 2015).

In line with prior studies that used fMRI (Doehrmann et al., 2013; Klumpp et al., 2013; Kujawa et al., 2015), we found that greater attention toward aversive stimuli at pre-treatment predicted superior response to CBT. Although speculative, it is possible that individuals who demonstrate increased attention toward aversive stimuli have a greater tolerance for (or are less avoidant of) aversive emotions, which could facilitate engagement with (or habituation to) these target emotions as part of CBT and therefore could lead to improved treatment response. Alternatively, individuals with greater attention toward aversive stimuli may benefit more from participating in treatments that involve engagement with difficult emotions because these treatments target the reduction of these very characteristics (i.e., one of the aims of CBT is to improve increase tolerance of negative thoughts and emotions). Consistent with this hypothesis, emotional disorders may impair the recruitment of prefrontal regions and filtering of negative, task-irrelevant information (Bishop et al., 2004 a,b, 2009; MacNamara et al., 2011; Peckham et al., 2010). Thus, larger LPPs to aversive distracters may signal attenuated top-down control of attention to aversive stimuli (Pessoa et al., 2002), suggesting that treatments that aim to improve this top-down control might be a particularly good match for these patients. That treatment outcome was most evident in the presence of negative targets and negative distracters suggests that conditions containing a high load of aversive stimuli (and thus an implicit need for regulation) may be necessary for differentiating which individuals are most likely to benefit from CBT. It also suggests the importance of assessing attention to aversive stimuli using behavioral tasks that contain valenced targets and distracters, as treatment outcome may not have been evident in the absence of distracters in the task used here.

In contrast with our findings, a related literature has suggested that cognitive flexibility is longitudinally associated with improved symptom course in anxiety and depression (e.g., Johnco et al., 2014; Stange et al., 2016). For example, a recent paper demonstrated that superior cognitive flexibility, as indexed by the N2 ERP, was associated with less-distressing intrusive symptoms following an analog trauma (Streb et al., 2016). In our study, the behavioral task evaluated engagement with aversive stimuli in variable locations (targets and distracters) using the LPP, rather than measuring cognitive flexibility per se, which could account for the different pattern of results found here. In addition, naturalistic predictors of symptom course are not necessarily the same as those that may predict response to treatment; for example, cognitive flexibility in general could facilitate reductions in symptom course in naturalistic contexts, but inflexibility with respect to emotional stimuli could represent a target representing greater room for improvement with treatment for emotional disorders. Thus, future research is needed to clarify the contexts in which cognitive flexibility and engagement with emotional stimuli may be associated with symptom course and outcome in naturalistic and treatment contexts.

The study involved a diagnostically-mixed sample of patients receiving treatment for primary SAD or MDD, suggesting that the mechanisms by which larger LPPs to aversive stimuli might be associated with treatment outcome could be at least partially overlapping for these disorders. Larger LPPs to aversive stimuli may indicate less avoidance of these stimuli, which may facilitate habituation in exposure-based treatments (e.g., Jaycox et al., 1998); individuals with greater avoidance (or smaller LPPs to aversive stimuli) may, on the other hand, be less ready for or less able to benefit from CBT (Tillfors et al., 2015; Waters & Kershaw, 2015). Patients who show a blunted pattern of responding to aversive stimuli (e.g., Proudfit et al., 2015; Bruder et al., 2012) may have difficulty with experiencing and modifying negative emotions in therapy, leading to poorer treatment response. Prior work has documented opposite patterns of LPPs to aversive stimuli among individuals with anxiety and depression in the absence of treatment (e.g., MacNamara et al., 2015; Proudfit et al., 2015). However, the mechanisms discussed here could help to explain why our pattern of results was consistent transdiagnostically and when predicting symptoms of anxiety and depression. Thus, across anxiety and depressive disorders, larger LPPs for aversive relative to neutral stimuli could be an indicator of the ability to engage with difficult emotions (Weinberg & Hajcak, 2011; Proudfit et al., 2013), which previously has been shown to facilitate response to CBT (Jaycox et al., 1998; Kashdan et al., 2006; Whelton, 2004).

This was the first study to identify the LPP as a predictor of response to CBT among a heterogeneous sample of patients with emotional disorders. Nevertheless, several limitations should be noted. The sample size was relatively small, which prevented us from determining if results differed by primary diagnosis. In addition, the lack of a wait-list control group means that our findings could be predictive of symptom-based change more broadly (e.g., remittance of symptoms due to the passage of time), rather than CBT-based change in particular. Future studies also would benefit from employing multiple treatments and control conditions to determine whether the LPP can be used to identify which patients are most likely to benefit from one treatment versus another, with the goal of personalized medicine (Tracy, Klonsky, & Proudfit, 2014). Although predictors of treatment response are necessarily the same as those that are changed by treatment (e.g., Doehrmann et al., 2013;

Klump et al., 2013; MacNamara et al., 2015; Phan et al., 2013), examining the degree to which neural responses change following treatment would help to elucidate these questions. It also will be important to evaluate the extent to which findings converge across different tasks and stimulus/distracter types (e.g., idiographic, loss-related stimuli). Relatedly, we did not include a passive viewing task in the present study, so we are not able to specify whether similar results would be found in this context. It is possible that rapidly presented stimuli are more likely to elicit larger LPPs among anxious individuals (e.g., MacNamara & Hajcak, 2009, 2010) than are pictures that are passively viewed (e.g., Weinberg & Hajcak, 2011); however, future work is needed to determine whether rapidly presented (vs. passively viewed) stimuli might be differentially associated with treatment response. Finally, although post-hoc tests only were conducted when interaction analyses were statistically significant, due to the preliminary nature of this study we did not apply correction for multiple comparisons when testing the simple effects of interactions.

In conclusion, the data provide evidence that patients with larger LPPs to aversive relative to neutral stimuli may be particularly likely to benefit from CBT for anxiety or depression. Results were not observed for behavioral measures, in line with the notion that neural measures may provide particularly sensitive means of assessing elaborative attentional processing of emotional stimuli (e.g., Doehrmann et al., 2013; Kujawa et al., 2016) and underscoring the importance of including such measures in future studies of treatment outcome prediction (Tracy et al., 2014).

Acknowledgments

This work was supported by National Institute of Mental Health (NIMH) K23MH093679 and Brain and Behavior Research Foundation (formerly NARSAD) Award to HK and in part by NIMH R01MH101497 (to KLP) and the Center for Clinical and Translational Research (CCTS) grant, UL1RR029879. JS is supported by NIMH 5T32MH067631-12. AM is supported by NIMH K23MH105553.

References

- Andreescu C, Aizenstein H. Predicting treatment response with functional magnetic resonance imaging. *Biological Psychiatry*. 2016; 79:262–263. [PubMed: 26796875]
- Barlow DH, Gorman JM, Shear MK, Woods SW. Cognitive-behavioral therapy, imipramine, or their combination for panic disorder: A randomized controlled trial. *JAMA*. 2000; 283:2529–36. [PubMed: 10815116]
- Bar-Haim Y, Lamy D, Glickman S. Attentional bias in anxiety: a behavioral and ERP study. *Brain and Cognition*. 2005; 59:11–22. [PubMed: 15919145]
- Bar-Haim Y, Lamy D, Pergamin L, Bakermans-Kranenburg MJ, van IJzendoorn MH. Threat-related attentional bias in anxious and nonanxious individuals: a meta-analytic study. *Psychological Bulletin*. 2007; 133:1–24. [PubMed: 17201568]
- Beck AT, Bredemeier K. A unified model of depression integrating clinical, cognitive, biological, and evolutionary perspectives. *Clinical Psychological Science*. 2016; 4(4):596–619.
- Beck, AT., Rush, AJ., Shaw, BF., et al. *Cognitive therapy of depression*. New York: Guilford; 1979. p. 1979
- Beesdo K, Bittner A, Pine DS, et al. Incidence of social anxiety disorder and the consistent risk for secondary depression in the first three decades of life. *Archives of General Psychiatry*. 2007; 64(8): 903–912. [PubMed: 17679635]
- Bishop S, Duncan J, Brett M, Lawrence AD. Prefrontal cortical function and anxiety: controlling attention to threat-related stimuli. *Nature Neuroscience*. 2004; 7:184–8. [PubMed: 14703573]

- Bishop SJ. Trait anxiety and impoverished prefrontal control of attention. *Nature Neuroscience*. 2009; 12:92–8. [PubMed: 19079249]
- Bishop SJ, Duncan J, Lawrence AD. State anxiety modulation of the amygdala response to unattended threat-related stimuli. *Journal of Neuroscience*. 2004; 24:10364–8. [PubMed: 15548650]
- Brown TA. Temporal course and structural relationships among dimensions of temperament and DSM-IV anxiety and mood disorder constructs. *Journal of Abnormal Psychology*. 2007; 116:313–28. [PubMed: 17516764]
- Bruder, GE., Kayser, J., Tenke, CE. Event-related brain potentials in depression: Clinical, cognitive and neurophysiologic implications. In: Luck, SJ., Kappenman, ES., editors. *The Oxford handbook of event-related potential components*. New York Oxford University Press: 2012. p. 563-592.
- Burkhouse KL, Kujawa A, Kennedy AE, et al. Neural reactivity to reward as a predictor of cognitive behavioral therapy response in anxiety and depression. *Depression and Anxiety*. in press.
- Busner J, Targum SD. The clinical global impressions scale. *Psychiatry*. 2007; 4:28–37.
- Canli T, Cooney RE, Goldin P, et al. Amygdala reactivity to emotional faces predicts improvement in major depression. *Neuroreport*. 2005; 16:1267–70. [PubMed: 16056122]
- Craske, MG., Barlow, DH., O'Leary, TA. *Mastery of your anxiety and worry*. Albany: Graywind Publications; 1992.
- Cuthbert BN. The RDoC framework: facilitating transition from ICD/DSM to dimensional approaches that integrate neuroscience and psychopathology. *World Psychiatry*. 2014; 13(1):28–35. [PubMed: 24497240]
- Desseilles M, Schwartz S, Dang-Vu TT, et al. Depression alters “top-down” visual attention: a dynamic causal modeling comparison between depressed and healthy subjects. *Neuroimage*. 2011; 54:1662–8. [PubMed: 20807578]
- Desseilles M, Baletau E, Sterpenich V, et al. Abnormal neural filtering of irrelevant visual information in depression. *Journal of Neuroscience*. 2009; 29:1395–403. [PubMed: 19193886]
- Dillon DG, Rosso IM, Pechtel P, et al. Peril and pleasure: an rdoc-inspired examination of threat responses and reward processing in anxiety and depression. *Depression and Anxiety*. 2014; 31:233–49. [PubMed: 24151118]
- Doehrmann O, Ghosh SS, Polli FE, et al. Predicting treatment response in social anxiety disorder from functional magnetic resonance imaging. *JAMA Psychiatry*. 2013; 70:87–97. [PubMed: 22945462]
- Etkin A, Schatzberg AF. Common abnormalities and disorder-specific compensation during implicit regulation of emotional processing in generalized anxiety and major depressive disorders. *American Journal of Psychiatry*. 2011; 168(9):968–78. [PubMed: 21632648]
- Eysenck, M. *Anxiety: The Cognitive Perspective*. Hove: Erlbaum; 1992.
- Eysenck M, Derakshan N, Santos R, Calvo M. Anxiety and cognitive performance: attentional control theory. *Emotion*. 2007; 7:336–353. [PubMed: 17516812]
- Fani N, Tone EB, Phifer J, et al. Attention bias toward threat is associated with exaggerated fear expression and impaired extinction in PTSD. *Psychological Medicine*. 2012; 42:533–43. [PubMed: 21854700]
- Fava M, Rush AJ, Alpert JE, et al. Difference in treatment outcome in outpatients with anxious versus nonanxious depression: a STAR*D report. *American Journal of Psychiatry*. 2008; 165:342–51. [PubMed: 18172020]
- First, MB., Spitzer, RL., Gibbon, M., Williams, JBW. *Structured Clinical Interview for DSM-IV Axis I Disorders, Clinician Version (SCID-CV)*. Washington, D.C: American Psychiatric Association; 1996.
- Fu CH, Williams SC, Cleare AJ, et al. Neural responses to sad facial expressions in major depression following cognitive behavioral therapy. *Biological Psychiatry*. 2008; 64:505–12. [PubMed: 18550030]
- Gibb BE, McGeary JE, Beevers CG. Attentional biases to emotional stimuli: Key components of the RDoC constructs of sustained threat and loss. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*. 2015; 171:65–80.
- Gyurak A, Patenaude B, Korgaonkar MS, Grieve SM, Williams LM, Etkin A. Frontoparietal activation during response inhibition predicts remission to antidepressants in patients with major depression. *Biological Psychiatry*. 2016; 79:274–81. [PubMed: 25891220]

- Hajcak G, MacNamara A, Olvet DM. Event-related potentials, emotion, and emotion regulation: an integrative review. *Developmental Neuropsychology*. 2010; 35:129–55. [PubMed: 20390599]
- Hajcak G, Dunning JP, Foti D. Neural response to emotional pictures is unaffected by concurrent task difficulty: an event-related potential study. *Behavioral Neuroscience*. 2007; 121:1156–62.
- Hajcak, G., Weinberg, A., MacNamara, A., Foti, D. ERPs and the study of emotion. In: Luck, SJ., Kappenman, ES., editors. *Oxford handbook of ERP components*. New York: Oxford University Press; 2012. p. 441–474.
- Hamilton M. A rating scale for depression. *Journal of Neurology, Neurosurgery, and Psychiatry*. 1960; 23:56–62.
- Hamilton M. The assessment of anxiety states by rating. *British Journal of Medical Psychology*. 1959; 32:50–5. [PubMed: 13638508]
- Hofmann SG, Asnaani A, Vonk IJ, et al. The efficacy of cognitive behavioral therapy: a review of meta-analyses. *Cognitive Therapy and Research*. 2012; 36:427–440. [PubMed: 23459093]
- Hofmann SG, Smits JA. Cognitive-behavioral therapy for adult anxiety disorders: a meta-analysis of randomized placebo-controlled trials. *Journal of Clinical Psychiatry*. 2008; 69:621–32. [PubMed: 18363421]
- Hope, DA., Heimberg, RG., Turk, CL. *Managing social anxiety: a cognitive-behavioral therapy approach*. New York: Oxford University Press; 2006.
- Jaycox LH, Foa EB, Morral AR. Influence of emotional engagement and habituation on exposure therapy for PTSD. *Journal of Consulting and Clinical Psychology*. 1998; 66:185–92. [PubMed: 9489273]
- Johnco C, Wuthrich VM, Rapee RM. The influence of cognitive flexibility on treatment outcome and cognitive restructuring skill acquisition during cognitive behavioural treatment for anxiety and depression in older adults: Results of a pilot study. *Behaviour Research and Therapy*. 2014; 57:55–64. [PubMed: 24828838]
- Kashdan TB, Barrios V, Forsyth JP, Steger MF. Experiential avoidance as a generalized psychological vulnerability: comparisons with coping and emotion regulation strategies. *Behaviour Research and Therapy*. 2006; 44:1301–20. [PubMed: 16321362]
- Kaufman J, Charney D. Comorbidity of mood and anxiety disorders. *Depression and Anxiety*. 2000; 12(1):69–76. [PubMed: 11098417]
- Kemp AH, Gordon E, Rush AJ, Williams LM. Improving the prediction of treatment response in depression: integration of clinical, cognitive, psychophysiological, neuroimaging, and genetic measures. *CNS Spectrums*. 2008; 13:1066–1088. [PubMed: 19179943]
- Kessler RC, Akiskal HS, Ames M, et al. Prevalence and effects of mood disorders on work performance in a nationally representative sample of U.S. workers. *American Journal of Psychiatry*. 2006; 163:1561–8. [PubMed: 16946181]
- Kessler RC, Berglund P, Demler O, et al. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *JAMA*. 2003; 289(23):3095–3105. [PubMed: 12813115]
- Kessler RC, Chiu WT, Demler O, et al. Prevalence, severity, and comorbidity of 12-month DSM-IV disorders in the National Comorbidity Survey Replication. *Archives of General Psychiatry*. 2005; 62:617–27. [PubMed: 15939839]
- Kircanski K, Joormann J, Gotlib IH. Attention to emotional information in social anxiety disorder with and without co-occurring depression. *Cognitive Therapy and Research*. 2015; 39:153–61.
- Klump H, Fitzgerald DA, Phan KL. Neural predictors and mechanisms of cognitive behavioral therapy on threat processing in social anxiety disorder. *Progress in Neuropsychopharmacology and Biological Psychiatry*. 2013; 45:83–91.
- Klump H, Keutmann MK, Fitzgerald DA, et al. Resting state amygdala-prefrontal connectivity predicts symptom change after cognitive behavioral therapy in generalized social anxiety disorder. *Biology of Mood and Anxiety Disorders*. 2014; 4:14. [PubMed: 25540682]
- Kujawa A, Swain JE, Hanna GL, et al. Prefrontal reactivity to social signals of threat as a predictor of treatment response in anxious youth. *Neuropsychopharmacology*. 2016

- Lang, P., Bradley, M., Cuthbert, B. International affective picture system (IAPS): affective ratings of pictures and instruction manual Technical Report A-6. University of Florida; Gainesville, FL: 2005.
- Leutgeb V, Schäfer A, Schienle A. An event-related potential study on exposure therapy for patients suffering from spider phobia. *Biological Psychology*. 2009; 82:293–300. [PubMed: 19751797]
- Li W, Zinbarg RE, Paller KA. Trait anxiety modulates supraliminal and subliminal threat: brain potential evidence for early and late processing influences. *Cognitive, Affective, and Behavioral Neuroscience*. 2007; 7:25–36.
- Liu Y, Huang H, McGinnis-Deweese M, et al. Neural substrate of the late positive potential in emotional processing. *Journal of Neuroscience*. 2012; 32:14563–72. [PubMed: 23077042]
- MacNamara A, Rabinack CA, Kennedy AE, et al. Emotion Regulatory Brain Function and SSRI Treatment in PTSD: Neural Correlates and Predictors of Change. *Neuropsychopharmacology*. 2015; 41:611–618. DOI: 10.1038/npp.2015.190 [PubMed: 26111649]
- MacNamara A, Ferri J, Hajcak G. Working memory load reduces the late positive potential and this effect is attenuated with increasing anxiety. *Cognitive, Affective, and Behavioral Neuroscience*. 2011; 11:321–31.
- MacNamara A, Hajcak G. Anxiety and spatial attention moderate the electrocortical response to aversive pictures. *Neuropsychologia*. 2009; 47:2975–80. [PubMed: 19576234]
- MacNamara A, Hajcak G. Distinct electrocortical and behavioral evidence for increased attention to threat in generalized anxiety disorder. *Depression and Anxiety*. 2010; 27:234–43. [PubMed: 20196100]
- MacNamara, A., Kappenman, ES., Black, SR., et al. Integrating behavioral and electrocortical measures of attentional bias toward threat. In: Barrett, KC.Fox, NA.GA, MorganFidler, DJ., Daunhauer, LA., editors. *Handbook of self-regulatory processes in development: New directions and international perspectives*. New York: Psychology Press; 2013. p. 215–246.
- MacNamara A, Proudfit GH. Cognitive load and emotional processing in generalized anxiety disorder: electrocortical evidence for increased distractibility. *Journal of Abnormal Psychology*. 2014; 123:557–65. [PubMed: 24933276]
- MacNamara A, Proudfit GH. Diagnostic and symptom-based predictors of emotion processing in generalized anxiety disorder and major depressive disorder: An event-related potential study. *Cognitive Therapy and Research*. in press.
- Martell, CR., Dimidjian, S., Herman-Dunn, R. *Behavioral activation for depression*. New York: Guilford Press; 2010.
- Mathews A, MacLeod C. Cognitive vulnerability to emotional disorders. *Annual Review of Clinical Psychology*. 2005; 1:167–95.
- Mathews A, MacLeod C. Induced processing biases have causal effects on anxiety. *Cognition and Emotion*. 2002; 16:331–354.
- Miller GA, Gratton G, Yee CM. Generalized implementation of an eye movement correction procedure. *Psychophysiology*. 1988; 25:241–243.
- Mineka S, Watson D, Clark LA. Comorbidity of anxiety and unipolar mood disorders. *Annual Review of Clinical Psychology*. 1998; 49:377–412.
- Moras K, Di Nardo PA, Barlow DH. Distinguishing anxiety and depression: Reexamination of the reconstructed Hamilton scales. *Psychological Assessment*. 1992; 4:224–227.
- Peckham AD, McHugh RK, Otto MW. A meta-analysis of the magnitude of biased attention in depression. *Depression and Anxiety*. 2010; 27:1135–42. [PubMed: 21049527]
- Pergamin-Hight L, Naim R, Bakermans-Kranenburg MJ, et al. Content specificity of attention bias to threat in anxiety disorders: a meta-analysis. *Clinical Psychology Review*. 2015; 35:10–18. [PubMed: 25462110]
- Pessoa L, Kastner S, Ungerleider LG. Attentional control of the processing of neutral and emotional stimuli. *Cognitive Brain Research*. 2002; 15:31–45. [PubMed: 12433381]
- Phan KL, Coccaro EF, Angstadt M, et al. Corticolimbic brain reactivity to social signals of threat before and after sertraline treatment in generalized social phobia. *Biological Psychiatry*. 2013; 73(4):329–336. [PubMed: 23164370]

- Pishyar R, Harris LM, Menzies RG. Responsiveness of measures of attentional bias to clinical change in social phobia. *Cognition and Emotion*. 2008; 22:1209–27.
- Proudfit, GH., Dunning, J., Foti, D., Weinberg, A. Temporal dynamics of emotion regulation. In: Gross, JJ., editor. *Handbook of emotion regulation*. New York: Guilford; 2013. p. 43–57.
- Proudfit GH, Bress JN, Foti D, et al. Depression and event-related potentials: Emotional disengagement and reward insensitivity. *Current Opinion in Psychology*. 2015; 4:110–113. [PubMed: 26462292]
- Ritchey M, Dolcos F, Eddington KM, et al. Neural correlates of emotional processing in depression: changes with cognitive behavioral therapy and predictors of treatment response. *Journal of Psychiatric Research*. 2011; 45:577–87. [PubMed: 20934190]
- Rodebaugh TL, Holaway RM, Heimberg RG. The treatment of social anxiety disorder. *Clinical Psychology Review*. 2004; 24(7):883–908. [PubMed: 15501560]
- Sabatinielli D, Lang PJ, Keil A, Bradley MM. Emotional perception: Correlation of functional MRI and event-related potentials. *Cerebral Cortex*. 2007; 17:1085–91. [PubMed: 16769742]
- Sanislow CA, Pine DS, Quinn KJ, et al. Developing constructs for psychopathology research: research domain criteria. *Journal of Abnormal Psychology*. 2010; 119:631–9. [PubMed: 20939653]
- Schneibel R, Brakemeier EL, Wilbertz G, et al. Sensitivity to detect change and the correlation of clinical factors with the Hamilton Depression Rating Scale and the Beck Depression Inventory in depressed inpatients. *Psychiatry Research*. 2012; 198:62–67. [PubMed: 22445070]
- Shear MK, Vander Bilt J, Rucci P, et al. Reliability and validity of a structured interview guide for the Hamilton Anxiety Rating Scale (SIGH-A). *Depression and Anxiety*. 2001; 13:166–178. [PubMed: 11413563]
- Siegle GJ, Carter CS, Thase ME. Use of fMRI to predict recovery from unipolar depression with cognitive behavior therapy. *American Journal of Psychiatry*. 2006; 163:735–8. [PubMed: 16585452]
- Simpson HB. The RDoC project: a new paradigm for investigating the pathophysiology of anxiety. *Depression and Anxiety*. 29:251–2.
- Stange JP, Connolly SL, Burke TA, Hamilton JL, Hamlat EJ, Abramson LY, Alloy LB. Inflexible cognition predicts first onset of major depressive episodes in adolescence. *Depression and Anxiety*. 2016; doi: 10.1002/da.22513
- Stein MB, Fuetsch M, Müller N, et al. Social anxiety disorder and the risk of depression: a prospective community study of adolescents and young adults. *Archives of General Psychiatry*. 2001; 58(3): 251–256. [PubMed: 11231832]
- Streb M, Mecklinger A, Anderson MC, Johanna LH, Michael T. Memory control ability modulates intrusive memories after analogue trauma. *Journal of Affective Disorders*. 2016; 192:134–142. [PubMed: 26724692]
- Tacikowski P, Nowicka A. Allocation of attention to self-name and self-face: An ERP study. *Biological Psychology*. 2010; 84:318–24. [PubMed: 20298741]
- Tillfors M, Furmark T, Carlbring P, Andersson G. Risk profiles for poor treatment response to internet-delivered CBT in people with social anxiety disorder. *Journal of Anxiety Disorders*. 2015; 33:103–9. [PubMed: 26087474]
- Tracy JL, Klonsky ED, Proudfit GH. How affective science can inform clinical science: An introduction to the special series on emotions and psychopathology. *Clinical Psychological Science*. 2014; 2:371–86.
- Waters AM, Kershaw R. Direction of attention bias to threat relates to differences in fear acquisition and extinction in anxious children. *Behaviour Research and Therapy*. 2015; 64:56–65. [PubMed: 25540863]
- Weinberg A, Hajcak G. Beyond good and evil: the time-course of neural activity elicited by specific picture content. *Emotion*. 2010; 10:767–82. [PubMed: 21058848]
- Weinberg A, Hajcak G. Electrocardiac evidence for vigilance-avoidance in Generalized Anxiety Disorder. *Psychophysiology*. 2011; 48:842–51. [PubMed: 21073479]
- Weinberg A, Perlman G, Kotov R, Hajcak G. Depression and reduced neural response to emotional images: Distinction from anxiety, and importance of symptom dimensions and age of onset. *Journal of Abnormal Psychology*. 2016; 125:26–39. [PubMed: 26726817]

- Whalen PJ, Johnstone T, Somerville LH, et al. A functional magnetic resonance imaging predictor of treatment response to venlafaxine in generalized anxiety disorder. *Biological Psychiatry*. 2008; 63:858–63. [PubMed: 17964548]
- Whelton WJ. Emotional processes in psychotherapy: Evidence across therapeutic modalities. *Clinical Psychology and Psychotherapy*. 2004; 11:58–71.

Highlights

- We evaluated attention toward aversive and neutral target and distracter stimuli.
- Event-related potentials (the late positive potential; LPP) were recorded pre-treatment.
- Patients received up to 12 sessions of CBT for social anxiety or major depression.
- Larger LPPs to aversive distracters (when targets were aversive) predicted treatment response.
- CBT may be most beneficial for individuals with attenuated top-down control of attention.

Figure 1. Grand average amplitudes at pooling of CP1, CP2, Cz, and Pz (panel A) and scalp distributions of amplitudes from 400-1000 ms after picture onset (panel B) for each trial type.

Figure 2.

Grand-average amplitudes for each trial type and spatial distributions of amplitude differences (from 400-1000ms after picture onset) for aversive minus neutral distracters under conditions of aversive targets, shown separately for Non-Responders (CGI-Severity and CGI-Improvement > 3; panel A) and Responders (CGI-Severity and CGI-Improvement < 3; panel B).

Figure 3.

Scatterplots of associations between pre-treatment LPP (difference between aversive and neutral distracters on aversive target trials) and post-treatment residual Hamilton Depression Rating Scale scores (HAM-D; controlling for pre-treatment HAM-D; panel A) and post-treatment Hamilton Anxiety Rating Scale scores (HAM-A; controlling for pre-treatment HAM-A; panel B), plotted by primary diagnosis (SAD = social anxiety disorder; MDD = major depressive disorder).

Table 1

Sample characteristics and diagnoses.

	Mean	SD
Age (years)	24.03	5.38
Education (years)	15.22	2.20
	N	%
Female	27	84.4
Race		
Caucasian	19	59.4
African American	1	3.1
Asian	8	25.0
More than one race	4	12.5
Hispanic or Latino/a	7	21.9
Primary Diagnosis		
Social Anxiety Disorder	20	62.5
Major Depressive Disorder	12	37.5
Any Current Diagnosis		
Social Anxiety Disorder	2	81.3
Major Depressive Disorder	13	40.6
Generalized Anxiety Disorder	9	28.1
Panic Disorder	8	25.0
Specific Phobia	3	9.4
Post-Traumatic Stress Disorder	4	12.5

Note. $N = 32$. CGI = Clinical Global Impression scale; HAM-D = Hamilton Depression Rating Scale; HAM-A = Hamilton Anxiety Rating Scale.

Table 2

Treatment outcome scores according to CGI responder status.

	Responder		Non-Responder	
	Mean	SD	Mean	SD
CGI Severity (pre-treatment)	4.00	0.00	4.36	0.57
CGI Severity (post-treatment)	1.64	0.50	3.57	0.75
CGI Improvement (post-treatment)	1.00	0.00	2.67	0.97
HAM-A (pre-treatment)	13.73	9.13	17.10	7.64
HAM-A (post-treatment)	2.82	2.82	8.67	5.08
HAM-D (pre-treatment)	8.55	6.70	11.00	6.82
HAM-D (post-treatment)	1.64	1.36	5.57	4.74

Note. $N = 32$. CGI = Clinical Global Impression scale; HAM-D = Hamilton Depression Rating Scale; HAM-A = Hamilton Anxiety Rating Scale.