Published in final edited form as:

Chemistry. 2016 December 19; 22(51): 18290–18315. doi:10.1002/chem.201603382.

Cyclobutenones and Benzocyclobutenones: Versatile Synthons in Organic Synthesis

Peng-hao Chen^[a] and Prof. Dr. Guangbin Dong^{[a],[b]}

^[a]Department of Chemistry University of Texas at Austin, 100 East 24th street, Austin, TX, 78712 (USA), <http://gbdong.cm.utexas.edu/>

^[b]Department of Chemistry University of Chicago, Chicago, Illinois 60637 (USA)

Abstract

Cyclobutenones, four-membered ketones bearing an unsaturated carbon–carbon double bond, and their structural sibling benzocyclobutenones, possess unique reactivity. Owing to their inherent high ring strain, such structures readily undergo ring opening under a variety of conditions, including thermolysis, photolysis, and transition metal catalysis, to afford reactive intermediates that can be trapped with nucleophiles, dienophiles, and unsaturated bonds. Their electron-deficient enone moieties are good electrophiles for facile nucleophilic addition. Such properties render cyclobutenones versatile synthons, serving as excellent coupling partners in a vast array of synthetically valuable transformations.

Graphical Abstract

Correspondence to: Guangbin Dong.

The ORCID identification number(s) for the author(s) of this article can be found under <http://dx.doi.org/10.1002/chem.201603382>.

Keywords

cycloaddition; electrocyclic reactions; enones; homogeneous catalysis; strained molecules

1. Introduction

Recent years have witnessed a dramatic increase in the use of four-membered ring compounds as building blocks for the rapid construction of complex molecules.^[1] Owing to their inherent ring strain, such small ring structures can undergo ring-opening reactions triggered by myriad reaction conditions, allowing them to serve as coupling partners and core structures in a variety of transformations. Of the four-membered ring compounds, the properties of structures with saturated carbon–carbon bonds have been extensively reviewed, but four-membered ring structures bearing unsaturated carbon–carbon bonds, namely cyclobutenones and benzocyclobutenones, have received less attention.^[2] Cyclobutenones and benzocyclobutenones can be conveniently prepared through [2+2] cycloadditions between alkynes or benzyne and ketenes or their equivalents,^[2] although alternative methods, such as intramolecular nucleophilic addition,^[3a] 4 π -electrocyclic ring closure,^[3b] and intramolecular transition metal-catalyzed cross-couplings,^[3c–e] are also known. This Minireview focuses on representative transformations of cyclobutenones and benzocyclobutenones; examples of cyclobutenols, cyclobutediones, and squaric acid derivatives, as well as preparation of cyclobutenones and benzocyclobutenones,^[1–3] will not be covered.

As an overview of the reactions of cyclobutenones and benzocyclobutenones (Scheme 1), early reports primarily involved photolysis and thermolysis, in which vinylketene is generally the reactive intermediate. Thus, cyclobutenones and benzocyclobutenones have been widely incorporated in transformations including [4+2]/[2+2] cycloadditions. In addition, given the nature of the conjugate enone moiety, both 1,2- and 1,4-nucleophilic additions have been investigated. Moreover, compared to cyclopentenones and cyclohexenones, cyclobutenones were recently found to be exceptional dienophiles for Diels–Alder reactions as such cycloadditions benefit from the release of ring strain. Finally, cyclobutenones and benzocyclobutenones have been frequently employed as substrates in C–C-activation reactions,^[4] in which transition metals, including nickel, ruthenium, rhodium and cobalt, can selectively activate either C1–C2 or C1–C4 carbon–carbon bonds.

2. Thermal Ring Opening

The electrocyclic reactions of cyclobutenes have been extensively studied and the stereoselectivity can be readily predicted. Terms like “conrotatory” and “disrotatory” are well defined and widely used to describe such transformations.^[5] However, these terms are inadequate for describing the electrocyclic ring opening transformation of cyclobutenones and benzocyclobutenones, as there is only one carbon center carrying sp³ hybridization. In the ring opening of cyclobutenones, the terms “inward” and “outward” are frequently employed to describe the rotation: if a substituent rotates “inward”, it becomes *cis* to the ketene moiety, whereas it rotates “outward” to be *trans* to the ketene moiety (Scheme 2).^[5]

Houk and coworkers explored the effect of C4 substituents on the rotation barrier under thermal conditions and found that, although the substituent effects are less pronounced compared to cyclobutenes, electron donating groups prefer to rotate “outward”, whereas electron-withdrawing groups tend to rotate “inward”.^[6] Such results can be attributed to the fact that, when rotating “outward”, the high-lying electron-donor orbital of the substituent can mix with the low lying σ^* (LUMO) of the breaking bond, thus stabilizing the intermediate and lowering the activation energy.

In 1956, Jenny and Roberts studied the racemization of optically active 2,4-dichloro-3-phenylcyclobutenone **1** (Scheme 3).^[7] They rationalized that a slowly formed vinylketene intermediate, which underwent rapid cyclization to give back the cyclobutenone, was responsible for the racemization. The putative vinylketene intermediate could either undergo rapid ring closure to furnish racemic **1** or react with acetic acid or acetate to give mixed anhydride **2**.

In 1975, Mayr reported that 2,4,4-trimethylcyclobutenone **3** also underwent facile ring opening in boiling cyclohexane, and the vinylketene intermediate was trapped in situ by ethyl vinyl ether to afford cyclobutanone **4** (Scheme 4).^[8] Huisgen and Mayr later reported that the vinylketene intermediate formed from cyclobutenone **5** can be intercepted by nucleophiles such as methanol and aniline, to give ester **6** or amide **7**.^[9]

In 1977, Ficini et al. reported an unexpected rearrangement during the cycloaddition of ynamines with cyclobutenones (Scheme 5).^[10] Unlike the cycloaddition of ynamines with cyclopentenones or cyclohexenones,^[11] the reaction of ynamines with strained cyclobutenone **8** did not afford the [2+2] product **9**. Instead, 3-amino-4-vinylcyclobutenones **10** were obtained in moderate yields. It was proposed that **10** came from intermediate **9**, through a concerted rearrangement involving one π bond and two σ bonds, and at low temperature, the disrotatory ring-opening of **9** to form phenols is thermally forbidden. However, upon heating, **10** was converted to polysubstituted phenols **11**. This transformation was proposed to involve thermal ring opening of the cyclobutenone followed by ring closure and aromatization.

Eight-membered rings can also be accessed from cyclobutenones. In 1982, the Danheiser group developed a formal [4+4] annulation process to synthesize unsymmetrical highly functionalized cyclooctadienones (Scheme 6).^[12] This reaction employed 1,3-dienes and vinylketenes as coupling partners, with cyclobutenones serving as the vinylketene precursor. Under the reaction conditions, the in situ generated vinylketene underwent a facile [2+2] cycloaddition to afford 2,3-divinylcyclobutanone **12**, which upon heating gave cyclooctadienone **13** through a [3,3]-sigmatropic rearrangement.

The Danheiser group in 1984 reported an efficient approach to access highly substituted aromatic compounds from cyclobutenones and heteroatom (O, N or S)-substituted alkynes **14** via four successive pericyclic reactions (Scheme 7).^[13] Upon heating above 80 °C, 4 π -electrocyclic ring opening of cyclobutenone reversibly generated a vinylketene intermediate **15**, which subsequently reacted with the electron-rich alkyne in a regiospecific [2+2] cycloaddition to give 4-vinyl cyclobutenones **16**. As reported in Ficini's work,^[10] these 4-

vinyl cyclobutenones can undergo a retro-4 π /6 π cyclization cascade to furnish highly substituted phenol derivatives.

Later, Danheiser et al. also successfully applied this benzannulation method to the total synthesis of Mycophenolic Acid **19** (Scheme 8).^[14] The key step in the synthesis involves the construction of pentasubstituted resorcinol by a cycloaddition cascade between cyclobutenone **17** and alkynyl ether **18**.

Although methoxy-substituted alkynes are excellent coupling partners for the benzannulation reaction with cyclobutenones, removal of the methyl group to generate free phenols is not trivial. In 1988, the groups of Kowalski^[15] and Danheiser^[16] independently developed an extended protocol using alkynyl silyl ethers **20** as substrates (Scheme 9). Under thermal conditions, the annulation proceeded with high yields to afford a variety of monosilyl-protected resorcinols. The silyl group can be smoothly removed under acidic conditions to give free phenols. By using this method, the Kowalski group accomplished an elegant synthesis of Δ^9 -6-tetrahydrocannabinol **21**. In both reports, 2,4-disubstituted cyclobutenones failed to furnish the desired annulation products.

Although alkynyl ethers have been demonstrated as good coupling partners in benzannulation with cyclobutenones, the use of ynamines for phenol synthesis has been less common.^[10] In Danheiser's cyclobutenone benzannulation with *N,N*-dimethyl ynamines (Scheme 10),^[13] allenes **22** were observed as byproducts. The formation of such allene products were suggested to have resulted from a zwitterionic intermediate **23**.^[17] This intermediate can either cyclize to provide the desired phenol product or afford allene **22** via an alkylideneoxete intermediate **24**.

Although the zwitterionic intermediate was able to give phenol products, the regioselectivity would be different from the electrocyclic reaction pathway (e.g., **25a** and **25b**), as depicted in Scheme 11. Considering the unusual rearrangement reported by Ficini et al. (e.g., **25c**),^[10] the benzannulation of vinylketenes from cyclobutenones could provide three different isomers.

To suppress these alternative regioisomeric products, less nucleophilic ynamides were employed in the benzannulation (Scheme 12).^[18] Compared to ynamines, the ynamides favor the desired concerted [2+2] cycloaddition with the ketene rather than carbonyl addition. Under thermal conditions, the benzannulation between cyclobutenones **26** and ynamides **27** proceeded with high regioselectivity. By using this method, Danheiser and co-workers accomplished the formal synthesis of (+)-FR900482 (**28**, Scheme 13).

In addition to the intermolecular coupling between cyclobutenones and unsaturated π bonds, intramolecular cyclizations by electrocyclic ring opening of cyclobutenones have also been developed. In 1975, Mayr reported that cyclobutenones **29** were converted into the corresponding α -naphthols **30** quantitatively in refluxing cyclohexane (Scheme 14).^[8] For cyclobutenone **31**, bearing two different substituents at the C4 position, a mixture of 49 % of α -naphthol **32**, 21 % of **33** and 30 % of **34** were isolated, indicating the formation of both *cis*- and *trans*-vinylketenes.

In 1985, Moore and co-workers developed the synthesis of polysubstituted benzoquinones through thermal rearrangement of 4-alkynylcyclobutenols **35** prepared from squaric acid derivatives (Scheme 15).^[19] Upon 4 π -electrocyclic ring opening, the resulting vinylketene intermediate **36** underwent ring closure to afford zwitterions **37** and/or **38**, eventually leading to products **39** and/or **40**. The selectivity was largely affected by the substituent: strong electron-withdrawing groups favored the formation of cyclopentenediones, whereas electron-donating groups facilitated the formation of benzoquinones.

Interestingly, an allyl migration occurred when the trimethylsilyl group was replaced with an allyl group (e.g., **41**).^[20] By using aryl-substituted substrates (**42**, Scheme 16), aryl-fused benzoquinones were formed upon concomitant oxidation with Ag₂O or Ce^{IV}. In all cases, the vinylketene intermediates were generated by an outward ring opening with regards to the oxygen substituent.

In 1988, Moore and Perri reported a detailed study on the substituent effect of the aryl group in the benzannulation reaction of 4-arylcyclobutenols (Scheme 17).^[21] When cyclobutenones **43** and **44** were subjected to the reaction conditions, the same naphthalene-fused benzoquinone product was observed. Given that the 1-position of naphthalene is more reactive towards electrophilic substitution, this result indicated that some electrophilic character was required in the transition state of this reaction. In addition, the electrocyclic ring opening of cyclobutenone **45** was expected to give phenol **46**. However, a mixture of quinones **47** and **48** was isolated instead, in which the ring closure of the vinylketene intermediate had selectively occurred at the carbon bearing the methoxy group followed by concomitant tautomerization or elimination of methanol.

Moreover, although in most cases the hydroxy group favors outward rotation during ring opening, the presence of the strongly electron-rich aryl group (e.g., substrate **49**) favors inward rotation, affording vinylketene intermediate **50**, which led to the formation of γ -lactones **51** and **52** after workup with silica gel (Scheme 18).

In 1992, Liebeskind and co-workers developed a one-pot method for α -vinylation or arylation followed by benzannulation from 4-chlorocyclobutenone **53** (Scheme 19).^[22] Through a Pd-catalyzed Stille coupling, a range of vinyl- and aryltin compounds were coupled with **53** to generate reactive cyclobutenone intermediate **54** in situ; subsequent thermal ring-opening and 6 π -cyclization furnished the corresponding phenols **55** in high yields. Interestingly, when vinylzirconium **56** was employed, upon thermolysis, the reaction selectively afforded one of the two regioisomers **57** or **58**, depending on the choice of catalyst system. For example, with 10 mol % [$\text{Pd}(\text{allyl})\text{Cl}$]₂ and 20 mol % PPh₃, phenol **57** was produced, whereas with 10 mol % $[\text{Pd}(\text{PPh}_3)_4]$ and 40 mol % of PPh₃, phenol **58** was produced. However, the reason for this selectivity is not yet clear.

Later, syntheses of 1,4-quinone derivatives by thermal ring opening of cyclobutenones were extensively studied with substituents including a range of vinyl,^[23,24] aryl, and heteroaryl^[25] groups. However, in most cases, the formation of quinones required an oxidation step (Scheme 20).

Aiming to avoid the oxidative modification, Harrowven and co-workers introduced a leaving group to the olefin moiety.^[26] Under microwave conditions, cyclobutenone **59** afforded the desired quinone **60** in the absence of any oxidant (Scheme 21). However, an unusual phenomenon was observed: on increasing the electron density of the incorporated olefin moiety, carbocyclic compound **61**, putatively through a carbonyl-ene reaction, became the major product. Interestingly, when 4-(*o*-styryl)cyclobutenones (e.g., **62**, Scheme 22) were subjected to the reaction conditions with elevated temperature, benzobicyclo[3.2.1] structures **63** were afforded in high yields, which was proposed to involve a diradical intermediate **64**.

Moore and co-workers later expanded this transformation to cyclobutenones bearing 4-allyl substituents (Scheme 23).^[27] Upon 4 π -electrocyclic ring opening, the resulting vinylketene intermediate **65** underwent a facile [2+2] cycloaddition to furnish the bicyclo[3.2.0]heptenones **66** as the final products. Substrates bearing free 4-OH (R=H), although they successfully underwent a similar ring expansion (e.g., **66g**), were rather limited in substrate scope as a 2-phenyl substituent would lead to an unusual product **66h**. Substrate **66e**, which undergoes facile deprotection to afford diketone **67**, was used to unveil the mechanism of the reaction with **66h**. Although it is stable under thermolysis in toluene, **67** was converted into tricycle **66h** in the presence of a catalytic amount of acetic acid.

The vinylketene intermediate can also be intercepted intra-molecularly by a σ -nucleophile. In 2009, Shimizu and co-workers developed the synthesis of β -lactams through thermal ring opening of aminocyclobutenone **68**.^[28] Upon 4 π -electrocyclic ring opening, the resulting amino vinylketene intermediate was attacked by the amine moiety to provide the β -lactams **69** in good yields. Although inherently favoring the *cis* isomer (3:1–4:1 without adding bases), the stereoselectivity can be well controlled: when 1,8-diazabicyclo[5.4.0]undec-7-ene (DBU) was added, *trans*-**69** was favored; in contrast, in the presence of 1,4-dimethylpiperazine, *cis*-**69** was afforded exclusively. In 2014, the same group developed the enantioselective version of this transformation by using as the starting materials chiral aminocyclobutenones **68** (Scheme 24),^[29] which were obtained from asymmetric reduction of the corresponding iminocyclobutenones **70** by using chiral phosphoric acid **71** as catalyst and dihydrobenzothiazole **72** as the reductant. Upon thermolysis with different bases, both *trans*- and *cis*- β -lactams were afforded in high yield and enantioselectivity.

3. Photolysis-Induced Ring Opening

In 1967, Baldwin and co-workers studied the ring opening of 2,4-dichloro-3-phenylcyclobutenone **73** (Scheme 25).^[30,31] Under thermal conditions, *E*- β,γ -unsaturated ester **74** was provided; in contrast, under irradiation by a high-pressure mercury lamp, *Z*- β,γ -unsaturated ester **75** was afforded. The stereoselectivity was tentatively explained by the calculated excited state of the reaction. Based on Kikuchi's work,^[32] the $n-\pi^*$ singlet excited state could lead to a perpendicular diradical, which caused an inward rotation.

The involvement of a vinylketene intermediate under photolysis conditions was observed by Chapman and Lassila by using infrared spectroscopy.^[33] Upon irradiation with a high-

pressure mercury lamp, 2,3,4,4-tetrachlorocyclobutenone **76** (Scheme 26) afforded a new carbonyl band at 2145 cm^{-1} , which was believed to be a ketene carbonyl peak of **77**.

In 1974, Toda and Todo reported the photochemical ring-opening of methylenecyclobutenone **78** (Scheme 27).^[34] Under UV irradiation, the ring opening of **78** afforded allene–ketene intermediate **79**. Similar to other vinylketene intermediates, **79** can be intercepted intramolecularly by a hydroxy group, affording furanone **80** as the final product. With methanol as solvent, however, photolysis provided isomeric compound **81** as the product.^[35] Control experiments showed that, although there was no conversion under the photolysis conditions, **80** could isomerize to **81** rapidly upon heating in the presence of a base.

Later, the same group demonstrated that, upon ring-opening, the allene–ketene intermediate **82** could undergo a rapid [2+2] cycloaddition with benzophenone to afford an oxetanone intermediate, which eventually led to product **83** after a concomitant elimination of diphenylketene **84** (Scheme 28).^[36] A similar transformation was realized under an oxygen atmosphere with elimination of carbon dioxide, albeit with a lower yield.

In 1984, Schiess et al. reported the cycloaddition of benzocyclobutenone **85** (Scheme 29) with various dienophiles under flash photolysis conditions.^[37] Upon UV irradiation in ether, benzocyclobutenone **85** underwent isomerization to vinylketene **86** with a lifetime of 170 ms at room temperature. Unlike simple vinylketenes, which typically react with electron-rich olefins, **86** acted as a diene and rapidly formed [4+2] cycloaddition products with diazo, activated carbonyl compounds or Michael acceptors, likely driven by forming aromatic ketones.

4. Nucleophile-Induced Ring Opening

Due to the α,β -unsaturated ketone moiety, cyclobutenones can be attacked by nucleophiles in a 1,2- or 1,4-addition fashion. The resulting alkoxide or enolate often triggers the subsequent ring opening of the four membered ring structures. In the 1950s, Roberts and co-workers studied the reaction of 3-phenyl-4,4-dichlorocyclobutenone **87** with hydroxide (Scheme 30).^[38] 1,2-Addition of hydroxide to the carbonyl moiety in **87** initiated ring opening through retro- 4π -cyclization, giving 4,4-dichloro-3-phenyl-2-butenic acid **88** as a final product in good yield. A similar transformation in the presence of *n*BuOH was later reported by Hassner and Dillon, which eventually led to the formation of the corresponding ester.^[39]

In 2001, Murakami et al. reported the ring opening of cyclobutenones triggered by silyl anions (Scheme 31).^[40] When 3-phenylcyclobutenone **89** was treated with a silylcuprate at $-78\text{ }^{\circ}\text{C}$, the 1,4-addition product could be smoothly trapped by acetic anhydride to afford vinyl acetate **90**, which, upon heating, afforded diene product **91** in an excellent yield as one stereoisomer. When silyllithium was used as a nucleophile, **89** underwent 1,2-addition to afford highly reactive intermediate **92**, which, upon treatment with acetic anhydride at $-78\text{ }^{\circ}\text{C}$, gave the ring-opening product **93** in excellent yield. Based on these results, the same

group later developed an efficient approach for the stereoselective synthesis of highly functionalized 1,3-dienes from cyclobutenones.^[41]

In 2004, Magomedov et al. reported a cascade reaction sequence for the synthesis of heavily functionalized cyclohexanones through the ring opening of cyclobutenone **94**, which was initiated by addition of vinyl lithium reagent **98** (Scheme 32).^[42] Upon 1,2-addition, alkoxide **95** underwent a charge-accelerated 4π conrotatory ring opening to generate triene **96**, which, through 6π ring closure and isomerization, eventually resulted in cyclohexenone **97** in high yield as the *trans* isomer. Reaction intermediates, such as α,β -unsaturated ketone **99** and cyclobutenol **100**, were successfully isolated by quenching the reaction at low temperature. As expected, treatment of cyclobutenol **100** with lithium diisopropylamide (LDA) provided cyclohexone **101 a** in 94 % yield. Other pericyclic rearrangements upon 1,2-addition and ring opening can also occur under the reaction conditions, leading to more diverse products (Scheme 33).

In 1993, Cammers-Goodwin reported the ring opening of cyclobutenones triggered by tributylphosphine in alcoholic solvents (Scheme 34).^[43] When 3-phenylcyclobutenone **94** was treated with a catalytic amount of tributylphosphine in methanol, the ring-opening product **102** was afforded, albeit contaminated by approximately 1 % yield of the conjugated ester **103**. Tributylphosphine was demonstrated to be crucial for the transformation as triethylamine failed to afford any desired product. Interestingly, the β,γ -unsaturated ester **102** can be converted into the conjugated product **103** upon gentle heating in alcohol with a base, indicating that the tributylphosphine catalyst led to the kinetic product. In contrast, under the same reaction condition, 3-pentylcyclobutenone gave methoxy-substituted cyclobutanone **104** in high yield. The ring-opening product **105** was afforded only when an increased concentration of tributylphosphine and higher temperature were applied. A mechanism was proposed whereby the reaction started with 1,4-addition of tributylphosphine to the cyclobutenone and concomitant protonation, followed by an S_N1 displacement by methanol to afford cyclobutanone **104**, or a Grob-type fragmentation to give the ring-opening product **102**.

Inspired by the phosphine-catalyzed ring opening reaction, in 2015 Zhang and co-workers reported the enantioselective synthesis of spirocycles **108** through a formal [4+2] cycloaddition between cyclobutenones **106** and isatylidenemalononitriles **107**, catalyzed by amino acid-derived chiral phosphines **109** (Scheme 35).^[44] Compared with phosphine catalysts bearing a single or weak double H-bond donor (e.g., **109 a–e**), phenylalanine-derived catalyst **109 f**, with a strong double H-bond donor moiety, proved optimal, affording the cycloadduct in excellent yield and good enantioselectivity. NaI was proposed to provide a counterion that stabilized the zwitterionic intermediate. The substitution pattern of cyclobutenones only had a marginal effect on reactivity and enantioselectivity.

Unlike Cammers-Goodwin's proposal (Scheme 34),^[43] a 1,2-addition of the phosphine catalyst to the cyclobutenone was proposed as the initial step (Scheme 36). The following ring opening led to a 1,4-dipolar intermediate **110**, which then underwent a stepwise addition/elimination sequence to generate the spiroannulation product.

Such a formal [4+2] cycloaddition with cyclobutenones can also be realized with N-heterocyclic carbene (NHC) catalysts. In 2015, Chi and co-workers reported that a chiral NHC carbene (e.g., **115**) catalyst effectively coupled imines **111** with cyclobutenones **112** in good yield and high enantioselectivity (Scheme 37).^[45] As proposed, the enolate intermediate **113**, formed after the 1,2-addition and concomitant ring opening, underwent a stepwise addition with imines to give lactam **114**, bearing a quaternary carbon center as a single diastereomer. Both cyclic sulfonyl imines (e.g., **116**) and *tert*-butyloxycarbonyl (Boc)-protected isatin-derived imines (e.g., **117**) are suitable substrates. Cyclobutenones with a substituent at the 2-position showed low reactivity in this transformation.

Later, Chi and co-workers also developed a formal [3+2] cycloaddition between 4-chlorocyclobutenones **118** and azomethine imines catalyzed by a chiral isothiourea catalyst **119** (Scheme 38).^[46] Unlike their earlier work,^[45] upon 1,2-addition and ring opening, the α -carbon of the enolate intermediate **120**, instead of the γ -carbon, attacked the azomethine imines and eventually provided heterocycle **121**. A variety of functional groups were well tolerated, giving high diastereo- and enantioselectivity, although bulky substrates lowered both the yields and enantioselectivities. For cyclobutenones bearing a proton or a methyl group at the C4 position or having 4,4-dichloro substituents, no cycloaddition was observed.

The nucleophile-initiated ring opening of benzocyclobutenones was first reported by Cava and Muth in 1960.^[47a] Unlike monocyclic cyclobutenones, the cleavage of benzocyclobutenones could occur at either proximal or distal position, resulting in carboxylic acids **122** and/or **123** (Scheme 39).^[47]

In 1997, the Suzuki group reported that lithium tetramethylpiperidide (LiTMP) could attack the carbonyl moiety of benzocyclobutenone **124**, leading to ring opening predominantly at the C1–C2 bond (Scheme 40).^[48] The resulting anionic intermediate **125** reacted with benzaldehyde smoothly to afford alkoxide intermediate **126**, which, upon intramolecular cyclization, resulted in lactone **127** in decent yields.

In 2011, the Aguilar and co-workers reported the coupling between benzocyclobutenone and Fischer carbene **128** (Scheme 41).^[49] Upon addition with an alkyllithium reagent, benzocyclobutenone **129** underwent ring opening at the proximal position to afford oxy-*ortho*-quinodimethane **130**, which attacks the Fischer carbene to provide intermediate **131**. Followed by a 1,2-metal migration, the seven-membered ring oxonium intermediate **132** was attacked intramolecularly by the alkoxide and then protonation to give bridged compound **133** as the final product. However, yields were generally low for this reaction.

In 2006, Matsuya et al. reported the rapid synthesis of 2,3-benzodiazepine derivatives **134** through a diazoanion-mediated benzocyclobutenone ring-opening transformation (Scheme 42).^[50] Upon oxyanion-facilitated ring opening, an 8π -electrocyclization followed by isomerization led to 2,3-benzodiazepines as the final products. This method was later widely used in the syntheses of benzodiazepine^[51] and unfused diazepine^[52] derivatives.

5. Transition-Metal-Catalyzed Ring opening

Transition metal catalysis has emerged as a powerful tool in the functionalization of chemically inert bonds. Owing to their high ring strain, cyclobutenones are widely used in transition-metal-catalyzed carbon–carbon bond activation reactions.

5.1 Reactions involving stoichiometric transition metals

The first example of transition-metal insertion into cyclobutenones was reported by Liebeskind and co-workers in 1990 (Scheme 43).^[53] Upon treatment with Wilkinson's catalyst, 3-substituted and 2,3-disubstituted cyclobutenones smoothly produced rhodacyclopentenones **135** in good yields through ring-opening at the C1–C4 bond. However, 4-substituted cyclobutenones only afforded a mixture of complex products under the reaction condition, likely due to the steric hindrance.^[54] Interestingly, benzocyclobutenones generally gave two regioisomeric rhodacyclopentenones **136** and **137**, which can arise from cleavage of either the C1–C2 or C1–C8 bond. When an isolated mixture of **136** and **137** was heated at 130 °C, conversion into **137** was observed, indicating that rhodacycle **137** came from **136** under the reaction conditions and was the thermodynamic product. These rhodacyclopentenones **135**–**137** were found to be reluctant to react with alkynes.

In 2013, Yamashita, Nozaki, and Murakami studied the reactivity of a rhodium complex **138** containing an electron-rich pincer-type ligand towards C–C activation (Scheme 44).^[55] Remarkably, the rhodium complex **139** underwent oxidative addition into the benzocyclobutenone C–C bond at room temperature with exclusive C1–C4 selectivity.

The insertion of cobalt complexes was also investigated (Scheme 45).^[54,56] Unlike [CoCl(PPh₃)₃] and [Co(η^5 -C₅H₅)(CO)₂], only the more electron-rich cobalt precatalyst [Co(η^5 -C₉H₇)(PPh₃)₃] **140** was able to react with cyclobutenones to afford η^4 -vinylketene π -complexes **141**. In sharp contrast to the reaction with rhodium (see above, Scheme 43), 3,4-disubstituted cyclobutenones afforded the η^4 -vinylketene products in higher yields as an *E/Z* isomeric mixture. Although the *E* isomers were kinetically favored, they could nevertheless be converted into their *Z* isomers with an elongated reaction time. These η^4 -vinylketene complexes (**141**) are spectroscopically different from the rhodacyclopentenones.^[54] In particular, the stretching frequency of the carbonyl group of **141** is substantially higher, at 1756–1787 cm^{−1}, whereas that of the rhodacyclopentenones appeared at around 1645 cm^{−1}.

Interestingly, 3-ethoxy-substituted cyclobutenone **142** gave cobaltacyclopentenone **143** as the only product (Scheme 46).^[54] A phenyl-substituted cobaltacyclopentenone **144** was obtained when the π complex **141 e** was subjected to a large excess of triphenylphosphine or to a CO atmosphere, although, in both cases, η^4 -vinylketene **141 e** could be regenerated by heating. The treatment of benzocyclobutenone with [Co(η^5 -C₉H₇)(PPh₃)₃] also afforded similar cobaltacyclopentenones **145** and **146**; however, no isomerization was observed when the isolated pure **145** and **146** were subjected to the reaction conditions.

When superstoichiometric ZnCl_2 was used as an additive, a new cobalt complex **147** was isolated with a nonconjugated enone moiety (Scheme 47). The exact reason remains unclear.

Three mechanisms were proposed regarding the formation of η^4 -vinylketene cobalt complexes.^[54] Mechanism 1 (Scheme 48) involves thermolysis of cyclobutenones to form vinylketene intermediate **148**, which was then trapped by the cobalt complex. This mechanism was excluded for two reasons. Firstly, cobalt insertion can happen at room temperature, whereas the thermolysis would be very slow. Secondly, the transformation can occur with isopropanol as solvent, in which the free vinylketene intermediate would be trapped rapidly by solvent instead of the cobalt species. Mechanism 2 is initiated by direct oxidative addition into the C1–C4 bond of cyclobutenones, followed by loss of PPh_3 to give the η^4 -vinylketene cobalt complexes. However, the conversion from isolated **149** into η^4 -vinylketene cobalt complexes was very slow at 60 °C, whereas the insertion of cyclobutenone to form the η^4 - π complex happened smoothly at this temperature. Mechanism 3, which is the most consistent with experimental results, proceeds by pre-coordination of the cyclobutenone carbon–carbon double bond to $[\text{Co}(\eta^5\text{-C}_9\text{H}_7)(\text{PPh}_3)_3]$, followed by phosphine loss and subsequent ring opening.

η^4 -Vinylketene cobalt complex **150** was found to react with a variety of alkynes to produce phenol derivatives (Scheme 49).^[56] When unsymmetrical alkynes were used, a mixture of regioisomers was obtained. 1,5-Cyclooctadiene (cod) was employed as an additive to inhibit alkyne trimerization. However, 3,4-disubstituted η^4 -vinylketene cobalt complex **151** was only able to react with highly electron-deficient dimethyl acetylenedicarboxylate to provide a phenol derivative along with the cyclohexadienone complex **152**.

5.2 Reactions involving catalytic transition metals

In 1991, Liebeskind and co-workers reported the first catalytic phenol synthesis from cyclobutenones and alkynes (Scheme 50).^[57] When the two coupling partners were subjected to a catalytic amount of $[\text{Ni}(\text{cod})_2]$ at 0 °C, phenol products **153** were rapidly formed. Such a transformation was proposed to be initiated by a nickel- η^4 -vinylketene or nickelcyclopentenone intermediate followed by migratory insertion and reductive elimination sequence. Although both 3-substituted and 3,4-disubstituted cyclobutenones were suitable substrates, 2,3-disubstituted cyclobutenones failed to provide the desired products. The tolerance of a tributylstannyl substituent (e.g., **153 e**) allows further functionalization of the product. In contrast to Danheiser's benzannulation,^[13] the nickel-catalyzed phenol synthesis can use non-activated alkynes and gives a different regioselectivity (e.g., *ortho*-**153** vs *para*-**153**). The size of the alkyne substituents only had a marginal effect on the regioselectivity, whereas the heteroatom-containing substrates (e.g., **153 b** and **153 d**) showed an increased regioselectivity, which was proposed to come from a chelation effect (e.g., **154**).

In 2011, Harrity and co-workers expanded the scope of the nickel-catalyzed benzannulation to alkynylboronates **155** (Scheme 51).^[58] Unlike the rhodium catalyst, which provided cyclobutenone dimerization **156** as the major product, $[\text{Ni}(\text{cod})_2]$ yielded the desired phenol boronic ester **157** in good yield and excellent regioselectivity, which allowed further

derivatization through cross-coupling reactions. As this method was compatible with 3,4-disubstituted cyclobutenones, the regioselectivities of the coupling with 3-substituted cyclobutenones were predominantly determined by the substituents on the alkynylboronates. For example, sp^3 -based substituents showed high selectivity for the regioisomer where the pinacolboranyl (Bpin) group was *meta* to the phenolic OH (e.g., **158 a**), whereas sp^2 -based substituents resulted in a modest level of selectivity for the regioisomer where the Bpin group was *ortho* to the phenolic OH (e.g., **158 b**).

In 2015, the same group reported a detailed study of the nickel-catalyzed benzannulation with alkynes bearing aryl substituents (Scheme 52).^[59] When norbornadiene was added as a promoter to enhance the catalyst lifetime, various alkynes were compatible, giving heavily substituted phenols in good yields, and the regioselectivity was predominantly controlled by the substituents on the alkynes: an alkyl group tends to be incorporated *ortho* to the phenol OH moiety, whereas an aryl group prefers to stay at the *meta* position. Unlike Liebeskind's work,^[57] the presence of heteroatoms only had a marginal effect on the regioselectivity.

Two mechanisms were proposed for this benzannulation reaction (Scheme 53).^[59] The first one is initiated by ring opening of the cyclobutenone to form either a metallocyclopentenone or a η^4 -vinylketene complex **159**, which would then undergo migratory insertion and reductive elimination to afford the phenol product. The second reaction pathway was initiated by cyclometallation between the carbonyl group and the alkyne to form metallacycle **160**. Similarly to the nickel-catalyzed cyclobutanone activation,^[60] subsequent β -carbon elimination and reductive elimination would provide the phenol products. While both mechanisms are possible, the isolation of ester **161**, when adding phenol to the standard reaction conditions, suggested involvement of a vinylketene-like intermediate.

In 1993, Huffman and Liebeskind developed a transitionmetal-catalyzed synthesis of medium-sized rings from cyclopropyl and cyclobutyl-substituted cyclobutenones **162** (Scheme 54).^[61] It was proposed that the rhodium or ruthenium first inserted into the C1–C4 bond of cyclobutenone **162**, which was followed by β -carbon elimination and reductive elimination to give cycloheptadienone **163**. Product **164** was also obtained, likely through an olefin isomerization from **163** by 1,5-hydrogen shift under thermal conditions.

In 2004, Kondo, Mitsudo, and co-workers developed a Rh/Ru-catalyzed dimerization of cyclobutenones (Scheme 55).^[62] When cyclobutenone **165** was treated with a catalytic amount of $[RuCl_2(CO)_3]_2$ in refluxing toluene, 2-pyranone **166** was afforded as a mixture of *E* and *Z* isomers. However, use of $[Rh(CO)_2Cl]_2$ provided the desired dimers in *E* configuration as the sole product.

In addition, with $[Rh(CO)_2Cl]_2$ as the catalyst, the reaction between cyclobutenone and norbornene predominantly gave the decarbonylative coupling product **167** in good yield. However, under a CO atmosphere, the direct coupling occurred to give cyclohexenone **168** (Scheme 56). Interestingly, when ^{13}CO was used, the corresponding ^{13}C -labeled product (**168 b**) was formed, indicating a facile yet reversible decarbonylation/reinsertion process.

In 2007, Kondo and co-workers reported a new phenol synthesis from cyclobutenones and electron-deficient olefins **169** with the help of a rhodium catalyst (Scheme 57).^[63] Tricyclohexylphosphine was employed as a ligand to increase the catalyst reactivity and suppress cyclobutenone dimerization. Although the precise mechanism remains unclear, it was proposed that, upon ring opening, the resulting η^4 -vinyl-ketene intermediate **170** underwent a Diels–Alder-type cyclization to afford cyclohexenone **171**, which, upon dehydrogenation, led to the phenol products with high regioselectivity.

In 2012, Xu and Dong developed a Rh-catalyzed intramolecular coupling between olefins and benzocyclobutenones, whereby insertion occurred selectively at the C1–C2 position (Scheme 58).^[64] Upon treatment with $[\{\text{Rh}(\text{cod})\text{Cl}\}_2]$ and 1,4-bis(diphenylphosphino)butane (dppb) as a bidentate phosphine ligand, benzocyclobutenone **172** tethered with an olefin moiety afforded the benzofused ring structure **173**. Ligands with a larger bite angle were found to give higher yields. This method is compatible with a variety of mono- and 1,1-disubstituted alkenes. For more challenging substrates, such as those containing longer tethers, aryl substituents or trisubstituted alkenes, ZnCl_2 was employed as a Lewis acid cocatalyst.

The enantioselective version of this transformation was later reported by the same group (Scheme 59).^[65] (*R*)-DTBM-SEG-PHOS and (*R,R*)-DIOP were found to be suitable chiral ligands to provide excellent enantioselectivity. The resulting products can be reduced to the corresponding saturated tricycles with four consecutive chiral centers (e.g., **174**).

To better understand the mechanism and the ligand effect responsible for the observed high reactivity and enantioselectivity, Liu and co-workers undertook a detailed computational study of this rhodium-catalyzed intramolecular carboacylation reaction (Scheme 60).^[66] According to the DFT calculations, instead of direct insertion into the C1–C2 bond of benzocyclobutenone (Scheme 60, path b), the cleavage is more likely to occur from cleavage of the C1–C8 bond (Scheme 60, path a), followed by a carbonyl migratory process with CO reinsertion being the rate-determining step (RDS). The increased reactivity of bidentate ligands with larger bite angles was attributed to release of the ligand–substrate repulsion in intermediate **175**. The enantioselectivity was attributed to the steric repulsion between the equatorial aryl group of the chiral ligand and the C8-methylene of the benzocyclobutenone substrate.

To enable insertion of more sterically hindered trisubstituted alkenes, such as **176**, Xu and Dong developed a new catalyst system that employed electron-deficient $[\{\text{Rh}(\text{CO})_2\text{Cl}\}_2]$ and $\text{P}(\text{C}_6\text{F}_5)_3$ (Scheme 61).^[67] By using this method, the tetracyclic core of the proposed cycloinnumakiol **177** was efficiently constructed. Ultimately, the syntheses of both diastereomers of the proposed structure were achieved, leading to the correct structural assignment of the natural product.

In addition to forming fused rings, spirocycles can also be constructed from olefin-tethered benzocyclobutenones through properly choosing the catalyst system (Scheme 62).^[68] It was anticipated that, after Rh insertion into the benzocyclobutenone C–C bond and subsequent

migratory insertion into the olefin, whereas reductive elimination would lead to fused-ring products, β -H elimination and decarbonylation would afford the spirocycle **178**.

When 10 mol % of $\text{P}(\text{C}_6\text{F}_5)_3$ was used alongside 5 mol % $[\{\text{Rh}(\text{CO})_2\text{Cl}\}_2]$, a variety of cyclic trisubstituted olefins, as well as electron-deficient enone moieties, can react to give spirocycles **179** in decent yields, whereas in some cases the products were isolated as isomeric mixtures of alkenes (e.g., **179 b**, **d**, **g**, and **h**; Scheme 63). Due to the near-neutral reaction conditions, this spirocyclization transformation tolerated functional groups including ketones, enamides, esters, dienes, benzyl and vinyl ethers, and even unprotected tertiary alcohols.

The intramolecular coupling between alkynes and benzocyclobutenones was also reported by Dong and co-workers (Scheme 64).^[69] The mechanism was proposed to be similar to the intramolecular alkene carboacylation.^[64–66] The resulting enone moiety from the “cut and sew” sequence underwent tautomerization to give β -naphthols as the final product. ZnCl_2 was again found to be an effective Lewis acid cocatalyst to assist medium-sized ring formation (e.g., **180 d** and **180 e**).

Interestingly, a decarbonylative coupling was realized by using DTBM-SEPHOS as the ligand under argon flow, whereby tricyclic indenenes (e.g., **181**) were isolated as the products (Scheme 65). Thus, a divergent approach from alkyne-tethered benzocyclobutenones was realized (Scheme 66).

Apart from coupling with nonpolar π bonds in substrates such as alkenes and alkynes, the Dong group later expanded the “cut and sew” transformation by utilizing more polar $\text{C}=\text{N}$ bonds, for example, in oximes, as intramolecular coupling partners (Scheme 67).^[70] The cationic $[\{\text{Rh}(\text{cod})(\text{CH}_3\text{CN})_2\}]\text{BF}_4$ was found to be the optimal precatalyst. High enantioselectivity and decent yields were achieved by using a combination of two chiral ligands. This method provides an unusual way to prepare chiral fused lactams.

In 2015, Martin and co-workers developed a nickel-catalyzed intermolecular coupling reaction between benzocyclobutenone and dienes or acetylenes (Scheme 68).^[71] Upon treatment with $[\text{Ni}(\text{cod})_2]$ as the precatalyst and tri(4-trifluoromethylphenyl)phosphine as the ligand, benzocyclobutenone **182** reacted smoothly with diene **183** to afford the [4+4] cyclization product **184 a** in excellent yield, with a trace amount of formal [4+2] addition product **185**. The coupling with unsymmetrical 1,3-dienes occurred in a regio- and diastereoselective fashion (e.g., **184 c** and **184 i**). For substrates having enolizable protons, an intramolecular transannular aldol reaction occurred under the reaction conditions, giving **186** as a single regioisomer and diastereoisomer. By using PPh_3 as the ligand, the coupling between benzocyclobutenone and diphenylacetylene **187** occurred smoothly to give enone **188** in excellent yields. In all cases, no C1–C4 cleavage was observed.

In 2015, Dong and co-workers reported a rhodium-catalyzed decarbonylative coupling of 3-amino-4,4-disubstitutedcyclo-butenones **189** and tethered alkenes to access [3.1.0]bicyclic products **190** (Scheme 69).^[72] Under thermal conditions, vinyl cyclobutanone product **191** was isolated, in the form of two conformational isomers, **191 a** and **191 a'**, presumably

because the C3–N bond is incapable of free rotation. In contrast, when a more electron-deficient precatalyst [$\{\text{Rh}(\text{CO})_2\text{Cl}\}_2$] was used, [3.1.0]bicyclic **190** was the predominant product. The transformation was proposed to be initiated by retro-4 π electrocyclization of the cyclobutenone moiety through thermolysis. The resulting vinylketene intermediate **192** either undergoes [2+2] with the tethered alkene moiety to give the cyclobutanone product or a rhodium-catalyzed cyclometallation to give five-membered rhodacycle **193**, which can undergo subsequent decarbonylative reductive elimination to form cyclopropane **190**. When isolated cyclobutanones **191 a** and **191 a'** were subjected to the [$\{\text{Rh}(\text{CO})_2\text{Cl}\}_2$] conditions, no formation of cyclopropane **190** was observed, which excluded the possibility of forming cyclopropane **190** by decarbonylation of cyclobutanone **191**. This transformation can tolerate different amide groups and tether linkages. Interestingly, when a 1,1-disubstituted olefin was used as the coupling partner, fused cyclobutanone was formed exclusively in high yield.

In 2015, by adding a cyclopropyl group to the benzocyclobutenone moieties, Yu and co-workers developed a rhodium-catalyzed [7+1] cycloaddition to form benzocyclooctenones (BCOs; Scheme 70).^[73] Synthesized by 1,2-addition of cyclopropyl Grignard reagents to benzocyclobutenones, benzocyclobutene **194** underwent facile ring opening to form diene intermediate **195**. Assisted by a rhodium catalyst, cyclometallation and then β -carbon elimination occurred to give intermediate **196**, which, upon CO insertion, afforded BCO **197** in good yield.

Although most transition-metal-catalyzed cyclobutenone and benzocyclobutenone-mediated transformations involve coupling with unsaturated units, Dong and co-workers recently reported a reagent-free catalytic ring expansion of cyclobutenones and benzocyclobutenones bearing 3-alkyl substituents (Scheme 71).^[74] As indicated in Liebeskind's work,^[53] the insertion of rhodium into the C1–C8 bond of benzocyclobutenones is kinetically favored. In the absence of coordination of unsaturated π moieties,^[64–70] concomitant β -H elimination could occur after the oxidative addition to give the acylrhodium species **198**, which, upon hydride re-insertion and reductive elimination, leads to ring-expanded five-membered ring products **199**.

By properly choosing the Rh precatalyst and bidentate phosphine ligands, benzocyclobutenones and simple cyclobutenones both afforded the desired ring-expansion product, that is, 1-indanones and cyclopentenones, in good yields (Scheme 72). The involvement of a β -H elimination/migratory insertion/reductive elimination sequence was supported by deuterium labeling experiments and the isolation of ring-opening olefin products **200** and **201**.

6. Diels–Alder Reactions

Besides having been widely investigated for ring-opening transformations, cyclobutenones also possess remarkable dienophilicity compared with cyclopentenones and cyclohexenones, which is promoted by the strain relief during the cycloaddition. Hence, cyclobutenones can efficiently participate in Diels–Alder reactions. The first Diels–Alder reaction that employed cyclobutenones as dienophiles was reported by Kelly and McNutt in 1975 (Scheme 73).^[75]

Treatment of 4,4-dimethylcyclobutenone **202** with 1,3-diphenylisobenzofuran **203** proceeded smoothly to afford adduct **204** in 62 % yield. $\text{BF}_3 \cdot \text{OEt}_2$ was used as a catalyst for the reaction between cyclopentadiene and **202**, from which the *exo* adduct **205** was isolated as the major product.

In 1991, Viehe, Sustmann, and co-workers reported the Diels–Alder reaction of 3-cyanocyclobutenone **206** (Scheme 74).^[76] Cycloaddition with a variety of dienes proceeded with high efficiency at ambient temperature.

In 2010, Danishefsky and co-workers reported the first Diels–Alder cycloaddition that incorporated the unsubstituted parent structure of cyclobutenones **207** (Scheme 75).^[77] A variety of dienes underwent cycloaddition rapidly and some reactions even occurred at $-40\text{ }^\circ\text{C}$. When less reactive dienes, such as furan and 2,3-dimethylbutadiene, were used, ZnCl_2 was added as a Lewis acid catalyst to promote the transformation.

7. Conclusion and Outlook

The high yet predictable reactivities of cyclobutenones and benzocyclobutenones allow them to serve as versatile synthons, providing two, three or four carbon atoms in a variety of synthetic transformations. Their participation in such a large variety of reactions, including Diels–Alder reactions, [2+2] electrocyclizations, and cycloadditions with 1,3-dipolar species, has allowed their use in the synthesis of various structurally attractive and biologically important molecules. The recent development of transition-metal-catalyzed C–C bond activations of cyclobutenones and benzocyclobutenones has led to a renaissance of the field for new reactivity discovery. With the advent of more facile routes to cyclobutenones and benzocyclobutenones, it is expected that more synthetically useful applications of these intriguing molecules will be developed.

Acknowledgments

We thank NIGMS (R01GM109054) and the Welch Foundation (F 1781) for research grants. G.D. is a Searle Scholar and Sloan Fellow. Prof. Dr. M. C. Young is greatly acknowledged for editing and proof-reading the manuscript.

References

1. For reviews on cyclobutanes, see: Namyslo JC, Kaufmann DE. Chem Rev. 2003; 103:1485–1537. [PubMed: 12683789] Sadana AK, Saini RK, Billups WE. Chem Rev. 2003; 103:1539–1602. [PubMed: 12683790] Nemoto H. Chem Pharm Bull. 2007; 55:961–974. [PubMed: 17603183] Seiser T, Saget T, Tran DN, Cramer N. Angew Chem Int Ed. 2011; 50:7740–7752. Angew Chem. 2011; 123:7884–7896.
2. For reviews on cyclobutenones and benzocyclobutenones, see: Belluš D, Ernst B. Angew Chem Int Ed Engl. 1988; 27:797–827. Angew Chem. 1988; 100:820–850. Martin R, Flores-Gaspar A. Synthesis. 2013:563–580. Kondo T, Mitsudo T-A. Chem Lett. 2005; 34:1462–1467. The term “benzocyclobutanone” was occasionally used. However, due to the unsaturated nature of the benzene moiety and the similar reactivity as cyclobutenones, the term “benzocyclobutenone” is considered a more accurate description.
3. a) Aidhen IS, Ahuja JR. Tetrahedron Lett. 1992; 33:5431–5432. b) Zhao YL, Yang SC, Di CH, Han XD, Liu Q. Chem Commun. 2010; 46:7614–7616. c) Álvarez-Bercedo P, Flores-Gaspar A, Correa A, Martin R. J Am Chem Soc. 2010; 132:466–467. [PubMed: 20038097] d) Martin R, Flores-

- Gaspar A. *Org Synth.* 2012; 89:159–169.e) Flores-Gaspar A, Gutiérrez-Bonet Á, Martín R. *Org Lett.* 2012; 14:5234–5237. [PubMed: 23030460] f) Mehta G, Kotha S. *Tetrahedron.* 2001; 57:625–659.h) Liebman JF, Greenberg A. *Chem Rev.* 1976; 76:311.
4. a) Souillart L, Cramer N. *Chem Rev.* 2015; 115:9410–9464. [PubMed: 26044343] b) Dong, G., editor. *Topics in Current Chemistry*, Vol. 346: C–C Bond Activation. Springer; Heidelberg: 2014. c) Chen F, Wang T, Jiao N. *Chem Rev.* 2014; 114:8613–8661. [PubMed: 25062400] d) Jones WD. *Nature.* 1993; 364:676–677.e) Murakami, M., Ito, Y. *Activation of Unreactive Bonds and Organic Synthesis.* Murai, S. Alper, H. Gossage, RA. Grushin, VV. Hidai, M. Ito, Y. Jones, WD. Kakiuchi, F. van Koten, G. Lin, YS. Mizobe, Y. Murai, S. Murakami, M. Richmond, TG. Sen, A. Sugimoto, M., Yamamoto, A., editors. Vol. 3. Springer; Heidelberg: 1999. p. 97–129.f) Rybtchinski B, Milstein D. *Angew Chem Int Ed.* 1999; 38:870–883. *Angew Chem.* 1999; 111:918–932.g) Perthuisot C, Edelbach BL, Zubris DL, Simhai N, Iverson CN, Müller C, Satoh T, Jones WD. *J Mol Catal A.* 2002; 189:157–168.h) van der Boom ME, Milstein D. *Chem Rev.* 2003; 103:1759–1792. [PubMed: 12744693] i) Miura, M., Satoh, T. *Palladium in Organic Synthesis.* Tsuji, J., editor. Vol. 14. Springer; Heidelberg: 2005. p. 1–20.j) Jun, C-H., Park, J-W. *Directed Metallation.* Chatani, N., editor. Vol. 24. Springer; Heidelberg: 2007. p. 117–143.k) Cramer N, Seiser T. *Synlett.* 2011:449–460.l) Murakami M, Matsuda T. *Chem Commun.* 2011; 47:1100–1105.m) Ruhland K. *Eur J Org Chem.* 2012:2683–2706.n) Dermenci A, Coe JW, Dong G. *Org Chem Front.* 2014; 1:567–581.
5. a) Moore HW, Decker OHW. *Chem Rev.* 1986; 86:821–830.b) Dolbier WR Jr, Koroniak H, Houk KN, Sheu C. *Acc Chem Res.* 1996; 29:471–477.
6. Niwayama S, Kallel EA, Sheu CM, Houk KN. *J Org Chem.* 1996; 61:2517–2522.
7. Jenny EF, Roberts JD. *J Am Chem Soc.* 1956; 78:2005–2009.
8. Mayr H. *Angew Chem Int Ed Engl.* 1975; 14:500–501. *Angew Chem.* 1975; 87:491–492.
9. Huisgen R, Mayr H. *J Chem Soc Chem Commun.* 1976:55–56.
10. Ficini J, Falou S, Dangelo J. *Tetrahedron Lett.* 1977; 18:1931–1934.
11. For a review on ynamines, see: Ficini J. *Tetrahedron.* 1976; 32:1449–1486.
12. Danheiser RL, Gee SK, Sard H. *J Am Chem Soc.* 1982; 104:7670–7672.
13. Danheiser RL, Gee SK. *J Org Chem.* 1984; 49:1672–1674.
14. Danheiser RL, Gee SK, Perez JJ. *J Am Chem Soc.* 1986; 108:806–810.
15. Kowalski CJ, Lal GS. *J Am Chem Soc.* 1988; 110:3693–3695.
16. Danheiser RL, Nishida A, Savariar S, Trova MP. *Tetrahedron Lett.* 1988; 29:4917–4920.
17. Delaunoy M, Ghosez L. *Angew Chem Int Ed Engl.* 1969; 8:72–73. *Angew Chem.* 1969; 81:33–34.
18. Mak XY, Crombie AL, Danheiser RL. *J Org Chem.* 2011; 76:1852–1873. [PubMed: 21322545]
19. Karlsson JO, Nguyen NV, Foland LD, Moore HW. *J Am Chem Soc.* 1985; 107:3392–3393.
20. Perri ST, Foland LD, Decker OHW, Moore HW. *J Org Chem.* 1986; 51:3067–3068.
21. Moore HW, Perri ST. *J Org Chem.* 1988; 53:996–1003.
22. Krysan DJ, Gurski A, Liebeskind LS. *J Am Chem Soc.* 1992; 114:1412–1418.
23. a) Ezcurra JE, Karabelas K, Moore HW. *Tetrahedron.* 2005; 61:275–286.b) Tiedemann R, Turnbull P, Moore HW. *J Org Chem.* 1999; 64:4030–4041.c) Onofrey TJ, Gomez D, Winters M, Moore HW. *J Org Chem.* 1997; 62:5658–5659.d) Heerding JM, Moore HW. *J Org Chem.* 1991; 56:4048–4050.e) Enhnen A, Karabelas K, Heerding JM, Moore HW. *J Org Chem.* 1990; 55:1177–1185.f) Foland LD, Decker OHW, Moore HW. *J Am Chem Soc.* 1989; 111:989–995.g) Xu SL, Taing M, Moore HW. *J Org Chem.* 1991; 56:6104–6109.h) Xia H, Moore HW. *J Org Chem.* 1992; 57:3765–3766.i) Xu SL, Moore HW. *J Org Chem.* 1992; 57:326–338.
24. a) Liu F, Liebeskind LS. *J Org Chem.* 1998; 63:2835–2844.b) Sun L, Liebeskind LS. *J Org Chem.* 1995; 60:8194–8203.c) Liebeskind LS, Granberg KL, Zhang J. *J Org Chem.* 1992; 57:4345–4352.d) Edwards JP, Krysan DJ, Liebeskind LS. *J Am Chem Soc.* 1993; 115:9868–9869.e) Gurski A, Liebeskind LS. *J Am Chem Soc.* 1993; 115:6101–6108.f) Kwan HL, Moore HW. *Tetrahedron Lett.* 1993; 34:235–238.g) Liebeskind LS, Wang J. *J Org Chem.* 1993; 58:3550–3556.h) Liebeskind LS, Bombrun A. *J Org Chem.* 1994; 59:1149–1159.
25. a) Harrowven DC, Pascoe DD, Demurtas D, Bourne HO. *Angew Chem Int Ed.* 2005; 44:1221–1222. *Angew Chem.* 2005; 117:1247–1248.b) Peña-Cabrera E, Liebeskind LS. *J Org Chem.* 2002;

- 67:1689–1691. [PubMed: 11871905] c) Perri ST, Dyke HJ, Moore HW. J Org Chem. 1989; 54:2032–2034.
26. Harrowven DC, Pascoe DD, Guy IL. Angew Chem Int Ed. 2007; 46:425–428. Angew Chem. 2007; 119:429–432.
27. a) Xu SL, Moore HW. J Org Chem. 1989; 54:6018–6021. b) Xu SL, Xia H, Moore HW. J Org Chem. 1991; 56:6094–6103.
28. Hachiya I, Yoshitomi T, Yamaguchi Y, Shimizu M. Org Lett. 2009; 11:3266–3268. [PubMed: 19572744]
29. Hachiya I, Ito A, Shimizu M. Asian J Org Chem. 2014; 3:614–618.
30. Baldwin JE, Mcdaniel C. J Am Chem Soc. 1967; 89:1537–1537.
31. Baldwin JE, Mcdaniel MC. J Am Chem Soc. 1968; 90:6118–6124.
32. Kikuchi O. Bull Chem Soc Jpn. 1982; 55:1669.
33. Chapman OL, Lassila JD. J Am Chem Soc. 1968; 90:2449–2450.
34. Toda F, Todo E. Chem Lett. 1974:1279–1280.
35. Toda F, Todo E. Bull Chem Soc Jpn. 1976; 49:2503–2505.
36. Toda F, Todo Y, Todo E. Bull Chem Soc Jpn. 1976; 49:2645–2646.
37. Schiess P, Eberle M, Huysfrancotte M, Wirz J. Tetrahedron Lett. 1984; 25:2201–2204.
38. a) Roberts JD, Kline GB, Simmons HE. J Am Chem Soc. 1953; 75:4765–4768. b) Silversmith EF, Roberts JD. J Am Chem Soc. 1956; 78:4023–4024. c) Silversmith EF, Kitahara Y, Caserio MC, Roberts JD. J Am Chem Soc. 1958; 80:5840–5845.
39. Hassner A, Dillon JL. J Org Chem. 1983; 48:3382–3386.
40. Murakami M, Miyamoto Y, Ito Y. Angew Chem Int Ed. 2001; 40:189–190. Angew Chem. 2001; 113:182–184.
41. Murakami M, Miyamoto Y, Ito Y. J Am Chem Soc. 2001; 123:6441–6442. [PubMed: 11427081]
42. Magomedov NA, Ruggiero PL, Tang Y. J Am Chem Soc. 2004; 126:1624–1625. [PubMed: 14871080]
43. Cammers-Goodwin A. J Org Chem. 1993; 58:7619–7621.
44. Li Y, Su X, Zhou W, Li W, Zhang J. Chem Eur J. 2015; 21:4224–4228. [PubMed: 25630968]
45. Li BS, Wang Y, Jin Z, Zheng P, Ganguly R, Chi YR. Nat Commun. 2015; 6:6207. [PubMed: 25652912]
46. Li B-S, Wang Y, Jin Z, Chi YR. Chem Sci. 2015; 6:6008–6012.
47. a) Cava MP, Muth K. J Am Chem Soc. 1960; 82:652–654. b) Bradley JC, Durst T. Can J Chem. 1995; 73:1660–1665. c) Gokhale A, Schiess P. Helv Chim Acta. 1998; 81:251–267.
48. Matsumoto T, Hamura T, Kuriyama Y, Suzuki K. Tetrahedron Lett. 1997; 38:8985–8988.
49. García-García P, Novillo C, Fernández-Rodríguez MA, Aguilar E. Chem Eur J. 2011; 17:564–571. [PubMed: 21207574]
50. Matsuya Y, Ohsawa N, Nemoto H. J Am Chem Soc. 2006; 128:13072–13073. [PubMed: 17017785]
51. Matsuya Y, Katayanagi H, Ohdaira T, Wei ZL, Kondo T, Nemoto H. Org Lett. 2009; 11:1361–1364. [PubMed: 19231851]
52. Sugimoto K, Hayashi R, Nemoto H, Toyooka N, Matsuya Y. Org Lett. 2012; 14:3510–3513. [PubMed: 22721410]
53. Huffman MA, Liebeskind LS, Pennington WT. Organometallics. 1990; 9:2194–2196.
54. Huffman MA, Liebeskind LS, Pennington WT. Organometallics. 1992; 11:255–266.
55. Masuda Y, Hasegawa M, Yamashita M, Nozaki K, Ishida N, Murakami M. J Am Chem Soc. 2013; 135:7142–7145. [PubMed: 23641905]
56. Huffman MA, Liebeskind LS. J Am Chem Soc. 1990; 112:8617–8618.
57. Huffman MA, Liebeskind LS. J Am Chem Soc. 1991; 113:2771–2772.
58. Auvinet AL, Harrity JP. Angew Chem Int Ed. 2011; 50:2769–2772. Angew Chem. 2011; 123:2821–2824.

59. Stalling T, Harker WR, Auvinet AL, Cornel EJ, Harrity JP. *Chem Eur J*. 2015; 21:2701–2704. [PubMed: 25491134]
60. Murakami M, Ashida S, Matsuda T. *J Am Chem Soc*. 2005; 127:6932–6933. [PubMed: 15884919]
61. Huffman MA, Liebeskind LS. *J Am Chem Soc*. 1993; 115:4895–4896.
62. Kondo T, Taguchi Y, Kaneko Y, Niimi M, Mitsudo TA. *Angew Chem Int Ed*. 2004; 43:5369–5372. *Angew Chem*. 2004; 116:5483–5486.
63. Kondo T, Niimi M, Nomura M, Wada K, Mitsudo T-a. *Tetrahedron Lett*. 2007; 48:2837–2839.
64. Xu T, Dong G. *Angew Chem Int Ed*. 2012; 51:7567–7571. *Angew Chem*. 2012; 124:7685–7689.
65. Xu T, Ko HM, Savage NA, Dong G. *J Am Chem Soc*. 2012; 134:20005–20008. [PubMed: 23171396]
66. Lu G, Fang C, Xu T, Dong G, Liu P. *J Am Chem Soc*. 2015; 137:8274–8283. [PubMed: 26051406]
67. Xu T, Dong G. *Angew Chem Int Ed*. 2014; 53:10733–10736. *Angew Chem*. 2014; 126:10909–10912.
68. Xu T, Savage NA, Dong G. *Angew Chem Int Ed*. 2014; 53:1891–1895. *Angew Chem*. 2014; 126:1922–1926.
69. Chen PH, Xu T, Dong G. *Angew Chem Int Ed*. 2014; 53:1674–1678. *Angew Chem*. 2014; 126:1700–1704.
70. Deng L, Xu T, Li H, Dong G. *J Am Chem Soc*. 2016; 138:369–374. [PubMed: 26674855]
71. Juliá-Hernández F, Ziadi A, Nishimura A, Martin R. *Angew Chem Int Ed*. 2015; 54:9537–9541. *Angew Chem*. 2015; 127:9673–9677.
72. Zhou X, Zafar I, Dong G. *Tetrahedron*. 2015; 71:4478–4483. [PubMed: 26034330]
73. a) Fu XF, Xiang Y, Yu ZX. *Chem Eur J*. 2015; 21:4242–4246. [PubMed: 25612241] b) Wang Y, Yu ZX. *Acc Chem Res*. 2015; 48:2288–2296. [PubMed: 26227886]
74. Chen P-H, Sieber J, Senanayake CH, Dong G. *Chem Sci*. 2015; 6:5440–5445. [PubMed: 28042458]
75. Kelly TR, McNutt RW. *Tetrahedron Lett*. 1975; 16:285–288.
76. Bienfait B, Coppemotte G, Merenyi R, Viehe HG, Sicking W, Sustmann R. *Tetrahedron*. 1991; 47:8167–8176.
77. a) Li X, Danishefsky SJ. *J Am Chem Soc*. 2010; 132:11004–11005. [PubMed: 20698657] b) Pham HV, Paton RS, Ross AG, Danishefsky SJ, Houk KN. *J Am Chem Soc*. 2014; 136:2397–2403. [PubMed: 24410341]

Biographies

Peng-hao Chen received his BSc. degree in Chemistry from the University of Science and Technology of China (P. R. China) in 2012 under the supervision of Professor Liu-zhu Gong. He is currently a PhD candidate in organic chemistry under the supervision of Professor Guangbin Dong at the University of Texas at Austin (USA). His research is focused on transition-metal-catalyzed C–C bond activation.

Guangbin Dong received his BSc. degree from Peking University (P. R. China) and completed his PhD in chemistry from Stanford University (USA) with Professor Barry M. Trost, where he was a Larry Yung Stanford Graduate Fellow. In 2009, he began research with Prof. Robert H. Grubbs at California Institute of Technology (USA) as a Camille and Henry Dreyfus Environmental Chemistry Fellow. In 2011, he joined the Department of Chemistry and Biochemistry at the University of Texas at Austin (USA) as an Assistant Professor and a CPRIT Scholar. Currenty, he is a Professor of Chemistry at the University of Chicago. His research interests entail the development of powerful chemical tools for addressing questions of biological importance.

Scheme 1.
General reaction profiles for cyclobutenones.

Scheme 2.
“Inward” and “outward” rotation.

Scheme 3.

Preliminary observation for vinylketene intermediate.

Scheme 4.

Vinylketene intermediates react with olefins and nucleophiles.

Scheme 5.
Ficini-type rearrangement with ynamines and cyclobutenones.

Scheme 6.

Formation of cyclooctane derivatives by sigmatropic rearrangement.

Scheme 7.

Phenol formation by [2+2] cycloaddition of cyclobutenones with heterodisubstituted alkynes.

Scheme 8.

Total synthesis of mycophenolic acid. MOM=methoxymethyl acetal.

Scheme 9.

Benzannulation with trialkylsiloxyalkynes as coupling partners. TIPS=triisopropylsilyl.

Scheme 10.

Allene side-product formation in Danheiser benzannulation.

Scheme 11.

Three different regioisomers involved in the benzannulation.

selected examples

toluene, 80°C, 1.5 h
then 110°C, 1.5 h
84%

toluene, 110°C, 10 h
53%

toluene, 80°C, 1.5 h
then 110°C, 1.5 h
79%

Scheme 12.

Ynamides as coupling partners for regioselective benzannulation.

Scheme 13.

Formal synthesis of (+)-FR900482. Teoc=2-(trimethylsilyl)ethoxycarbonyloxy, TBS=*t*-butyldimethylsilyl.

Scheme 14.

Intramolecular cyclization followed by thermal ring opening.

Scheme 15.

Benzoquinone synthesis involving silyl or ally migration.

Scheme 16.
Benzoquinone synthesis involving intramolecular cyclization.

Scheme 17.
Substituent effects on benzoquinone synthesis.

Scheme 18.

Hydroxy group inward rotation in the presence of an electron-donating group.

selected examples

Scheme 19.

Phenol synthesis by one-pot Pd-catalyzed cross-coupling/benzannulation sequence.

Scheme 20.
Oxidation-free benzoquinone synthesis.

Scheme 21.

Substituent-controlled formation of carbocyclic ring vs. quinone derivatives.

Scheme 22.

Formation of benzobicyclo[3.2.1] structure via diradical intermediate.

Scheme 23.

Bicyclo[3.2.0]heptenone formation from 3-allylcyclobutenones.

Scheme 24.

Base-controlled chiral lactam formation through enantioselective reduction of iminocyclobutenones. PMP = *p*-methoxyphenyl.

Scheme 25.

Inward and outward ring opening by thermolysis and photolysis.

Scheme 26.
Formation of vinylketene under photolysis.

Scheme 27.

Intramolecular nucleophilic attack after ring opening.

Scheme 28.
Ring opening with concomitant one-carbon loss.

Scheme 29.

Reaction of benzocyclobutenones with activated carbonyl and dienophiles.

Scheme 30.

Early observation of nucleophile-induced ring opening.

Scheme 31.

Silyl nucleophile induces stereoselective synthesis of 1,3-dienes.

Scheme 32.
Lithium salt induced electrocyclic reactions.

Scheme 33.

Pericyclic rearrangements involved in the ring opening. ERC=electrocyclic ring closure.

Scheme 34.
Tributylphosphine-catalyzed ring opening of cyclobutenones.

Scheme 35.

Chiral phosphine-catalyzed enantioselective spirocyclization. PMB=*p*-me-thoxybenzyl.

Scheme 36.

Mechanism for chiral phosphine-catalyzed enantioselective spirocyclization.

Scheme 37.
Chiral N-heterocyclic carbene-catalyzed enantioselective lactam synthesis.

Scheme 38.

Chiral isothioureacatalyzed formal [3+2] cycloaddition.

Scheme 39.

Early observation of nucleophile-induced ring opening of benzocyclobutenones.

Scheme 41.

Lithium salt-induced ring opening trapped by Fischer carbenes.

selected example

Scheme 42.

Diazoanion-induced ring opening and synthesis of 2,3-benzodiazepine derivatives.

Scheme 43.

The first example of transition-metal insertion into cyclobutenones.

Scheme 44.

PBP pincer rhodium complex-enabled selective insertion of benzocyclobutenones.

selected examples

Scheme 45.

η^4 -Vinylketene complex formation through insertion of cobalt complexes into cyclobutenones.

Scheme 46.

Cobaltacyclopentenone complex formation with cyclobutenones and benzocyclobutenones.

Scheme 47. Formation of different cobaltacyclopentenones by addition of ZnCl_2 .

Scheme 48.

Three mechanisms proposed for the formation of η^4 -vinylketene complexes.

Scheme 49.
Stoichiometric cobalt-enabled benzannulation reaction.

Scheme 50.

Nickel-catalyzed benzannulation of cyclobutenones and alkynes.

selected examples

Scheme 51.

Nickel-catalyzed benzannulation with alkynylboronates.

Scheme 52.
Regioselectivity in nickel-catalyzed benzannulation.

Scheme 53.
Mechanism for nickel-catalyzed benzannulation.

Scheme 54.
Rhodium-catalyzed medium-size ring synthesis.

Scheme 55.

Dimerization of 2,3-disubstituted cyclobutenones.

Scheme 56.

Direct/decarbonylative coupling between cyclobutenones and norbornene.

Scheme 57.

Phenol synthesis from cyclobutenones and electron-deficient alkenes.

Scheme 58.

Rhodium-catalyzed carboacylation for the synthesis of benzofused ring systems. Values in parentheses refer to the yields based on recovered starting material.

Scheme 59.

Enantioselective intramolecular carboacylation. THS=tetrabutylammonium hydrogen sulfate; TPAP=tetrapropylammonium perruthenate; NMO=*N*-methylmorpholine-*N*-oxide. Values in parentheses refer to the yields based on recovered starting material.

Scheme 60.
Computationally derived proposal of the mechanism.

Scheme 61.

“Cut-and-sew”-strategy enabling total synthesis of proposed structure of cycloinumakiol.

DIAD=diisopropyl azodicarboxylate.

Scheme 62.
Rhodium-catalyzed decarbonylative spirocyclization.

Scheme 63.

Substrate scope for decarbonylative spirocyclization.

selected examples

180b, 53%

180c, 93%

180d, 4%, no ZnCl_2 , 24 h
91%, 1 equiv ZnCl_2 , 24 h

180e, 0%, no ZnCl_2 , 24 h
64%, 1 equiv ZnCl_2 , 24 h

180f, 23%

180g, 40%

180h, 78%

180i, 56%

Scheme 64.

Catalytic direct intramolecular coupling of alkynyl benzocyclobutenone. Dppp=1,3-bis(diphenylphosphino)propane.

selected examples

Scheme 65.

Catalytic decarbonylative intramolecular coupling of alkynyl benzocyclobutenone. Values in parentheses refer to the yields based on recovered starting material.

Scheme 66.

Rhodium-catalyzed divergent syntheses of fused β -naphthol and indene scaffolds.

selected examples

Scheme 67.

Enantioselective rhodium-catalyzed carboacylation of C=N bonds. SDP=7,7'-bis(diphenylphosphino)-2,2',3,3'-tetrahydro-1,1'-spirobiindene.

Scheme 68.

Nickel-catalyzed intramolecular coupling with acetylenes and dienes.

Scheme 69.

Rhodium-catalyzed formation of [3.1.0]bicyclic structures.

selected examples

Scheme 70.

Rhodium-catalyzed [7+1] cycloaddition. $[\text{Si}]$ =trialkylsilyl.

Scheme 71.

Rhodium-catalyzed ring expansion of cyclobutenones.

Scheme 72.

Substrate scope and mechanism study for the cyclobutenone ring expansion.

Scheme 73.

Diels–Alder cycloadditions of 4,4-dimethylcyclobutenone.

Scheme 74.

Diels–Alder cycloaddition involving 3-cyanocyclobutenone.

O=C1C=CC2C1C=C3C2C(=O)C=C3

Chemical structure of a bicyclic ketone, specifically a phenyl-substituted bicyclo[2.2.2]octan-2-one derivative, with two phenyl groups attached to the bridgehead carbons.

CC1=C(C)C2C(C1)C(=O)C2