Published in final edited form as:

Arch Immunol Ther Exp (Warsz). 2016 December ; 64(6): 455–462. doi:10.1007/s00005-016-0403-0.

A Special Connection between $\gamma\delta$ T Cells and Natural Antibodies?

Willi K. Born^{1,2}, Yafei Huang³, Wanjiang Zeng⁴, Raul M. Torres², and Rebecca L. O'Brien^{1,2}

¹Department of Biomedical Research, National Jewish Health, Denver, CO 80206

²Department of Immunology and Microbiology, University of Colorado Health Sciences Center, Aurora, CO 80045

³Joint Laboratory for Stem Cell Engineering and Technology Transfer, School of Basic Medicine, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430030, P.R. China

⁴Department of Obstetrics and Gynecology, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, 430030, China

Abstract

Natural antibodies (NAbs) play an important role in early host defense, autophagy and tissue remodeling, and in immune regulation. They arise spontaneously (without specific immunization), and are already present at birth. NAbs are produced by B1 B cells, MZ B cells and other B cell types. They include all major Ig subclasses but IgM antibodies are prevalent, especially early in development. NAbs may be poly-specific, recognize particular auto-antigens, or detect neo-determinants such as those exposed during apoptosis or generated by oxidation. NAbs do not require cognate T cell help but depend on soluble mediators produced by T cells. Our recent studies suggest that $\gamma\delta$ T cells may have a special relationship with NAbs, and play a prominent role in their regulation, in part through the fine-tuning of IL-4-levels. The spontaneously activated state of these cells likely enables their cytokine production and other functions in the absence of external stimulation. Ontogenetically, the earlier arising $\gamma\delta$ T cells are better positioned than $\alpha\beta$ T cells to shape the developing repertoire of NAbs. Intriguingly, ligand specificities of NAbs and $\gamma\delta$ T cell receptors appear to be overlapping, perhaps allowing $\gamma\delta$ cognate help for certain NAb specificities. Via NAbs, $\gamma\delta$ T cells could exert a regulatory influence on numerous processes in health and disease.

Introduction

Prior to any immunization, circulating antibodies already exist in normal healthy humans and mice (Avrameas, 1991; Lutz et al., 2008). They are present at birth, and in mice have been shown to arise under germ-free conditions. These natural antibodies (NAbs) include all Ig subclasses. Natural IgM is prevalent particularly early in ontogeny and has been studied most extensively. B1 B cells appear to be the primary source of NAbs in mice (Savage and

Baumgarth, 2015), but other B cell-types, most notably marginal zone B cells (Durand et al., 2009), contribute as well (Avrameas and Selmi, 2013; Avrameas, 2016). Although sometimes referred to as “non-specific”, NAbS in fact may be mono- or poly-specific, and recognize auto-antigens, neo-antigens and certain foreign antigens. Importantly, recent studies revealed that NAbS play a critical role in the early host defense against pathogens, protection against malignancy, tissue homeostasis and immune regulation. Available information about NAbS and treatment opportunities with intravenous immunoglobulin (IVIG) has been expertly reviewed by others (Ehrenstein and Notley, 2010; Gronwall and Silvermann, 2014; Madi et al., 2012; McCoy et al., 2006; Panda and Ding, 2015; Rahyab et al., 2011; Schwartz-Albiez et al., 2009; Vas et al., 2013). It is only summarized here, followed by a discussion of new data suggesting that $\gamma\delta$ T cells become involved in regulating NAbS and their functional activity.

Specificity and function of natural antibodies

The primary B cell repertoire is not random (Perlmutter et al., 1985). Evidence for recurrent VH gene rearrangements in mice, humans and other vertebrates, at a time when the repertoire is not yet affected by antigenic selection, has been reported (reviewed in (Vas et al., 2013)). Use of microarray chips allowing the simultaneous detection of antibodies specific for up to 300 defined self antigens revealed that IgM repertoires in cord blood were very similar between individuals indicating that different humans are born with the same autoantibodies produced in utero regardless of variances in IgG autoantibodies found in their mothers (Madi et al., 2009). The mechanisms responsible for this consistency are not yet fully understood.

The early IgM antibodies in mice and humans are mostly germline-encoded and are produced by CD5⁺ B1a B cells (Savage and Baumgarth, 2015). They carry specificities for common bacterial Ags, auto-antigens, certain phospholipids, DNA and several cell membrane proteins. Table 1 lists some of the specificities of these NAbS (Air, 2015; Basnet et al., 2010; Bohn et al., 1994; Bovin, 2013; Buneva et al., 2013; Chen et al., 2009; Chikazawa et al., 2013; Chou et al., 2009; Durrbach et al., 2007; Fukuda et al., 2004; Hamanova et al., 2014; Hardy and Hayakawa, 2005; Kalyanaraman et al., 1982; Kulik et al., 2009; Lebon et al., 2011; Li et al., 2009; Llorente et al., 1999; Morales-Buenrostro et al., 2008; Posner et al., 1981; Robert-Guroff et al., 1982; Sauerborn et al., 2011; Shilova et al., 2011; Silvermann, 2015; Skurnik et al., 2012; Toth et al., 1984; Tsiantoulas et al., 2013; Tuominen et al., 2006; Turunen et al., 2012; Wang et al., 2013; Xu et al., 2013). Other important targets of NAbS include the Thomsen-Friedenreich tumor antigen (CD176) (Ulsemer et al., 2013), neural gangliosides relevant in Guillain-Barre syndrome (Boffey et al., 2004) and amyloid in Alzheimer’s disease (Kayed et al., 2011). Intriguingly, many of the autoreactive specificities are similarly found throughout evolution (Flajnik and Rumfelt, 2000; Gonzalez et al., 1988; Marchalonis et al., 1993). The non-random nature of the natural repertoire suggests that these early antibodies might be programmed to enable normal development, ensure essential functions and protect against common pathogens. Indeed, natural IgM has been characterized as protective in infections with Influenza virus, *Pseudomonas aeruginosa* and *Streptococcus pneumoniae* (reviewed in (Ehrenstein and Notley, 2010)). Natural IgM also affects B cell development. Mice genetically deficient in

surface (s) IgM exhibit changes in B cell subsets including B1a B cells and MZ B cells along with increased splenic and impaired peritoneal B cell survival (Baker and Ehrenstein, 2002; Notley et al., 2010). Deficiency in sIgM has also been linked to increases in autoimmunity and atherosclerosis in both humans and mouse models. Self-reactive IgE exacerbates interferon responses in autoimmunity (Henault et al., 2016). Figure 1 lists some of the known functions of NAb (Benatuil et al., 2005; Chen et al., 2009; Elluru et al., 2014; Matter and Ochsenbein, 2008; Panda et al., 2013; Pires et al., 2010; Rapaka et al., 2010; Stager et al., 2003; Veljkovic et al., 2010; Xu et al., 2008; Zabel et al., 2013), (McCoy et al., 2006), (Britschgi et al., 2009; Dimayuga et al., 2009; Elvington et al., 2012; Frostegard, 2010; Galkina and Ley, 2009; Gounopoulos et al., 2007; Gronwall et al., 2012; Shimomura et al., 2008; Sjoberg et al., 2009; Warrington and Rodrigues, 2010), (Aksentijevich et al., 1991; Lobo et al., 2014; Nogueira-Martins and Mariano, 2010; Warrington and Lewis, 2011).

Influence of T cells and cytokines

B cells are quite capable of producing antibodies in the absence of any T cell help, including isotypes that require class switch recombination. However, in our mouse facility, mice lacking all T cells (B6.TCR- $\beta^{-/-}/\delta^{-/-}$) had serum total Ig levels that reached only about 1/4 of those in wt mice. Levels of IgM and surprisingly, IgE antibodies were nearly normal, whereas IgA and IgG antibodies were significantly reduced, especially IgG1 (Huang et al., 2015). Cytokines produced by non-B cells can determine development and steady state levels of NAb. For example, IL-18 can boost production of protective natural IgM (reviewed in (Ehrenstein and Notley, 2010)). Natural IgE is also produced in the absence of MHCII cognate help but its production depends on IL-4 (McCoy et al., 2006). Their cytokine dependence subjects NAb to the non-specific regulatory influence of other cell types.

A connection between $\gamma\delta$ T cells and IgE production in non-immunized mice

Lymphocytes expressing $\gamma\delta$ TCRs ($\gamma\delta$ T cells) represent a lineage whose biological role remains poorly defined. Functionally, they resemble $\alpha\beta$ T cells (Bonneville et al., 2010) but in terms of specificity and mechanism of TCR-ligand-recognition, they share properties with B cells (Chien et al., 1996; Zeng et al., 2012). In several studies with mice carrying distinct mutations affecting TCR-signaling, increased non-immune IgE levels could be correlated with increases in $\gamma\delta$ T cells (Nunez-Cruz et al., 2003). In particular, this was observed in mice deficient in the Tec family tyrosine kinase Itk, which are unable to mount conventional IL-4- or IL-4 plus IL-13 producing T helper 2 (Th2) responses (Felices et al., 2009; Qi et al., 2009). Nevertheless, such mice had elevated levels of serum IgE and increased numbers of germinal center B cells, a phenotype found to be dependent on the presence of $\gamma\delta$ T cells. $\gamma\delta$ T cells in Itk $^{-/-}$ mice produced more Th2 cytokines and expressed at elevated levels co-stimulatory molecules important for B cell help, which suggested that they directly promote B cell activation and Ig class switching. In the same year, data generated in our lab with TCR-signaling competent mice revealed that IgE levels (both pre- and post-immunization)

are determined by the composition of $\gamma\delta$ T cells. Although mice deficient in all $\gamma\delta$ T cells had slightly reduced levels of serum IgE, others lacking only $V\gamma 4^+$ and $V\gamma 6^+$ subsets had vastly increased background IgE levels as well as exacerbated IgE responses following immunization (Huang et al., 2009). In these mice, $V\gamma 1^+$ and especially the NKT-like IL-4-producing $V\gamma 1/V\delta 6.3^+$ $\gamma\delta$ T cells (Gerber et al., 1999) were much increased, and so were serum levels of IL-4 (Huang et al., 2015). In fact, the genetically $\gamma\delta$ -deficient mice (B6.TCR- $V\gamma 4/6^{-/-}$) closely resembled $Itk^{-/-}$ mice. Together, these observations implied that the composition of $\gamma\delta$ T cells can play a critical role in setting steady state levels of IgE antibodies in non-immunized mice.

Effect of $\gamma\delta$ T cells on serum Ig levels in non-immunized mice

After these findings, we extended the analysis to other Ig classes (Huang et al., 2015). Our data suggest that the net effect of $\gamma\delta$ T cells on total serum Ig in normal non-immunized C57BL/6 mice is slightly enhancing as adult animals genetically deficient in all $\gamma\delta$ T cells (B6.TCR- $\delta^{-/-}$) had approximately 2-fold lower Ig levels. Net total Ig levels in these mice are lower mainly due to significantly reduced levels of IgG1 and IgG2b. Further insight came from the analysis of partially $\gamma\delta$ T cell-deficient mouse strains. Thus, mice lacking $V\gamma 1^+$ $\gamma\delta$ T cells exhibit total Ig levels that are even further diminished than those in B6.TCR- $\delta^{-/-}$ mice, and show much reduced levels of IgM, IgG3, IgG1 and IgG2b. IgG2c and IgA were diminished also but to a lesser degree. Hence, $V\gamma 1^+$ $\gamma\delta$ T cells appear to enhance the production of several Ig subclasses in non-immunized mice. On the other hand, mice deficient in $V\gamma 4^+$ and $V\gamma 6^+$ $\gamma\delta$ T cells exhibit much increased total Ig levels, mainly due to increases in IgM, IgG1, IgG2b and IgG2c. IgE antibodies are even more elevated but, due to their low concentration, contribute only little to overall increases in Ig levels. IgG3 and IgA are not affected. Hence, $V\gamma 4^+$ and perhaps $V\gamma 6^+$ $\gamma\delta$ T cells, two subsets in mice distinguished by their ability to produce IL-17 (O'Brien et al., 2009), appear to exert an inhibitory effect on most Ig production under steady state conditions. Our data suggest that this inhibition in non-immunized mice is mediated in large part through regulating $V\gamma 1^+$ $\gamma\delta$ T cells, their IL-4 production, and IL-4-dependent IL-4 production by $\alpha\beta$ T cells (Huang et al., 2015). Consistently, for some Ig subclasses, most obviously IgM, but also IgG2b and IgG2c, this regulation is still evident even in the absence of $\alpha\beta$ T cells. Finally, it has been shown that $V\gamma 1^+$ $\gamma\delta$ T cells kill tissue macrophages and thus resolve inflammatory immune responses (Carding and Egan, 2000; Dalton et al., 2003). As macrophages can be a source of IL-18, this cytotoxic activity might enable indirect control of IL-18-dependent antibody-types, at least in inflammation.

Comparing serum Ig levels in non-immunized adult wt C57BL/6 mice with mice lacking all $\gamma\delta$ T cells (B6.TCR- $\delta^{-/-}$) or all $\alpha\beta$ T cells (B6.TCR- $\beta^{-/-}$) initially showed that the absence of $\alpha\beta$ T cells affects levels of total serum Ig and most Ig subclasses more than does the absence of $\gamma\delta$ T cells. IgE antibodies in non-immunized mice were a notable exception: They were diminished in the absence of all $\gamma\delta$ T cells but not in the absence of all $\alpha\beta$ T cells (Huang et al., 2015). However, mice lacking only subsets of $\gamma\delta$ T cells revealed far greater changes in Ig levels, suggestive of potent enhancing and suppressive regulation by these cells (Huang et al., 2015). This powerful regulation is all the more noteworthy

considering that $\gamma\delta$ T cells in circulation and in the lymphoid tissues represent a much smaller cell population than $\alpha\beta$ T cells.

$\gamma\delta$ T cells affect the repertoire of antibodies in non-immunized mice

The observation that changes in $\gamma\delta$ T cells differentially affect levels of Ig subclasses in non-immunized mice suggests an influence on the repertoire and specificities of NAb. To explore this possibility, we examined antibodies specific for chromatin, DNA and nuclear antigens in non-immunized mice (Huang et al., 2015). We found that the same change in $\gamma\delta$ T cells that increased all Ig subclasses except IgG3 and IgA - a genetic deficiency in $V\gamma4+$ and $V\gamma6+$ $\gamma\delta$ T cells, also led to substantial increases in anti chromatin antibodies. However, the anti chromatin antibodies were still further increased relative to the elevated total Ig levels, indicating a change in the composition of the elevated antibodies, and in their specificity. We also found a clear increase in antibodies detecting dsDNA/histone complexes and smaller increases in antibodies specific for ssDNA, and anti-nuclear autoantibodies. Finally, cell transfer and depletion experiments showed the change in antibody composition/specificity to be driven by the dys-regulated $V\gamma1+$ $\gamma\delta$ T cells derived from non-immunized B6.TCR- $V\gamma4/6-/-$ mice. Hence, $\gamma\delta$ T cells affect the antibody repertoire in non-immunized mice (Huang et al., 2015).

Do $\gamma\delta$ T cells have a special connection with natural antibodies?

The above-mentioned observations clearly implicate $\gamma\delta$ T cells in the regulation of antibodies independently of immunization, and thus establish their involvement with NAb. However, it remains to be determined if $\gamma\delta$ T cells have a unique functional connection with NAb. Do they play a role distinct from that of other T cells, in the development and maintenance of these antibodies? Such a role might even help to explain the evolutionary conservation of $\gamma\delta$ T cells. We suggest that such a special connection is probable, for the following reasons: Firstly, unlike the majority of unstimulated $\alpha\beta$ T cells, which are antigenically naïve and metabolically inactive, many peripheral $\gamma\delta$ T cells in non-immunized mice already exist in a state of moderate activation (Tough and Sprent, 1998). This activated/memory phenotype could support background levels of cytokine production capable of sustaining some B cell differentiation and spontaneous antibody production. Secondly, in mouse ontogeny, $\gamma\delta$ T cells develop roughly at the same time as the earliest B cells, and for a short period of time, the fetal liver gives rise to precursors of both cell-types (Havran and Allison, 1988; Whitlock et al., 1985). $\alpha\beta$ T helper cells, on the other hand, arise several days later (Marrack et al., 1988). Moreover, $\gamma\delta$ thymus-emigrants are functionally more mature than $\alpha\beta$ thymus-emigrants (Jin et al., 2009; Narayan et al., 2012). In particular, $\gamma\delta$ thymocytes are already sufficiently differentiated to produce polarized cytokines including IL-4 and IL-13 (Jin et al., 2009). Thus, $\gamma\delta$ T cells are enabled to support and modulate B cell development and antibody production from the start, by supplying cytokines prior to any immunization. In addition, $\gamma\delta$ T cells might interact with B cells via ligands expressed on B cells such as members of the CD1 family of molecules (Delia et al., 1988; Duan et al., 2008; Sonoda and Stein-Streilein, 2002), which are recognized by $\gamma\delta$ T cells (Dieude et al., 2011; Porcelli et al., 1992; Russano et al., 2006). Lastly, one might consider the early idea that B1 B cells and $\gamma\delta$ T cells occupy equivalent positions in a

“layered immune system” (Herzenberg and Herzenberg, 1989). We actually suspect that there is resemblance or overlap between antigen specificities of NAb and $\gamma\delta$ T cells. Thus, both recognize poly-anionic ligands including certain synthetic polypeptides (Cady et al., 2000) but also DNA/protein (unpublished data) and phospholipid/protein complexes (Born et al., 2003; Dieude et al., 2011; Lafer et al., 1981), and recent studies suggest a shared specificity for insulin (Aydintug et al., 2014; Liu et al., 2002; Zhang et al., 2010). Thus, in addition to non-specific cytokine support, at least some $\gamma\delta$ T cells conceivably could provide help to B cells via cognate interactions between $\gamma\delta$ TCR, antigen and BCR. Hence, the extent to which antigen specificities of $\gamma\delta$ TCRs and NAb might be related in non-immunized mice would be revealing with regard to the biological role of $\gamma\delta$ T cells.

Modulating NAb and their contribution to immune responsiveness via $\gamma\delta$ T cells?

Although B cells can spontaneously produce antibodies in the absence of all T cells, T cells affect antibody production when they are present. The assumption that $\gamma\delta$ T cells modulate levels and repertoire of NAb invites additional points: First, $\gamma\delta$ T cells in humans undergo large changes in population size and composition during ontogeny (Parker et al., 1990), due to genetic variation, in the course of hematopoietic transplantation (Airoldi et al., 2015), and as a consequence of diseases (Bank and Marcu-Malina, 2013; Pauza et al., 2015). Thus, one might expect based on the findings in mice that such changes in human $\gamma\delta$ T cells have an effect on human B cells and their natural production of immunoglobulins: Large variation of NAb might occur as a consequence of the changes in human $\gamma\delta$ T cells. Furthermore, similar to the mice, specific targeting of the relevant human $\gamma\delta$ T cells, directly or through their regulators, might effectively change human NAb, complement IVIG in certain cases, and at times might be preferable over targeting all B cells (e.g. using antibody drugs against CD19) or entire antibody classes (e.g. antibody drugs against IgE).

Acknowledgments

We would like to thank Drs. Lawrence Wysocki and Thiago Detanico for helpful discussions.

Abbreviations

NAb	natural antibody
TCR	T cell receptor
IL-4	interleukin 4

References

- Air GM. Influenza virus antigenicity and broadly neutralizing epitopes. *Curr Opin Virol.* 2015; 11:113–121. [PubMed: 25846699]
- Airoldi I, Bertaina A, Prigione I, et al. Gammadelta T cell reconstitution after HLA-haploidentical hematopoietic transplantation depleted of TCR-alpha-beta+/CD19+ lymphocytes. *Blood.* 2015; 125:2349–2358. [PubMed: 25612623]

- Aksentijevich I, Sachs DH, Sykes M. Natural antibodies can inhibit bone marrow engraftment in the rat — mouse species combination. *J Immunol.* 1991; 147:4140–4146. [PubMed: 1753089]
- Avrameas S. Natural autoantibodies: from “horror autotoxicus” to “gnoti seauton”. *Immunology Today.* 1991; 12:154–159. [PubMed: 1715166]
- Avrameas S, Selmi C. Natural autoantibodies in the physiology and pathophysiology of the immune system. *J Autoimmun.* 2013; 41:46–49. [PubMed: 23384670]
- Avrameas SC. Autopolyreactivity confers a holistic role in the immune system. *Scand J Immunol.* 2016; doi: 10.1111/sji12414
- Aydintug MK, Zhang L, Wang C, et al. Gammadelta T cells recognize the insulin B:9-23 peptide antigen when it is dimerized through thiol oxidation. *Molecular Immunology.* 2014; 60:116–128. [PubMed: 24853397]
- Baker N, Ehrenstein MR. Cutting edge: selection of B lymphocyte subsets is regulated by natural IgM. *J Immunol.* 2002; 169:6686–6690. [PubMed: 12471099]
- Bank I, Marcu-Malina V. Quantitative peripheral blood perturbations of gammadelta T cells in human disease and their clinical implications. *Clinic Rev Allerg Immunol.* 2013; doi: 10.1007/s12016-013-8391-x
- Basnet NB, Ide K, Tahara H, et al. Deficiency of N-glycolylneuraminic acid and Galalpha1-3Galbeta1-4GlcNAc epitopes in xenogeneic cells attenuates cytotoxicity of human natural antibodies. *Xenotransplantation.* 2010; 17:440–448. [PubMed: 21158945]
- Benatuil L, Kaye JA, Rich RF, et al. The influence of natural antibody specificity on antigen immunogenicity. *European Journal of Immunology.* 2005; 35:2638–2647. [PubMed: 16082726]
- Boffey J, Nicholl D, Wagner ER, et al. Innate murine B cells produce anti-disialosyl antibodies reactive with *Campylobacter jejuni* LPS and gangliosides that are polyreactive and encoded by a restricted set of unmutated V genes. *J Neuroimmunol.* 2004; 152:98–111. [PubMed: 15223242]
- Bohn J, Roggenbuck D, Settmacher U, et al. Binding of natural human IgM auto-antibodies to human tumor cell lines and stimulated normal T lymphocytes. *Immunol Lett.* 1994; 39:187–194. [PubMed: 7912225]
- Bonneville M, O'Brien RL, Born WK. Gammadelta T cell effector functions: a blend of innate programming and acquired plasticity. *Nature Reviews Immunology.* 2010; 10:467–478.
- Born WK, Vollmer M, Reardon C, et al. Hybridomas expressing gammadelta T-cell receptors respond to cardiolipin and beta2-glycoprotein 1 (apolipoprotein H). *Scand J Immunol.* 2003; 58:374–381. [PubMed: 12950685]
- Bovin NV. Natural antibodies to glycans. *Biochemistry (Mosc).* 2013; 78:786–797. [PubMed: 24010841]
- Britschgi M, Olin CE, Johns HT, et al. Neuroprotective natural antibodies to assemblies of amyloidogenic peptides decrease with normal aging and advancing Alzheimer's disease. *Proceedings of the National Academy of Sciences of the United States of America.* 2009; 106:12145–12150. [PubMed: 19581601]
- Buneva VN, Krasnorutskii MA, Nevinsky GA. Natural antibodies to nucleic acids. *Biochemistry (Mosc).* 2013; 78:127–143. [PubMed: 23581984]
- Cady CT, Lahn M, Vollmer M, et al. Response of murine $\gamma\delta$ T cells to the synthetic polypeptide poly-Glu⁵⁰Tyr⁵⁰. *J Immunology.* 2000; 165:1790–1798. [PubMed: 10925256]
- Carding SR, Egan PJ. The importance of $\gamma\delta$ T cells in the resolution of pathogen-induced inflammatory immune responses. *Immunological Reviews.* 2000; 173:98–108. [PubMed: 10719671]
- Chen Y, Park Y-B, Patel E, et al. IgM antibodies to apoptosis-associated determinants recruit C1q and enhance dendritic cell phagocytosis of apoptotic cells. *J Immunol.* 2009; 182:6031–6043. [PubMed: 19414754]
- Chien Y-H, Jores R, Crowley MP. Recognition by $\gamma\delta$ T cells. *Annu Rev Immunol.* 1996; 14:511–532. [PubMed: 8717523]
- Chikazawa M, Otaki N, Shibata T, et al. An apoptosis-associated mammary protein deficiency leads to an enhanced production of IgM antibodies against multiple damage-associated molecules. *PLoS One.* 2013; 8:e68468. Print 0062013. doi: 10.61371/journal.pone.0068468 [PubMed: 23874637]

- Chou MY, Fogelstrand L, Hartvigsen K, et al. Oxidation-specific epitopes are dominant targets of innate natural antibodies in mice and humans. *J Clin Invest*. 2009; 119:1335–1349. [PubMed: 19363291]
- Dalton JE, Pearson J, Scott P, et al. The interaction of gamma delta T cells with activated macrophages is a property of the Vg1 subset. *J Immunol*. 2003; 171:6488–6494. [PubMed: 14662848]
- Delia D, Cattoretti G, Fontanella E, et al. CD1c but neither CD1a nor CD1b molecules are expressed on normal, activated, and malignant human B cells: identification of a new B-cell subset. *Blood*. 1988; 72:241–247. [PubMed: 3260523]
- Dieude M, Striegl H, Tyznik AJ, et al. Cardiolipin binds to CD1d and stimulates CD1d-restricted gammadelta T cells in the normal murine repertoire. *J Immunol*. 2011; 186:4771–4781. [PubMed: 21389252]
- Dimayuga PC, Cesena FH, Chyu KY, et al. Natural antibodies and complement modulate intimal thickening after arterial injury. *Am J Physiol Regul Integr Comp Physiol*. 2009; 297:R1593–15600. [PubMed: 19776252]
- Duan B, Niu H, Xu Z, et al. Intrafollicular location of marginal zone/CD1d(hi) B cells is associated with autoimmune pathology in a mouse model of lupus. *Lab Invest*. 2008; 88:1008–1020. [PubMed: 18607347]
- Durand CA, Hartvigsen K, Fogelstrand L, et al. Phosphoinositide 3-kinase p110 delta regulates natural antibody production, marginal zone and B-1 B cell function, and autoantibody responses. *J Immunol*. 2009; 183:5673–5684. [PubMed: 19843950]
- Durrbach A, Baple E, Preece AF, et al. Virus recognition by specific natural antibodies and complement results in MHC I cross-presentation. *European Journal of Immunology*. 2007; 37:1254–1265. [PubMed: 17407191]
- Ehrenstein MR, Notley CA. The importance of natural IgM: scavenger, protector and regulator. *Nature Reviews Immunology* Epub ahead of print. 2010
- Elluru SR, Kaveri SV, Bayry J. Regulation of human dendritic cell function by natural anti-CD40 antibodies. *Methods Mol Biol*. 2014; 1155:47–54. [PubMed: 24788172]
- Elvington A, Atkinson C, Kulik L, et al. Pathogenic natural antibodies propagate cerebral injury following ischemic stroke in mice. *J Immunol*. 2012; 188:1460–1468. [PubMed: 22198950]
- Felices M, Yin CC, Kosaka Y, et al. Tec kinase Itk in gammadelta T cells is pivotal for controlling IgE production in vivo. *Proc Natl Acad Sci (USA)*. 2009; 106:8308–8313. [PubMed: 19416854]
- Flajnik MF, Rummel LL. Early and natural antibodies in non-mammalian vertebrates. *Current Topics in Microbiology and Immunology*. 2000; 252:233–240. [PubMed: 11125480]
- Frostegard J. Low level natural antibodies against phosphorylcholine: a novel risk marker and potential mechanism in atherosclerosis and cardiovascular disease. *Clin Immunol*. 2010; 134:47–54. [PubMed: 19748321]
- Fukuda K, Takao Y, Miyazaki Y, et al. New type of natural antibodies reactive to cytotoxic T lymphocyte-directed cancer vaccine peptides. *Immunobiology*. 2004; 209:245–253. [PubMed: 15518336]
- Galkina E, Ley K. Immune and inflammatory mechanisms of atherosclerosis (*). *Annu Rev Immunol*. 2009; 27:165–197. [PubMed: 19302038]
- Gerber DJ, Azuara V, Levraud J-P, et al. IL-4-producing $\gamma\delta$ T cells that express a very restricted TCR repertoire are preferentially localized in liver and spleen. *J Immunol*. 1999; 163:3076–3082. [PubMed: 10477572]
- Gonzalez R, Charlemagne J, Mahana W, et al. Specificity of natural serum antibodies present in phylogenetically distinct fish species. *Immunology*. 1988; 63:31–36. [PubMed: 3338819]
- Gounopoulos P, Merki E, Hansen LF, et al. Antibodies to oxidized low density lipoprotein: epidemiological studies and potential clinical applications in cardiovascular disease. *Minerva Cardioangiol*. 2007; 55:821–837. [PubMed: 18091649]
- Gronwall C, Silvermann GJ. Natural IgM: Beneficial autoantibodies for the control of inflammatory and autoimmune disease. *J Clin Immunol*. 2014; 34(Suppl 1):S12–S21. [PubMed: 24691998]
- Gronwall C, Vas J, Silverman GJ. Protective roles of natural IgM antibodies. *Front Immunol*. 2012

- Hamanova M, Zdrzilova Dubska L, Valik D, et al. Natural antibodies against alpha(1,3) galactosyl epitope in the serum of cancer patients (article in Czech). *Epidemiol Mikrobiol Imunol*. 2014; 63:130–133. [PubMed: 25025679]
- Hardy RR, Hayakawa K. Development of B cells producing natural antibodies to thymocytes and senescent erythrocytes. *Springer Semin Immunopathol*. 2005; 26:363–375. [PubMed: 15611857]
- Havran W, Allison JP. Developmentally ordered appearance of thymocytes expressing different T cell antigen receptors. *Nature*. 1988; 335:443–445. [PubMed: 2458531]
- Henault J, Riggs JM, Karnell JL, et al. Self-reactive IgE exacerbates interferon responses associated with autoimmunity. *Nature Immunol*. 2016; 17:196–203. [PubMed: 26692173]
- Herzenberg LA, Herzenberg LA. Toward a layered immune system. *Cell*. 1989; 59:953–954. [PubMed: 2688900]
- Huang Y, Heiser RA, Detanico TO, et al. Gammadelta T cells affect IL-4 production and B-cell tolerance. *Proceedings of the National Academy of Sciences of the United States of America*. 2015; 112:E39–E48. [PubMed: 25535377]
- Huang Y, Jin N, Roark CL, et al. The influence of IgE-enhancing and IgE-suppressive gammadelta T cells changes with exposure to inhaled ovalbumin. *J Immunol*. 2009; 183:849–855. [PubMed: 19542369]
- Jin N, Roark CL, Miyahara N, et al. Allergic airway hyperresponsiveness-enhancing gammadelta T cells develop in normal untreated mice and fail to produce IL-4/13, unlike Th2 and NKT cells. *J Immunol*. 2009; 182:2002–2010. [PubMed: 19201853]
- Kalyanaraman VS, Sarngadharan MG, Nakao Y, et al. Natural antibodies to the structural core protein (p24) of the human T-cell leukemia (lymphoma) retrovirus found in sera of leukemia patients in Japan. *Proceedings of the National Academy of Sciences of the United States of America*. 1982; 79:1653–1657. [PubMed: 6951204]
- Kayed R, Jackson GR, Estes DM, et al. Alzheimer's disease: review of emerging treatment role for intravenous immunoglobulins. *J Cent Nerv Syst Dis*. 2011; 3:67–73. [PubMed: 23861639]
- Kulik L, Fleming SD, Moratz C, et al. Pathogenic natural antibodies recognizing annexin IV are required to develop intestinal ischemia-reperfusion injury. *J Immunol*. 2009; 182:5363–5373. [PubMed: 19380783]
- Lafer EM, Rauch J, Andrzejewski C Jr, et al. Polyspecific monoclonal lupus autoantibodies reactive with both polynucleotides and phospholipids. *Journal of Experimental Medicine*. 1981; 153:897–909. [PubMed: 6972993]
- Lebon A, Verkaik NJ, Labout JA, et al. Natural antibodies against several pneumococcal virulence proteins in children during the pre-pneumococcal-vaccine era: the generation R study. *Infect Immun*. 2011; 79:1680–1687. [PubMed: 21282409]
- Li S, Waer M, Billiau AD. Xenotransplantation: role of natural immunity. *Transpl Immunol*. 2009; 21:70–74. [PubMed: 18992342]
- Liu E, Moriyama H, Abiru N, et al. Anti-peptide autoantibodies and fatal anaphylaxis in NOD mice in response to insulin self-peptides B:9-23 and B:13-23. *J Clin Invest*. 2002; 110:1021–1027. [PubMed: 12370280]
- Llorente M, Sanchez-Palomino S, Manes S, et al. Natural human antibodies retrieved by phage display libraries from healthy donors: polyreactivity and recognition of human immunodeficiency virus 1 gp120 epitopes. *Scand J Immunol*. 1999; 50:270–279. [PubMed: 10447936]
- Lobo PI, Brayman KL, Okusa MD. Natural IgM anti-leukocyte autoantibodies (IgM-ALA) regulate inflammation induced by innate and adaptive immune mechanisms. *J Clin Immunol*. 2014; 34(Suppl 1):S22–29. [PubMed: 24711004]
- Lutz HU, Binder CJ, Kaveri S. Naturally occurring auto-antibodies in homeostasis and disease. *Trends in Immunol*. 2008; 30:43–51. [PubMed: 19058756]
- Madi A, Bransburg-Zabary S, Kenett DY, et al. The natural autoantibody repertoire in newborns and adults: a current overview. *Adv Exp Med Biol*. 2012; 750:198–212. [PubMed: 22903676]
- Madi A, Hecht I, Bransburg-Zabary S, et al. Organization of the autoantibody repertoire in healthy newborns and adults revealed by system level informatics of antigen microarray data. *Proceedings of the National Academy of Sciences of the United States of America*. 2009; 106:14484–14489. [PubMed: 19667184]

- Marchalonis JJ, Hohman VS, Thomas C, et al. Antibody production in sharks and humans: a role for natural antibodies. *Dev Comp Immunol*. 1993; 17:41–53. [PubMed: 8449250]
- Marrack P, Kuschner E, Born W, et al. The development of helper T cell precursors in mouse thymus. *J Immunol*. 1988; 140:2508–2514. [PubMed: 2965721]
- Matter MS, Ochsenbein AF. Natural antibodies target virus-antibody complexes to organized lymphoid tissue. *Autoimmun Rev*. 2008; 7:480–486. [PubMed: 18558366]
- McCoy KD, Harris NL, Diener P, et al. Natural IgE production in the absence of MHC class II cognate help. *Immunity*. 2006; 24:329–339. [PubMed: 16546101]
- Morales-Buenrostro LE, Terasaki PI, Marino-Vazquez LA, et al. “Natural” human leukocyte antigen antibodies found in nonalloimmunized healthy males. *Transplantation*. 2008; 86:1111–1115. [PubMed: 18946350]
- Narayan K, Sylvia KE, Malhotra N, et al. Intrathymic programming of effector fates in three molecularly distinct gammadelta T cell subtypes. *Nature Immunol* published online April. 2012; 1:2012.doi: 10.1038/ni.2247
- Nogueira-Martins MF, Mariano M. B-1 cell participation in T-cell-mediated alloimmune response. *Immunobiology*. 2010; 215:264–274. [PubMed: 19581018]
- Notley CA, Baker N, Ehrenstein MR. Secreted IgM enhances B cell receptor signaling and promotes splenic but impairs peritoneal B cell survival. *J Immunol*. 2010; 184:3386–3393. [PubMed: 20207998]
- Nunez-Cruz S, Aguado E, Richelme S, et al. LAT regulates gammadelta T cell homeostasis and differentiation. *Nature Immunology*. 2003; 4:999–1008. [PubMed: 12970761]
- O’Brien RL, Roark CL, Born WK. IL-17-producing gammadelta T cells. *European Journal of Immunology*. 2009; 39:662–666. [PubMed: 19283718]
- Panda S, Ding JL. Natural antibodies bridge innate and adaptive immunity. *J Immunol*. 2015; 194:13–20. [PubMed: 25527792]
- Panda S, Zhang J, Tan NS, et al. Natural IgG antibodies provide innate protection against ficolin-opsonized bacteria. *EMBO Journal*. 2013; 32:2905–2919. [PubMed: 24002211]
- Parker CM, Groh V, Band H, et al. Evidence for extrathymic changes in the T cell receptor γ/δ repertoire. *J Exp Med*. 1990; 171:1597–1612. [PubMed: 2185330]
- Pauza CD, Poonia B, Li H, et al. GammaDelta T cells in HIV disease: past, present and future. *Frontiers in Immunology*. 2015; 5:1–11.
- Perlmutter RM, Kearney JF, Chang SP, et al. Developmentally controlled expression of immunoglobulin V_H genes. *Science*. 1985; 227:1597–1601. [PubMed: 3975629]
- Pires AE, Afonso AF, Queiros A, et al. Treatment with polyclonal immunoglobulin during T-cell reconstitution promotes naive T-cell proliferation. *J Immunother*. 2010; 33:618–625. [PubMed: 20551835]
- Porcelli S, Morita CT, Brenner MB. CD1b restricts the response of human CD4⁺ T lymphocytes to a microbial antigen. *Nature*. 1992; 360:593–597. [PubMed: 1281285]
- Posner LE, Robert-Guroff M, Kalyanaraman VS, et al. Natural antibodies to the human T cell lymphoma virus in patients with cutaneous T cell lymphomas. *Journal of Experimental Medicine*. 1981; 154:333–346. [PubMed: 6973601]
- Qi Q, Xia M, Hu J, et al. Enhanced development of CD4⁺ gammadelta T cells in the absence of Itk results in elevated IgE production. *Blood*. 2009; 114:564–571. [PubMed: 19443662]
- Rahyab AS, Alam A, Kapoor A, et al. Natural antibody - Biochemistry and functions. *Glob J Biochem*. 2011; 2:283–288. [PubMed: 25309852]
- Rapaka RR, Ricks DM, Alcorn JF, et al. Conserved natural IgM antibodies mediate innate and adaptive immunity against the opportunistic fungus *Pneumocystis murina*. *Journal of Experimental Medicine*. 2010; 207:2907–2919. [PubMed: 21149550]
- Robert-Guroff M, Fahey KA, Maeda M, et al. Identification of HTLV p19 specific natural human antibodies by competition with monoclonal antibody. *Virology*. 1982; 122:297–305. [PubMed: 6293176]

- Russano AM, Agea E, Corazzi L, et al. Recognition of pollen-derived phosphatidyl-ethanolamine by human CD1d-restricted gamma delta T cells. *J Allergy Clin Immunol*. 2006; 117:1178–1184. [PubMed: 16675349]
- Sauerborn M, van de Vosse E, Delawi D, et al. Natural antibodies against bone morphogenic proteins and interferons in healthy donors and in patients with infections linked to type-1 cytokine responses. *J Interferon Cytokine Res*. 2011; 31:661–669. [PubMed: 21612442]
- Savage HP, Baumgarth N. Characteristics of natural antibody-secreting cells. *Ann N Y Acad Sci*. 2015; 1362:132–142. [PubMed: 26104151]
- Schwartz-Albiez R, Monteiro RC, Rodriguez M, et al. Natural antibodies, intravenous immunoglobulin and their role in autoimmunity, cancer and inflammation. *Clin Exp Immunol*. 2009; 158(Suppl 1): 43–50. [PubMed: 19883423]
- Shilova NV, Navakouski MJ, Huflejt M, et al. Changes in the repertoire of natural antibodies caused by immunization with bacterial antigens. *Biochemistry (Mosc)*. 2011; 76:862–866. [PubMed: 21999548]
- Shimomura Y, Mizoguchi E, Sugimoto K, et al. Regulatory role of B-1 B cells in chronic colitis. *Int Immunol*. 2008; 20:729–737. [PubMed: 18375938]
- Silvermann GJ. Protective natural antibodies to apoptotic cells: evidence of convergent selection of recurrent innate-like clones. *Ann N Y Acad Sci*. 2015; 1362:164–175. [PubMed: 25990717]
- Sjoberg BG, Su J, Dahlbom I, et al. Low levels of IgM antibodies against phosphorylcholine-A potential risk marker for ischemic stroke in men. *Atherosclerosis*. 2009; 203:528–532. [PubMed: 18809177]
- Skurnik D, Kropec A, Roux D, et al. Natural antibodies in normal human serum inhibit *Staphylococcus aureus* capsular polysaccharide vaccine efficacy. *Clin Infect Dis*. 2012; 55:1188–1197. [PubMed: 22806596]
- Sonoda KH, Stein-Streilein J. CD1d on antigen-transporting APC and splenic marginal zone B cells promotes NKT cell-dependent tolerance. *European Journal of Immunology*. 2002; 32:848–857. [PubMed: 11870629]
- Stager S, Alexander J, Kirby AC, et al. Natural antibodies and complement are endogenous adjuvants for vaccine-induced CD8+ T-cell responses. *Nature Medicine*. 2003; 9:1287–1292.
- Toth FD, Szabo B, Vaczi L, et al. Natural antibodies to simian type-C viruses and human retrovirus HTLV in patients with lymphoid malignancies. *Acta Microbiol Hung*. 1984; 31:387–392. [PubMed: 6100223]
- Tough DF, Sprent J. Lifespan of γ/δ T cells. *Journal of Experimental Medicine*. 1998; 187:357–365. [PubMed: 9449716]
- Tsiantoulas D, Gruber S, Binder CJ. B-1 cell immunoglobulin directed against oxidation-specific epitopes. *Front Immunol*. 2013; 3:415. eCollection 02012. doi: 10.3389/fimmu.2012.00415 [PubMed: 23316200]
- Tuominen A, Miller YI, Hansen LF, et al. A natural antibody to oxidized cardiolipin binds to oxidized low-density lipoprotein, apoptotic cells, and atherosclerotic lesions. *Arterioscler Thromb Vasc Biol*. 2006; 26:2096–2102. [PubMed: 16794225]
- Turunen SP, Kumm O, Harila K, et al. Recognition of *Porphyromonas gingivalis* gingipain epitopes by natural IgM binding to malondialdehyde modified low-density lipoprotein. *PLoS One*. 2012; 7:e34910. Epub 0032012 Apr 0034915. doi: 10.1371/journal.pone.0034919 [PubMed: 22496875]
- Ulsemer P, Henderson G, Toutounian K, et al. Specific humoral immune response to the Thomsen-Friedenreich tumor antigen (CD176) in mice after vaccination with the commensal bacterium *Bacteroides ovatus* D-6. *Cancer Immunol Immunother*. 2013; 62:875–887. [PubMed: 23381581]
- Vas J, Gronwall C, Silverman GJ. Fundamental roles of the innate-like repertoire of natural antibodies in immune homeostasis. *Frontiers in Immunology*. 2013 Feb 5.4(4):4. eCollection 02013. doi: 10.3389/fimmu.2013.00004 [PubMed: 23386848]
- Veljkovic M, Dopsaj V, Stringer WW, et al. Aerobic exercise training as a potential source of natural antibodies protective against human immunodeficiency virus-1. *Scand J Med Sci Sports*. 2010; 20:469–474. [PubMed: 19538533]

- Wang C, Turunen SP, Kumm O, et al. Natural antibodies of newborns recognize oxidative stress-related malondialdehyde acetaldehyde adducts on apoptotic cells and atherosclerotic plaques. *International Immunology*. 2013; 25:575–587. [PubMed: 23900424]
- Warrington AE, Rodrigues M. Method of identifying natural antibodies for remyelination. *J Clin Immunol*. 2010; 30(Suppl 1):S50–S55. [PubMed: 20387101]
- Warrington RJ, Lewis KE. Natural antibodies against nerve growth factor inhibit in vitro prostate cancer cell metastasis. *Cancer Immunol Immunother*. 2011; 60:187–195. [PubMed: 20976447]
- Whitlock C, Denis K, Robertson D, et al. In vitro analysis of murine B-cell development. *Annu Rev Immunol*. 1985; 3:213–235. [PubMed: 3904770]
- Xu Z, Qiu Q, Tian J, et al. Coagulation factor X shields adenovirus type 5 from attack by natural antibodies and complement. *Nature Medicine*. 2013; 19:452–457.
- Xu Z, Tian J, Smith JS, et al. Clearance of adenovirus by Kupffer cells is mediated by scavenger receptors, natural antibodies, and complement. *J Virol*. 2008; 82:11705–11713. [PubMed: 18815305]
- Zabel F, Kundig TM, Bachmann MF. Virus-induced humoral immunity: on how B cell responses are initiated. *Curr Opin Virol*. 2013; 3:367–362.
- Zeng X, Wei YL, Huang J, et al. Gammadelta T cells recognize a microbial encoded B cell antigen to initiate a rapid antigen-specific interleukin-17 response. *Immunity*. 2012; 37:524–534. [PubMed: 22960222]
- Zhang L, Jin N, Nakayama M, et al. Gamma delta T cell receptors confer autonomous responsiveness to the insulin-peptide B:9–23. *J Autoimmun*. 2010; 34:478–484. [PubMed: 20080388]

Figure 1. Diverse functions of natural antibodies
 Marked in red: NAb with pathogenic effect.

Table 1
Broad Target Range of Natural Antibodies

Reference numbers in the table refer to list of references following the main body of the text. Marked in red: NAbs with pathogenic effect.

Category	Antigens recognized	species	References
Pathogen-derived	Poly-N-acetyl glucosamine (Staph. aureus capsule)	human	Skurnik et al. 2012
	Porphyromonas gingivalis gingipain		Turunen et al. 2012
	Pneumococcal virulence proteins		Lebon et al. 2011
	Adenovirus type 5	mouse	Xu et al. 2013
	HIV	human	Llorente et al. 1999
	HTLV, simian type C virus p24 structural core protein p19		Toth et al. 1984 Kalyanaraman et al. 1982 Robert-Guroff et al. 1982 Posner et al. 1981
	Influenza		Air 2015
	Measles virus		Durrbach et al. 2007
autologous	Apoptotic cells including thymocytes and senescent erythrocytes and their antigens phosphorylcholine	mouse, human	Chou et al. 2009 Silverman et al. 2015 Chikazawa et al. 2013 Hardy et al. 2005 Chen et al. 2009
	Tumor cells, activated T cells	human	Bohn et al. 1994
	Proliferation-related self peptides		Fukuda et al. 2004
	Annexin IV	mouse, human	Kulik et al. 2009
	Nucleic acids	human	Buneva et al. 2013 Henault et al. 2016
	Glycans, sialoglycans	chicken, human	Shilova et al. 2011 Bovin et al. 2013
	Bone morphogenic protein	human	Sauerborn et al. 2011
	IFN- β , IFN- γ		Sauerborn et al. 2011
	α (1,3)-galactosyl		Hamanova et al. 2014
	CD40		Elluru et al. 2014
allogeneic	HLA		Morales-Buenrostro et al. 2008
xenogenic	α Gal		Li et al. 2009
	N-glycolylneuraminic acid	mouse	Basnet et al. 2010
Oxidation-induced	Apoptotic cells	mouse, human	Chou et al. 2009
	Oxidized lipids, cardiolipin	mouse	Tsiantoulas et al. 2013 Tuominen et al. 2006
	Malondialdehyde acetaldehyde adducts	human	Wang et al. 2013