Published in final edited form as:

J Nat Prod. 2006 May ; 69(5): 845–846. doi:10.1021/np050554l.

Aegicerin, the First Oleanane Triterpene with Potent Antimycobacterial Activity, Isolated from *Clavija procera*

Rosario Rojas^{†,‡}, Luz Caviedes[‡], José C. Aponte[§], Abraham J. Vaisberg[‡], Walter H. Lewis^{||}, Gerardo Lamas[⊥], César Sarasara[∇], Robert H. Gilman^{○,‡}, and Gerald B. Hammond^{*,§,‡}

[†]Departamento de Ciencias Farmacéuticas, Universidad Peruana Cayetano Heredia, Lima, Perú

[‡]Departamento de Microbiología y Laboratorios de Investigación y Desarrollo, Universidad Peruana Cayetano Heredia

[§]Department of Chemistry, University of Louisville, Louisville, Kentucky, 40292

^{||}Department of Biology, Washington University, St. Louis MO 63130

[⊥]Departamento de Entomología, Museo de Historia Natural Universidad Nacional Mayor de San Marcos, Lima, Perú

[∇]Confederación de Nacionalidades Amazónicas del Perú (CONAP), Lima, Perú

[○]Department of International Health, Johns Hopkins School of Public Health Baltimore, Maryland, 21205

Abstract

An ethanol extract of the Peruvian plant *Clavija procera*, a member of the rare Theophrastaceae family, was fractionated using a colorimetric bioassay-guided protocol against *Mycobacterium tuberculosis* (MTB), yielding the oleanane triterpenoid aegicerin (1) as the active constituent. Its MIC values ranged between 1.6 to 3.12 µg/mL against 37 different sensitive and resistant MTB strains (21 susceptible clinical isolates, 2 INH-resistant clinical isolates and 9 MDR clinical isolates).

Graphical abstract

*To whom correspondence should be addressed. Tel: (+1)502-852-5998. Fax: (+1)502-852-3899. gb.hammond@louisville.edu.

[‡]R. R. and G. B. H. share senior authorship.

Tuberculosis (TB) is the cause of 2 million deaths per year worldwide, with 98% occurring in developing countries. One of the highest incidence rates in the Americas occurs in Peru (228 cases per 100,000 population in 1999).^{1,2} The HIV epidemic has worsened this situation, especially in resource-poor countries. In the developing world, about 50% of HIV seropositive individuals may be co-infected with TB.³ Inadequate, incomplete or improperly supervised treatment regimens and co-infection with HIV have caused the emergence of multidrug-resistant (MDR) strains of *Mycobacterium tuberculosis* (MTB).⁴ Due to the urgent need to discover new drugs for the treatment of tuberculosis, several colorimetric methods for rapid screening of pure compounds or crude natural product extracts have been developed.⁵⁻⁷ The microplate Alamar blue assay (MABA) is a rapid, non-radioactive and inexpensive method that is based in the color change of the redox dye Alamar blue for the detection of viable *M. tuberculosis*.^{8,9} Members of our group have developed a variant of this assay that uses tetrazolium bromide instead of the more expensive Alamar blue dye.¹⁰ This tetrazolium microplate assay (TEMA) determines MIC values as quickly (6 to 7 days) and accurately as the MABA procedure.

Northern Peru, particularly the eastern slopes of the Andes Mountains and adjacent upper Amazon basin, the area from which *Claviija procera* B. Stahl (Theophrastaceae) originates, is exceedingly rich in diverse woody plants. This area has evolved into the richest source of biodiversity, exceeding the plant diversity of sub-Saharan Africa and Southeast Asia.¹¹ Only a small percentage of these Peruvian species has been investigated chemically and/or biologically. Using the TEMA assay, a group of 591 ethanol extracts from Peruvian plants was screened for antimycobacterial activity in vitro against the sensitive MTB H37Rv strain.¹² Thirty seven (6.3%) extracts showed MIC values $\leq 100 \mu\text{g/mL}$. The most active corresponded to the extract from the stems and bark of *C. procera* with a MIC of $12.5 \mu\text{g/mL}$. The genus and family of this species are poorly known. Indeed, no previous phytochemical or pharmacological investigations have been reported for *C. procera*. This plant is traditionally used by the Aguaruna people for the wound healing of cutaneous leishmaniasis. The bioassay-guided isolation of the ethanol extract of *C. procera* yielded an active fraction (MIC = $3.12 \mu\text{g/mL}$) for which final purification resulted in the isolation of aegicerin (**1**) and protoprimumagenin A (**2**). Compounds **1** and **2**, were previously isolated from *Aegiceras* spp.^{13,14} and from *Embelia schimperi*.¹⁵

Aegicerin (**1**) was investigated for its in vitro cytotoxic activity against Vero cells ($GI_{50} > 40 \mu\text{g/mL}$) using previously reported methodologies.¹⁶⁻¹⁸ (The GI_{50} is the concentration at which aegicerin inhibits the growth of cells by 50%). Evaluation of the antimycobacterial activity in vitro was performed using the TEMA method.¹⁰ The skeleton of **1** and **2** is not very common; it consists of an oleanane-type pentacyclic triterpene skeleton with a methyleneoxy bridge between C-13 and C-17. Aegicerin (**1**) was tested against a total of 37 different sensitive and resistant MTB strains (21 susceptible clinical isolates, 2 INH-resistant clinical isolates, and 9 MDR clinical isolates). Its MIC ranged between 1.6 to 3.12 $\mu\text{g/mL}$ (Table 1). Conversely, protoprimulagenin A (**2**) was inactive (MIC = 200 $\mu\text{g/mL}$).

Comparison of these structures **1** and **2** and their MIC values demonstrates that the presence of a carbonyl group at C-16 is crucial for antimycobacterial activity. Although a number of triterpenes possessing mild to strong antimycobacterial activity are known,¹⁹⁻²⁹ this work marks the first time that an oleanane-type triterpene has shown such a consistently high level of activity against a large panel of both: sensitive and resistant strains. While aegicerin (**1**) has been previously isolated, its wide-ranging anti-TB potential has not been realized.³⁰ In summary, using an inexpensive and reproducible colorimetric assay, a potent antimycobacterial triterpenoid, aegicerin (**1**), was isolated from a plant species belonging to a rare genus and family. Aegicerin (**1**) was found to possess excellent activity against a large number of MDR-MTB strains.

Experimental Section

General Experimental Procedures

The structures of the isolated compounds were identified by nuclear magnetic resonance (NMR; Varian Inova 500), ^1H NMR (500 MHz), ^{13}C NMR (125 MHz) and 2D-NMR analysis in CDCl_3 ; mass spectrometry (HREIMS at the Nebraska Center for Mass Spectrometry); and by comparison of spectroscopic data with those reported in the literature.¹¹ Column chromatography was carried out over silica gel (70-230 mesh, Merck). Fractions obtained from column chromatography were monitored by TLC (silica gel 60 F_{254} , Merck). Preparative TLC was carried out on Merck silica gel F_{254} plates (20 $\times$ 20 cm, 0.5 mm thick).

Plant Material

Clavija procera (Theophrastaceae) was collected near the community of Sukutim, province of Bagua, Department of Amazonas in August 1998 (vouchers #19191 kept at the Museo de Historia Natural, Universidad Nacional Mayor de San Marcos, Lima, Peru, and Missouri Botanical Garden, St. Louis, MO). The plant was identified by one of us (W.H.L.)

Extraction and Isolation

Dried stems and bark of *C. procera* (110 g) were extracted with 95% ethanol. This ethanol extract (6.25 g) was partitioned with the aid of solvents of different polarities. The activity was concentrated in the solvent system corresponding to 90% methanol-water. After concentration, this purified extract was chromatographed on a silica gel column using a hexane/ethyl acetate gradient. Five fractions (A – E) were obtained and evaluated for anti-MTB activity in vitro. The most active fraction (C, MIC = 3.12 µg/mL) was purified by silica gel preparative TLC using hexane-chloroform-methanol (25:10:2; triple development) resulting in the isolation of aegicerin (**1**, 5.7 mg, 0.005% yield) and protoprimulagenin A (**2**, 2.1 mg, 0.002% yield).

Bioassays

For the preparation of the inoculum, a suspension of MTB was made by mixing growth from slants (20 – 30 days old) with 100 µL of Tween 80 into 0.2% bovine serum albumin (Sigma Chemical Co., St. Louis, MO). Turbidity of the suspension was then adjusted to a McFarland standard No. 1 (3×10^7 CFU/mL) by adding Tween 80 and bovine serum albumin. The bacterial suspension (300 µL) was further transferred to 7.2 mL of 7H9GC broth (4.7 g of Middlebrook 7H9 broth base (Difco, Detroit, MI), 20 mL of 10% glycerol, 1 g of Bacto Casitone (Difco), 880 mL of distilled water, 100 mL OADC (oleic acid, albumin, dextrose, catalase) (Remel, Lenexa, KS)). For the bioassay, the extracts or pure compounds were resuspended in DMSO at a concentration of 4 mg/mL (stock solution). These stock solutions were further diluted with appropriate volumes of 7H9GC broth to yield final concentrations of 0.4 to 25 µg/mL. Final drug concentration ranges of standard antibiotics used as positive controls were 0.125 to 32 µg/mL for isoniazid and 0.063 to 16 µg/mL for rifampin. The plant extract or drug (100 µL) was mixed in the wells with 100 µL of bacterial inoculum, resulting in a final bacterial concentration of approx. 1.2×10^6 CFU/mL. The wells in column 11 served as inoculum-only controls. Solvent (DMSO) was included in every experiment as a negative control. The plates were sealed in plastic bags and then incubated at 37 °C for 5 days. On day 5, 50 µL of the tetrazolium-Tween 80 mixture {1.5 mL of tetrazolium [3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide] (Aldrich Chemical Co., Milwaukee, WI) at a dilution of 1 mg/mL in absolute ethanol and 1.5 mL of 10% Tween 80} was added to the wells and the plate was incubated at 37 °C for 24 h. After this incubation period, the growth of the microorganism was visualized by the change in color of the dye from yellow to purple. The tests were carried out in triplicate. MIC is defined as the lowest drug or plant extract concentration that prevents the aforementioned change in color.

Acknowledgments

The authors are grateful to the Iberoamerican Program for Science and Technology (CYTED), Project X. 11:PIBATUB (grant to R. R.), to the NIH and NSF (ICBG grant U01TW00331 to W.H.L., G.L. and A.J.V.), and the University of Louisville, Office of the Vice-Provost for Research (G.B.H.) for their financial support. We acknowledge the spectroscopic support provided by Dr. Neal Stolowich, Department of Chemistry, University of Louisville. Our gratitude is extended to the Aguaruna people for generously sharing their traditional medicinal information. In particular, we acknowledge the assistance of organizational/clan leaders, Apus, and members of OCCAAM, FAD, FECONARIN, and OAAM.

References and Notes

1. Dye C, Scheele S, Dolin P, Pathania V, Raviglion MC. *J Am Med Assoc.* 1999; 282:677–686.
2. Suárez PG, Watt CJ, Alarcón E, Portocarrero J, Zavala D, Canales R, Luelmo F, Espinal MA, Dye C. *J Infect Dis.* 2001; 184:473–478. [PubMed: 11471105]
3. Gazzard B. *Clin Med.* 2001; 1:62–68.
4. Zumla A, Grange J. *Br Med J.* 1998; 316:1962–1964. [PubMed: 9641941]
5. Yajko DM, Madej JJ, Lancaster MV, Sanders CA, Cawthon VL, Gee B, Babst A, Hadley WK. *J Clin Microbiol.* 1995; 33:2324–2327. [PubMed: 7494021]
6. Gomez-Florez R, Gupta S, Tamez-Guerra R, Mehta RT. *J Clin Microbiol.* 1995; 33:1842–1846. [PubMed: 7665657]
7. Palomino JC, Martin A, Camacho M, Guerra H, Swings J, Portaels F. *Antimicrob Agents Chemother.* 2002; 46:2720–2722. [PubMed: 12121966]
8. Collins LA, Franzblau C. *Antimicrob Agents Chemother.* 1997; 41:1004–1009. [PubMed: 9145860]
9. Franzblau SG, Witzig RS, McLaughlin JC, Torres P, Madico G, Hernandez A, Degnan MT, Cook MB, Quenzer VK, Ferguson RM, Gilman RH. *J Clin Microbiol.* 1988; 36:362–366.
10. Caviedes L, Delgado J, Gilman RH. *J Clin Microbiol.* 2002; 40:1873–1874. [PubMed: 11980982]
11. Gentry AH. *Proc Natl Acad Sci USA.* 1988; 85:156–159. [PubMed: 16578826]
12. Rojas, R., Caviedes, L., Lock, O., Gilman, RH. *Recent Progress in Medicinal Plants.* Govil, JN., editor. Vol. 13. Studium Press LLC; Houston: 2005. p. 429–441.
13. Zhang D, Wu J, Zhang S, Huang J. *Fitoterapia.* 2005; 76:131–133. [PubMed: 15664478]
14. Rao KV. *Tetrahedron.* 1964; 20:973–977.
15. Machocho AK, Kiprono PC, Grinberg S, Bittner S. *Phytochemistry* 2003. 62:573–577.
16. Boyd MR, Paull KD. *Drug Dev Res.* 1995; 34:91–109.
17. Grever MR, Schepartz SA, Chabner BA. *Sem Oncol.* 1992; 19:622–638.
18. Skehan P, Storeng R, Scudiero D, Monks A, McMahon J, Vistica D, Warren JT, Bokesch H, Kenney S, Boyd MR. *J Nat Cancer Inst.* 1990; 82:1107–1112. [PubMed: 2359136]
19. Akihisa T, Franzblau SG, Ukiya M, Okuda H, Zhang F, Yasukawa T, Suzuki T, Kimura Y. *Biol Pharm Bull.* 2005; 28:158–160. [PubMed: 15635183]
20. Caldwell CG, Franzblau SG, Suarez E, Timmermann BN. *J Nat Prod.* 2000; 63:1611–1614. [PubMed: 11141098]
21. Cantrell CL, Rajab MS, Franzblau SG, Fischer NH. *J Nat Prod.* 1999; 62:546–548. [PubMed: 10217705]
22. Newton SM, Lau C, Wright CW. *Phytother Res.* 2000; 14:303–322. [PubMed: 10925394]
23. Okunade AL, Elvin-Lewis MPF, Lewis WH. *Phytochemistry.* 2004; 65:1017–1032. [PubMed: 15110681]
24. Wächter GA, Valcic S, Flagg ML, Franzblau SG, Montenegro G, Suarez E, Timmermann BN. *Phytomedicine.* 1999; 6:341–345. [PubMed: 11962541]
25. Woldemichael GM, Gutierrez-Lugo MT, Franzblau SG, Wang Y, Suarez E, Timmermann BN. *J Nat Prod.* 2004; 67:598–603. [PubMed: 15104489]
26. Lall N, Marion Meyer JJ, Wang Y, Bapela NB, van Rensburg CEJ, Fourie B, Franzblau SG. *Pharm Biol.* 2005; 43:353–357.
27. Copp BR. *Nat Prod Rep.* 2003; 20:535–557. [PubMed: 14700198]

28. Wächter GA, Franzblau SG, Montenegro G, Hoffmann JJ, Maiese WM, Timmermann BN. *J Nat Prod.* 2001; 64:1463–1464. [PubMed: 11720535]
29. Cantrell CL, Rajab MS, Franzblau SG, Fronczek FR, Fischer NH. *Planta Med.* 1999; 65:732–734. [PubMed: 10630115]
30. This result resonates with Cordell and Colvard's observation that of all ethnomedical reports on about 14,300 species of plants in NAPRALERT, there has been no compound isolated and no biological work conducted for 8387 (58.6%) of them. See: Cordell GA, Colvard MD. *J Ethnopharmacol.* 2005; 100:5–14. [PubMed: 16009517]

Table 1Antimycobacterial Activity of INH, RIF and Aegicerin (**1**)

strains (number of strains tested)	MIC (µg/mL)		
	INH ^a	RIF ^b	aegicerin (1)
<i>Mycobacterium tuberculosis</i>			
H37Rv	0.125	0.063	3.1
susceptible clinical isolates (21)	0.125	0.063	1.6-3.1
INH-resistant clinical isolates (2)	2-4	0.063-0.125	1.6-3.1
MDR clinical isolates (13)	4->32	2->16	1.6-3.1

^aINH = isoniazid.^bRIF = rifampin