CORRESPONDENCE

Screening for Bowel Cancer: Increasing Participation via Personal Invitation

A Randomized Intervention Study

by Dr. sc. hum. Michael Hoffmeister, Dr. sc. hum. Bernd Holleczeck, Dr. sc. hum. Nadine Zwink, Dr. sc. hum. Christian Stock, Dipl.-Inform. Med. Christa Stegmaier, and Prof. Dr. med. Hermann Brenner in issue 6/2017

Missing Information

Hoffmeister et al. studied the effect of a written invitation to participate in bowel cancer screening on participation rates in a randomized controlled trial (RCT) (1). Since the tested materials were not published alongside the article, interpreting the results is impossible. It is a widespread problem that especially the details of non-medication-based interventions are not reported to a satisfactory degree (2). Materials used should always be published—for example, as a web appendix.

The name of the SAMS Study („Saarland gegen Darmkrebs – Machen Sie mit!“- *Saarland Against Bowel Cancer—Join In!*) implies that the intervention is such that people are invited to participate. This is in direct contradiction to current health political objectives. In the context of the German National Cancer Plan, several objectives were formulated between 2008 and 2011 on the basis of a wide consensus, in order to strengthen patients' competence; these include the provision of evidence-based information and participatory decision making, among others (3). In the meantime, the bowel cancer screening foundation *Stiftung LebensBlicke* has articulated in its declaration of Mannheim: „All entitled persons have the right to balanced information on the benefits and risks of bowel cancer screening, which allows for informed decision-making.“

These requirements are currently being implemented in the context of the Law on the Further Development of the Early Detection of Cancer and Quality Assurance Through Clinical Cancer Registries, which provides for „comprehensive and comprehensible information for insurance scheme members as regards the benefits and risks of the relevant procedure“ (§ 25a paragraph 1 No. 2 SGB V [social security statutes]). Relevant information for organized bowel cancer screening has been developed with the help of citizens and experts (4).

If the materials used in the SAMS Study did not adhere to these standards, an opportunity was missed to ensure that the requirement for evidence-based information and participatory decision making is gaining default status.

DOI: 10.3238/arztebl.2017.0426a

REFERENCES

1. Hoffmeister M, Holleczeck B, Zwink N, Stock C, Stegmaier C, Brenner H: Screening for bowel cancer: increasing participation via personal invitation—a randomized intervention study. *Dtsch Arztebl Int* 2017; 114: 87–93.
2. Glasziou P, Altman DG, Bossuyt P, et al.: Reducing waste from incomplete or unusable reports of biomedical research. *Lancet* 2014; 383: 267–76.
3. Bundesministerium für Gesundheit: Ziele des Nationalen Krebsplans. www.bundesgesundheitsministerium.de/themen/praevention/nationaler-krebsplan/oeffentlichkeitsarbeit/handlungsfelder/ziele-des-nationalen-krebsplans.html (last accessed on 17 February 2017).
4. Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen: Einladungsschreiben und Entscheidungshilfen zum Darmkrebs-Screening: Abschlussbericht. www.iqwig.de/de/projekte-ergebnisse/projekte/gesundheitsinformation/p15-01-einladungsschreiben-und-entscheidungshilfe-zum-darmkrebs-screening.6620.html#overview (last accessed on 17 February 2017).

Roland Buechter, Dr. rer. medic. Klaus Koch, Regina Will

Ressort Gesundheitsinformation, Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen (IQWiG), Köln
roland.buechter@iqwig.de

Preventing Bowel Cancer Through Hormones

„Targeted invitations can markedly increase participation rates in cancer screening“—this was the conclusion of the authors of the article (1). The results continue to be sobering for women.

As a result of the written invitation, participation rates in screening colonoscopies rose from 4.5% to 6.2%. The authors list selection factors. The editorial points out that only 1.9% of those who had been sent a written invitation attended colonoscopy screening within the year's screening period. This is disappointing for women as bowel cancer is the third most common cancer-related cause of death in women. This means that postmenopausal women need a hormonal alternative.

The estrogen receptor beta (ER-beta) is known to have anti-proliferative and anti-inflammatory characteristics. The effects of these can be substantially increased by using ER-beta agonists, an approach that is already being used in bowel cancer therapy (2). In order for this approach to be successful, the need is for enough ER-beta, whose density decreases notably after the menopause as a result of prolonged estrogen deficiency. Hormone replacement immediately after the menopause can prevent such a receptor loss. The bowel mucosa benefits, thanks to the anti-proliferative and anti-inflammatory characteristics of the ER-beta, or, indirectly, from estrogen substitution, which protects against receptor loss.

The evidence is sound: the 2016 international guideline on hormone replacement points to hormonal bowel cancer prevention. A study reported by Long et al. (3) is cited, which reports a halved risk (relative risk 0.47) after 9–14 years of hormone

replacement. This preventive potential can be used as one of many co-indications; the risk for healthy women is very low. The negative reports of the Women's Health Initiative (WHI) on hormone replacement therapy since 2002 are now history. After 14 years, the 2016 WHI study initiators apologized for an incorrect evaluation and unnecessary scaremongering (4).

In conclusion: the effort involved in increasing screening participation rates should be balanced by providing hormonal bowel cancer prevention.

DOI: 10.3238/arztebl.2017.0426b

REFERENCES

1. Hoffmeister M, Hollecsek B, Zwink N, Stock C, Stegmaier C, Brenner H: Screening for bowel cancer: increasing participation via personal invitation—a randomized intervention study. *Dtsch Arztebl Int* 2017; 114: 87–93.
2. Nguyen-Vu T, Wang J, Mesmar F, et al.: Estrogen receptor beta reduces colon cancer metastasis through a novel miR-205-PROX1 mechanism. *Oncotarget* 2016; 7: 42159–71.
3. Long MD, Martin CF, Galanko JA, Sandler RS: Hormone replacement therapy, oral contraceptive use, and distal large bowel cancer: a population-based case-control Study. *Am J Gastroenterol* 2010; 105: 1843–50.
4. Manson JE, Kaunitz AM: Menopause management-getting clinical care back on track. *N Engl J Med* 2016; 374: 803–6.

Prof. Dr. med. Dipl. Psych. J. Matthias Wenderlein
Universität Ulm
wenderlein@gmx.de

In Reply:

Büchter and colleagues assume on the basis of the title of our study (“Saarland gegen Darmkrebs – Machen Sie mit!”) that the interventions that we tested contradict current health policy aims. We disagree in this point. The supreme objective of health policy in colorectal cancer screening is to lower colorectal cancer incidence and mortality. Accordingly, in addition to the improvement of information offers, the objectives of the German National Cancer Plan (2) include primarily an improved uptake of the offered screening examinations. We focused on both of these objectives in our model project, which was planned in 2011 and conducted from 2012–2015. The information materials were developed at the time the study was being planned, on the basis of the information brochure of the Federal Joint Committee (*Gemeinsamer Bundesausschuss*, G-BA) and with support from the experts of the Cancer Information Service (*Krebsinformationsdienst*, KID) at the German Cancer Research Center (*Deutsches Krebsforschungszentrum*).

We agree with the authors of the letter to the editor that everyone should receive balanced information about the benefits and risks and about the offers of colorectal cancer screening according to the claim that is anchored in the law on early detection and registration of cancer (KFRG). To the best of our knowledge and according to the experts involved, this was implemented in an adequate manner. Speaking of a missed opportunity in our study because we did not consider

invitation letters and brochures that were developed in the meantime with the involvement of the correspondents from the German Institute for Quality and Efficiency in Health Care (*Institut für Qualität und Wirtschaftlichkeit im Gesundheitswesen*, IQWiG) and that were published in December 2016 seems inappropriate. Furthermore, our study materials were made available to the authors from IQWiG in the middle of 2016, and for this reason a balanced interpretation should have been possible for them.

With the SAMS study, we have, for the first time, presented comprehensive empirical data on the efficacy of different invitation schemes in Germany. Regardless of potential advantages and disadvantages of IQWiG's information materials, which thus far have not been empirically tested in a comparable way, a reduction in the incidence and mortality of colorectal cancer should be the primary concern of all parties involved. A large proportion of the more than 60 000 new cases and the more than 25 000 deaths due to colorectal cancer in Germany per year could be prevented by improving the utilization of screening offers. The decline in colorectal cancer incidence and mortality that is becoming apparent (3) should be strengthened and accelerated as soon as possible by means of an organized screening program (4). Our study aimed to contribute robust empirical evidence to answer the question of how this may best be achieved. We are happy to make the study materials we used (invitation letters, brochures) available to the public and to *Deutsches Ärzteblatt* for publication as a web appendix.

We thank Mr Wenderlein for his comments that we read with interest, but which do not contribute new insights to the subject of our study. DOI: 10.3238/arztebl.2017.0427

REFERENCES

1. Hoffmeister M, Hollecsek B, Zwink N, Stock C, Stegmaier C, Brenner H: Screening for bowel cancer: increasing participation via personal invitation—a randomized intervention study. *Dtsch Arztebl Int* 2017; 114: 87–93.
2. Bundesministerium für Gesundheit. Ziele des Nationalen Krebsplans. www.bundesgesundheitsministerium.de/themen/praevention/nationaler-krebsplan/oeffentlichkeitsarbeit/handlungsfelder/ziele-des-nationalen-krebsplans.html (last accessed on 19 April 2017).
3. Brenner H, Schrotz-King P, Hollecsek B, Katalinic A, Hoffmeister M: Declining bowel cancer incidence and mortality in Germany. *Dtsch Arztebl Int* 2016; 113: 101–6.
4. Brenner H, Stock C, Hoffmeister M: Colorectal cancer screening: the time to act is now. *BMC Med* 2015; 13: 262.

PD Dr. sc. hum. Michael Hoffmeister
Abteilung Klinische Epidemiologie und Altersforschung
Deutsches Krebsforschungszentrum (DKFZ), Heidelberg
m.hoffmeister@dkfz.de

Dr. sc. hum. Bernd Hollecsek
Krebsregister Saarland, Saarbrücken

PD Dr. sc. hum. Christian Stock
Institut für Medizinische Biometrie und Informatik, Universität Heidelberg

Prof. Dr. med. Hermann Brenner
Abteilung Klinische Epidemiologie und Altersforschung
Deutsches Krebsforschungszentrum (DKFZ), Heidelberg

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.