Regional Differences in the Prevalence of Major Opportunistic Infections among Antiretroviral-Naïve Human Immunodeficiency Virus Patients in Japan, Northern Thailand, Northern Vietnam, and the Philippines

Louie Mar A. Gangcuangco,^{1,2†} Ikumi Sawada,^{1†} Naho Tsuchiya,^{1,3} Cuong D. Do,⁴ Thanh Thuy T. Pham,⁴ Archawin Rojanawiwat,⁵ Marissa Alejandria,⁶ Katerina Leyritana,⁶ Yoshiyuki Yokomaku,⁷ Panita Pathipvanich,⁸ and Koya Ariyoshi^{1*}

¹Institute of Tropical Medicine (NEKKEN), Graduate School of Biomedical Science, Nagasaki University, Nagasaki, Japan; ²Bridgeport Hospital-Yale New Haven Health, Bridgeport, Connecticut; ³Tohoku Medical Megabank Organization, Tohoku University, Miyagi, Japan; ⁴Bach Mai Hospital, Hanoi, Vietnam; ⁵Ministry of Public Health, Nonthaburi, Thailand; ⁶Philippine General Hospital, University of the Philippines, Manila, Philippines; ⁷National Hospital Organization Nagoya Medical Center, Nagoya, Japan; ⁸Lampang Hospital, Lampang, Thailand

Abstract. To identify regional differences in the distribution of opportunistic infections (OIs) among human immunodeficiency virus (HIV)-infected patients in Asia, the medical records of antiretroviral therapy (ART)-naïve patients who attended the following tertiary hospitals from 2003 to 2011 were reviewed: Nagoya Medical Center (NMC, Nagoya, Japan), Lampang Hospital (LPH, Lampang, northern Thailand), Bach Mai Hospital (BMH, Hanoi, northern Vietnam), and Philippine General Hospital (PGH, Manila, Philippines). Logistic regression analyses were performed to identify associations between country of origin and risk of major OIs. In total, 1,505 patients were included: NMC, $N = 365$; LPH, $N = 442$; BMH, $N = 384$; and PGH, $N = 314$. The median age was 32 years, and 73.3% of all patients were male. The median CD4 count was 200 cells/ μ L. Most patients at NMC and PGH were men who have sex with men. Injection drug users were most common at BMH (35.7%). *Mycobacterium tuberculosis* (TB) was most common at PGH ($N = 75$) but was rare at NMC ($N = 4$). *Pneumocystis pneumonia* (PCP) prevalence was highest at NMC ($N = 74$) and lowest at BMH ($N = 13$). Multivariable logistic regression showed increased odds of TB at PGH (adjusted odds ratio [aOR] = 42.2, 95% confidence interval [CI] = 14.6–122.1), BMH (aOR = 12.6, CI = 3.9–40.3), and LPH (aOR = 6.6, CI = 2.1–21.1) but decreased odds of PCP at BMH (aOR = 0.1, CI = 0.04–0.2) and LPH (aOR = 0.2, CI = 0.1–0.4) compared with those at NMC. The cryptococcosis risk was increased at LPH (aOR = 6.2, CI = 0.9–41.0) compared with that at NMC. Cytomegalovirus (CMV) retinitis prevalences were similar in all countries. OI prevalence remained high among ART-naïve patients in our cohort. The risks of TB, PCP, and cryptococcosis, but not CMV retinitis, differed between countries. Improved early HIV detection is warranted.

INTRODUCTION

The clinical characteristics of human immunodeficiency virus (HIV)-infected patients before initiating antiretroviral therapy (ART) vary between developed and developing countries and are influenced by sociodemographic factors, geographic distribution of pathogens,^{1,2} access to health care,³ host genetic factors,⁴ and virus subtypes.⁵

A range of HIV-associated risk groups has previously been identified in Asia; these groups include men who have sex with men (MSM) in Japan and the Philippines, injection drug users (IDUs) in Vietnam and southern China, and female sex workers and their clients in other Asian countries.⁶ Among newly diagnosed HIV-infected patients, the prevalence of acquired immune deficiency syndrome (AIDS) also varies widely from 10% in Taiwan⁷ to 30% in Japan⁸ and New Zealand⁹ and 53% in Bangkok, Thailand.¹⁰ A study in Cambodia reported that 100% of newly diagnosed HIV-infected patients had AIDS.¹¹

The major opportunistic infections (OIs) in Asia are extrapulmonary *Mycobacterium tuberculosis* (TB), cryptococcosis, and *Pneumocystis pneumonia* (PCP).^{6,12} The predominant AIDS-defining illnesses in Thailand are TB, PCP, and cryptococcosis,¹³ whereas those in Japan are PCP, cytomegalovirus (CMV) infection, and candidiasis

according to a recent report from the Japanese Ministry of Health, Labor and Welfare.¹⁴ In two different HIV care facilities in Ho Chi Minh City, Vietnam, oral candidiasis and TB were reported to be major OIs, but PCP was reported to be uncommon.^{15,16} Disseminated *Penicillium marneffei* infection, that is, penicilliosis, is another well-known unique opportunistic mycosis in the northern Indochina region.⁶ However, there have been no comprehensive reports regarding HIV-related illnesses from the Philippines in the ART era.

Despite the diversity of results produced in individually published studies, a systematic comparison of the prevalence of HIV-related OIs among Asian countries has not been performed. Accordingly, we aimed to describe the demographic and clinical characteristics of newly diagnosed HIV patients at four tertiary hospitals providing HIV care in Japan, northern Thailand, northern Vietnam, and the Philippines and to investigate the factors associated with major HIV-related OIs adjusting for country of origin, age, sex, HIV-related risk behaviors, and immune status at the time of HIV diagnosis.

MATERIALS AND METHODS

Study design and data collection. We included all ART-naïve patients ≥ 15 years old with an available initial CD4 + T-cell count (CD4 count) who were enrolled/registered at the following sites: Nagoya Medical Center (NMC; Nagoya, Japan), Lampang Hospital (LPH; Lampang, northern Thailand), Bach Mai Hospital (BMH; Hanoi, northern Vietnam), and Philippine General Hospital (PGH; Manila, Philippines).

* Address correspondence to Koya Ariyoshi, Institute of Tropical Medicine, Nagasaki University, 1-12-4 Sakamoto, P.O. Box 852-8523, Nagasaki, Japan. E-mail: kari@nagasaki-u.ac.jp

† These authors contributed equally to this work.

Age, sex, self-reported mode of HIV transmission, baseline CD4 count (cells/ μ L), clinical stage at the time of HIV diagnosis, and OIs diagnosed at the initial visit before initiation of ART were collected from patients' medical records. Patients who were initially registered at the hospitals during the following periods were included: NMC, October 2008 to November 2011; LPH, January 2003 to December 2010; BMH, October 2009 to September 2010; and PGH, January 2004 to December 2011.

Study sites. All four study sites are government-funded hospitals that provide HIV treatment within each country. NMC is a tertiary hospital located in Nagoya City, one of the largest cities in Japan. NMC is a referral hospital for the AIDS treatment core hospitals that are located in the central region of Japan, serving three other surrounding prefectures. LPH is the largest government referral hospital in northern Thailand and is located in Lampang Province, which has the fifth highest prevalence of HIV in Thailand.¹⁷ LPH has an HIV clinic that has provided HIV testing and medical treatment since 1995. BMH is the largest tertiary hospital in northern Vietnam. The Department of Infectious Diseases officially launched the HIV Outpatient Clinic and the HIV Admission Ward in November 2009, which provides free ART. BMH also functions as a voluntary counseling and testing site. PGH is the largest tertiary hospital in the Philippines and is located in the capital city of Manila. In 1995, PGH established the STD-AIDS Guidance Intervention Prevention Unit, where both walk-in patients and interhospital referrals can receive anonymous testing for HIV as well as ART.

Reporting of OIs. In this study, clinical stage was categorized based on the World Health Organization-revised clinical staging of HIV/AIDS.¹⁸ Each OI was diagnosed at each study site as outlined by their local or international guidelines, which included the following: the anti-HIV treatment guidelines by the Ministry of Health, Labor and Welfare, Japan,¹⁹ the Clinical Management of HIV/AIDS Patients at Day Care Centers in LPH,²⁰ the Guidelines for HIV/AIDS Diagnosis and Treatment by the Vietnam Ministry of Health,²¹ and the Clinical Practice Guidelines for the Diagnosis, Treatment, Prevention and Control of Tuberculosis in Adult Filipinos (Supplemental Table 1).²² All OIs diagnosed before the initiation of ART were collected. Disease recurrences were not included.

Statistical analyses. Chi-square tests were used to identify differences in the categorical variables between countries, and Kruskal-Wallis tests were performed to compare continuous variables. The proportion of HIV-related illnesses was calculated in patients with stage 3 or 4 HIV/AIDS. Of the HIV-related illnesses, TB, PCP, cryptococcosis, CMV retinitis, and herpes zoster were considered as major HIV-related infections in Asia based on previous publications⁶ and the fact that their diagnoses were less likely to be influenced by limited access to endoscopy, pathological examinations, or advanced microbiological techniques such as polymerase chain reaction or blood culture of *Mycobacterium*. Their prevalences were determined with a 95% confidence interval (CI) calculated using the Wilson score method. Logistic regression analyses were conducted to evaluate the factors associated with major HIV-related infections in all HIV-infected patients. Multivariable logistic regression models were adjusted for age at registration, sex, self-reported mode of HIV

transmission, and country of origin. Age was divided by a factor of 10 to indicate the odds ratio per increase in decade. All analyses were conducted with STATA IC version 12.0 (Stata Corp., College Station, TX).

Ethical considerations. Institutional approval was obtained from the ethics committees of NMC and the Institute of Tropical Medicine, Nagasaki University. De-identified data from previously approved cohort studies were obtained from the collaborators at PGH, LPH, and BMH. All procedures adhered to country-specific laws on HIV and AIDS, including the Japanese regulations regarding HIV disclosure and the Philippine AIDS Prevention and Control Act. No patients were directly contacted or re-examined for this study.

RESULTS

Demographic and clinical characteristics. Data from 1,510 newly registered ART-naïve HIV-infected patients were available from the four study sites (NMC, $N = 367$; LPH, $N = 442$; BMH, $N = 387$; and PGH, $N = 314$). After excluding patients younger than 15 years (two at NMC and three at BMH), 1,505 patients were included in the analysis.

The demographic and clinical characteristics of the patients are summarized in Table 1. Overall, the median (interquartile range) age at registration was 32 years (27–39 years), and this age varied significantly by study site, with younger patients visiting PGH ($P < 0.001$). Approximately, 90% of patients at NMC were Japanese, and half of the rest were Asian foreigners. Most patients at LPH, BMH, and PGH were Thai, Kinh, and Filipino, respectively. The majority of the patients at NMC and PGH were male, and the proportions of male patients at NMC and PGH differed significantly from those of the other sites ($P < 0.001$). The different sex distributions were likely due to differences in risk group: PGH and NMC patients were mostly MSM, whereas LPH patients were mostly heterosexual. Although the majority of BMH patients were heterosexual, the second most common risk group was IDUs, the vast majority of whom were male (98.3%). Overall, 753 patients (52.7%) had already progressed to advanced immune deficiency; 49.8% ($N = 750$) had a CD4 count < 200 cells/ μ L and 29.4% ($N = 442$) had AIDS-defining illnesses (stage 4).

Pattern of OIs and HIV-related illnesses. Considerable variation was observed in the distribution of HIV-related illnesses. Figure 1 shows the proportions of major HIV-related infections among the patients with clinical stage 3 or 4 HIV/AIDS as indicated in Table 1. TB, including both pulmonary TB (PTB) and extrapulmonary TB, was the most common OI at PGH and BMH. In contrast, TB was rather rare at NMC. The prevalence of PCP was highest at NMC and lowest at BMH. Cryptococcosis was most frequently diagnosed at LPH. CMV retinitis and herpes zoster were diagnosed at all sites, although CMV antigenemia ($N = 35$, 27.3%) and CMV enteritis ($N = 2$, 1.6%) were diagnosed only at NMC. The prevalence of herpes zoster was only slightly higher at LPH than at the other hospitals. HIV encephalopathy and malignant lymphoma were diagnosed only at NMC. Interestingly, Kaposi's sarcoma was observed only at NMC and PGH, and its prevalence showed a significant association with MSM: seven patients with Kaposi's sarcoma among 552 MSM (1.3%) versus two patients among

TABLE 1
Characteristics of all patients from four hospitals

	NMC (N = 365)	LPH (N = 442)	BMH (N = 384)	PGH (N = 314)	P value
Age at registration	37 [31, 45]	34 [29, 40]	31 [28, 35]	28 [25, 33]	< 0.001
Male	343 (94.0)	213 (48.2)	246 (64.1)	301 (95.9)	< 0.001
Risk behavior					
Heterosexual	68 (18.6)	395 (89.4)	228 (59.4)	36 (12.1)	
MSM/bisexual	263 (72.1)	12 (2.7)	3 (0.8)	258 (86.9)	
IDU	19 (5.2)	12 (2.7)	137 (35.7)	3 (1.0)	
Others	15 (4.1)	23 (5.3)	16 (4.2)	0 (–)	< 0.001
CD4 + T cell count (cells/ μ L)	234 [58, 386]	194 [29, 391]	111 [40, 337]	266 [74, 400]	< 0.001
CD4 + T cell count category (cells/ μ L)					
≥ 200	203 (55.6)	220 (49.8)	142 (37.0)	190 (60.5)	
51–200	75 (20.6)	79 (17.9)	122 (31.8)	57 (18.2)	
≤ 50	87 (23.8)	143 (32.4)	120 (31.3)	67 (21.3)	< 0.001
Clinical stage at the first visit					
Stage 1	226 (61.9)	221 (50.0)	141 (36.7)	128 (40.8)	
Stage 2	11 (3.0)	31 (7.0)	31 (8.1)	14 (4.5)	
Stage 3/4	128 (35.1)	186 (42.1)	211 (55.0)	104 (33.1)	
No information	0 (–)	4 (0.9)	1 (0.3)	68 (21.7)	< 0.001

BMH = Bach Mai Hospital; IDU = injecting drug user; LPH = Lampang Hospital; MSM = men having sex with men; NMC = Nagoya Medical Center; PGH = Philippine General Hospital. Values are given as actual count (%) for categorical variables or median [interquartile range] for continuous variables. If no information, “–” is displayed in the number column. CD4 is not taken into consideration in the staging.

936 non-MSM (0.2%) ($P = 0.011$). Esophageal candidiasis was commonly diagnosed at NMC and moderately diagnosed at BMH. As expected from previous reports,²³ penicilliosis was observed exclusively at BMH and LPH (Supplemental Table 2).

Factors associated with TB, PCP, cryptococcosis, CMV retinitis, and herpes zoster. Logistic regression analyses were performed to identify the factors associated with the five major HIV-related infections. All patients from the four hospitals were included in the analysis. Injection drug use, low CD4 count, and hospital of registration were significantly associated with TB in both the univariable and multivariable analyses (Table 2). Hospital of registration substantially affected the risk of TB diagnosis, with the highest adjusted odds ratio (aOR) for PGH (aOR = 42.2, 95% CI = 14.6–122.1), followed by BMH (aOR = 12.6, 95% CI = 3.9–40.3) and LPH (aOR = 6.6, 95% CI 2.1–21.1). Injection drug use remained independently associated with

TB in the final multivariable logistic regression model. Factors associated with PCP in the univariable analysis were age, male gender, injection drug use, and CD4 count (Table 2). In the final multivariable model, only CD4 count and hospital of registration remained significantly associated with PCP. Notably, patients from LPH (aOR = 0.2, 95% CI = 0.1–0.4) and BMH (aOR = 0.1, 95% CI = 0.04–0.2) had significantly lower odds of PCP diagnosis than patients from NMC. The factors associated with cryptococcosis in the univariable analysis were low CD4 count and registration at LPH (Table 2). In the final multivariable model, a trend toward an increased risk of cryptococcosis was observed among patients at LPH (aOR = 6.2, 95% CI = 0.9–41.0). In contrast to the other OIs investigated, there were no significant regional differences in the diagnosis of CMV retinitis, even after adjusting for CD4 count (Table 2). Similarly, there were no regional differences in the risk of herpes zoster (Table 2).

FIGURE 1. Prevalence of major human immunodeficiency virus (HIV)-related infections among patients infected with clinical stage 3 or 4 HIV at each hospital. NMC = Nagoya Medical Center (N = 128), LPH = Lampang Hospital (N = 186), BMH = Bach Mai Hospital (N = 211), and PGH = Philippine General Hospital (N = 104). TB = tuberculosis, PCP = *Pneumocystis pneumonia*, CMV = cytomegalovirus. TB includes pulmonary TB and extrapulmonary TB. The presence of both conditions was counted as one case. Error bars represent 95% confidence intervals calculated using the Wilson score method. P values were calculated using χ^2 tests.

TABLE 2
Associated factors for the major HIV-related infections among all patients

Risk factors	Crude OR (95% CI, <i>P</i> value)	Adjusted OR (95% CI, <i>P</i> value)
Tuberculosis		
Age*	1.0 (0.8–1.2, 0.891)	1.3 (1.1–1.5, 0.034)
Male	1.9 (1.3–2.8, 0.002)	1.3 (0.8–2.2, 0.320)
Risk behavior		
Heterosexual	1.0	1.0
MSM/bisexual	1.2 (0.9–1.8, 0.285)	0.8 (0.4–1.6, 0.441)
IDU	3.0 (1.9–4.6, < 0.001)	2.0 (1.1–3.6, 0.013)
CD4+ T cell count category (cells/ μ L)		
≥ 200	1.0	1.0
51–200	3.0 (2.0–4.5, < 0.001)	2.5 (1.9–4.0, < 0.001)
≤ 50	4.0 (2.7–5.9, < 0.001)	4.0 (2.6–6.1, < 0.001)
Hospital		
Nagoya Medical Center	1.0	1.0
Lampang Hospital	8.7 (3.1–24.7, < 0.001)	6.6 (2.1–21.1, 0.001)
Bach Mai Hospital	19.4 (7.1–53.8, < 0.001)	12.6 (3.9–40.3, < 0.001)
Philippine General Hospital	28.3 (10.2–78.5, < 0.001)	42.2 (14.6–122.1, < 0.001)
<i>Pneumocystis jirovecii</i> pneumonia		
Age*	1.5 (1.3–1.7, < 0.001)	1.2 (0.9–1.4, 0.141)
Male	2.1 (1.3–3.2, 0.001)	1.0 (0.6–1.8, 0.921)
Risk behavior		
Heterosexual	1.0	1.0
MSM/bisexual	1.6 (1.1–2.2, 0.007)	0.8 (0.4–1.4, 0.388)
IDU	0.4 (0.2–0.8, 0.016)	0.5 (0.2–1.3, 0.154)
CD4+ T cell count category (cells/ μ L)		
≥ 200	1.0	1.0
51–200	8.7 (4.5–16.8, < 0.001)	10.4 (5.3–20.5, < 0.001)
≤ 50	23.9 (13.0–43.9, < 0.001)	35.3 (18.7–66.6, < 0.001)
Hospital		
Nagoya Medical Center	1.0	1.0
Lampang Hospital	0.4 (0.8–0.6, < 0.001)	0.2 (0.1–0.4, < 0.001)
Bach Mai Hospital	0.1 (0.1–0.3, < 0.001)	0.1 (0.04–0.2, < 0.001)
Philippine General Hospital	0.6 (0.4–0.9, 0.013)	0.7 (0.4–1.2, 0.226)
Cryptococcosis		
Age*	1.0 (0.7–1.4, 0.996)	0.8 (0.5–1.3, 0.330)
Male	1.0 (0.5–2.2, 0.974)	1.0 (0.4–2.6, 0.984)
Risk behavior		
Heterosexual	1.0	1.0
MSM/bisexual	0.4 (0.1–1.0, 0.051)	3.4 (0.5–24.4, 0.221)
IDU	0.9 (0.3–2.8, 0.917)	2.6 (0.6–10.9, 0.202)
CD4+ T cell count category (cells/ μ L)		
≥ 200	1.0	1.0
51–200	4.6 (0.8–25.1, 0.080)	4.7 (0.8–26.5, 0.081)
≤ 50	27.1 (6.4–114.4, < 0.001)	25.9 (6.0–112.6, < 0.001)
Hospital		
Nagoya Medical Center	1.0	1.0
Lampang Hospital	3.4 (1.4–8.5, 0.008)	6.2 (0.9–41.0, 0.058)
Bach Mai Hospital	0.6 (0.2–2.3, 0.447)	0.8 (0.9–7.6, 0.877)
Philippine General Hospital	–	–
CMV retinitis		
Age*	1.1 (0.7–1.7, 0.553)	1.0 (0.6–1.6, 0.918)
Male	0.8 (0.3–2.2, 0.738)	0.4 (0.1–1.5, 0.157)
Risk behavior		
Heterosexual	1.0	1.0
MSM/bisexual	1.2 (0.5–3.2, 0.698)	1.2 (0.3–5.2, 0.822)
IDU	0.5 (0.1–3.7, 0.474)	0.6 (0.1–6.2, 0.691)
CD4+ T cell count category (cells/ μ L)		
≥ 200	1.0	1.0
51–200	11.5 (1.3–98.8, 0.026)	13.5 (1.5–117.3, 0.018)
≤ 50	26.2 (3.4–199.9, 0.002)	32.8 (4.2–253.0, 0.001)
Hospital		
Nagoya Medical Center	1.0	1.0
Lampang Hospital	0.5 (0.1–1.6, 0.227)	0.3 (0.1–1.5, 0.145)
Bach Mai Hospital	0.5 (0.2–1.9, 0.326)	0.3 (0.1–1.8, 0.203)
Philippine General Hospital	0.8 (0.3–2.6, 0.749)	0.9 (0.3–3.0, 0.838)
Herpes zoster		
Age*	1.1 (0.8–1.4, 0.712)	1.0 (0.7–1.3, 0.999)
Male	1.1 (0.6–2.0, 0.702)	1.4 (0.7–3.0, 0.341)

(continued)

TABLE 2
Continued

Risk factors	Crude OR (95% CI, <i>P</i> value)	Adjusted OR (95% CI, <i>P</i> value)
Risk behavior		
Heterosexual	1.0	1.0
MSM/bisexual	0.9 (0.5–1.5, 0.635)	1.3 (0.4–3.7, 0.664)
IDU	0.7 (0.3–1.8, 0.424)	0.9 (0.3–2.9, 0.901)
CD4+ T cell count category (cells/ μ L)		
≥ 200	1.0	1.0
51–200	1.6 (0.9–3.0, 0.111)	1.6 (0.8–3.0, 0.152)
≤ 50	1.0 (0.5–1.9, 0.985)	0.9 (0.5–1.8, 0.729)
Hospital		
Nagoya Medical Center	1.0	1.0
Lampang Hospital	1.8 (0.9–3.6, 0.078)	2.6 (0.9–7.8, 0.084)
Bach Mai Hospital	0.7 (0.3–1.7, 0.450)	0.9 (0.3–3.2, 0.873)
Philippine General Hospital	0.9 (0.4–2.1, 0.787)	0.9 (0.4–2.4, 0.927)

CI = confidence interval; HIV = human immunodeficiency virus; IDU = injection drug user; MSM = men having sex with men; OR = odds ratio.

* Age was divided by a factor of 10 to indicate OR per decade increase in age.

DISCUSSION

This is the first study to comprehensively compare OIs among ART-naïve patients between four hospitals in Japan, northern Thailand, northern Vietnam, and the Philippines since the introduction of ART. The findings demonstrated considerable regional differences in the prevalence of major OIs, such as TB, PCP, and cryptococcosis, even after adjusting for age, sex, HIV-associated risk behavior, and CD4 count. We found a high prevalence of AIDS-defining OIs at our study sites despite the widespread availability of ART in all four countries during the study period. The distribution of OIs in each hospital mirrored the OI prevalences found in the respective reports by each country,^{8,13–16,24–26} indicating that our study population was representative.

TB is the most common HIV-related OI in Asia and is especially common in resource-limited settings.²⁷ The risk of TB in the four hospitals in our study paralleled the prevalence of PTB in the general population of each country; from 1990 to 2010, the estimated PTB prevalence was 502 per 100,000 individuals in the Philippines, 334 in Vietnam, 212 in Thailand, and 27 in Japan.²⁸ Interestingly, the incidence of TB has been shown to increase inversely proportionally to per capita gross domestic product.²⁹ The occurrence of TB among HIV-infected patients in developing regions is largely affected by the burden of TB in their surroundings,³⁰ as overcrowded living conditions place them at risk for frequent exposure to *Mycobacterium tuberculosis*.³¹ As previously reported,³² we found an increased prevalence of TB among IDUs. This increased risk can be attributed to the poor hygiene and ventilation in areas where injection drugs are used.³² The adverse immunomodulatory effects of illicit drugs have also been implicated in the increased susceptibility of IDUs to TB.³³

Consistent with national data from Japan, PCP was the most common AIDS-defining illness at NMC.¹⁴ Although PCP is considered prevalent in developed countries,³⁴ the aOR of PCP at PGH was similar to that at NMC in our study. In contrast, PCP was rare at BMH, which is consistent with previously published reports from Vietnam.^{15,16} *Pneumocystis jirovecii* is a ubiquitous pathogen, but its prevalence is highest in Europe and the United States.³⁵ The increased frequency of PCP diagnosis in developed countries could be due to differences in access to cytological examinations of respiratory pathogens, which are typically performed via bronchoalveolar

lavage; this differential access could contribute to the reportedly low prevalences of PCP found in lower-income countries.³⁴ Host genetic factors also affect susceptibility to PCP, although the precise mechanisms underlying these associations have not yet been illustrated. Possible protective genetic factors include apolipoprotein B mRNA-editing enzyme catalytic polypeptide-like 3F, CCR5-Delta32 mutation heterozygosity, and Dectin-1.^{36–38} Possible genetic factors increasing susceptibility to PCP include genotypes producing low mannose-binding lectin, CCRL2-F167Y mutation, and histidine-histidine homozygosity of the Fc segment receptors of IgG.^{39–41} A cohort study conducted in the Netherlands also indicated the potential impact of these host factors on PCP development.⁴² Therefore, these factors may have affected the regional differences in PCP identified in our study.

Penicilliosis was reported only at LPH and BMH; there were no known cases of penicilliosis at PGH or NMC during the study period. To date, only a few imported cases from Thailand have been reported in Japan.⁴³ Despite its typical presentation of skin lesions, penicilliosis is difficult to diagnose in nonendemic regions. A heightened awareness of the unique infections in neighboring countries is important given the recent increase in population mobility. Oropharyngeal candidiasis is one of the most common OIs in Asia¹² and was the most prevalent at NMC. The diagnosis of esophageal candidiasis was relatively rare at LPH and PGH, although its prevalence may have been underestimated due to limited access to upper endoscopy.

LPH showed the highest prevalence of cryptococcosis, which is consistent with the previously reported prevalence of cryptococcosis among AIDS patients in Thailand from 1994 to 1998.¹³ The endemicity of *Cryptococcus neoformans* in northern Thailand is typically attributed to frequent exposure to aerosolized feces from avian vectors.⁴⁴ A recent study from Chiang Mai University Hospital, Thailand, showed a lower prevalence of cryptococcal meningitis (5.2%),⁴⁵ likely concurrent with the recent improvements in living conditions and/or access to ART.

Consistent with previous reports from Europe, we did not find regional variations in CMV retinitis or herpes zoster.¹ However, the prevalence of CMV diseases in resource-limited settings was likely underestimated, as CMV infections other than retinitis were reported only at NMC. CMV viral load and antigen measurements are not routinely obtained in the work-up of HIV patients in resource-limited settings and should be a point of improvement in the next decade.

Most noninfectious AIDS-defining illnesses, lymphoma in particular, were diagnosed at NMC. Recently, the number of AIDS-defining malignancies (ADMs) has been increasing in Japan together with the increasing number of HIV-infected patients.¹⁴ Retrospective studies in Thailand have reported a small number of ADMs: Kiertiburanakul and others⁴⁶ found that 1.8% of HIV-infected patients had ADMs; Rojanawiwat and others⁴⁷ identified no cases of primary central nervous system (CNS) lymphoma despite frequent detection of Epstein-Barr virus in cerebrospinal fluid; and Chariyalertsak and others¹³ reported few cases of primary CNS lymphoma (0.1%) and Kaposi's sarcoma (0.2%) among the AIDS-defining illnesses occurring between 1994 and 1998. In Vietnam and the Philippines, information about ADMs is scarce. This trend in low prevalence of ADMs in resource-limited settings may be attributed to their nonspecific symptoms and/or the high mortality of patients with concomitant HIV-related OIs. HIV risk behaviors can strongly influence the prevalence of Kaposi's sarcoma. Human herpesvirus-8 (HHV-8), the oncovirus associated with Kaposi's sarcoma, is believed to be transmitted through anal intercourse. Hence, MSM are documented to have a higher risk of HHV-8 infection and Kaposi's sarcoma,^{48,49} which is consistent with the findings among patients at NMC and PGH, where MSM constituted the majority of the cohort.

This study has several limitations. First, the data collection was not performed simultaneously. Overall, patient information from 2003 to 2011 was used. However, this information was limited to 3 years at NMC and 1 year at BMH. Second, the numbers of sites and subjects were not sufficient to discuss the tendency of HIV infection across Asia. Third, the analyses were limited by the completeness of medical records, and the OI diagnoses may have been affected by variations in clinical practices at each study site. Fourth, the availability and accessibility of diagnostic procedures differed between NMC and the other three sites. To minimize this limitation, we focused the regression analyses on HIV-related illnesses that were less likely to be influenced by limited resources. Finally, the HIV subtypes of the study population were not available. Because our study sites are key major hospitals in each region, the HIV subtype was considered identical to its national tendency: the major HIV subtypes are B in Japan,⁵⁰ B and CRF01_AE in the Philippines,^{51,52} and CRF_01AE in Thailand and Vietnam.⁵³ There is no evidence that differences in subtype affect the incidence of OIs.

This is the first study to systematically compare the regional differences in major OIs in four Asian countries using logistic regression models. We found differences in the diagnoses of TB, PCP, and cryptococcosis between our study sites, whereas no regional differences in CMV retinitis or herpes zoster incidence were observed. These differences may be attributed to the geographical distribution of the pathogens, differences in HIV-related risk factors, clinical characteristics of the patients, and the availability of diagnostic procedures.

Received October 3, 2016. Accepted for publication February 22, 2017.

Published online April 24, 2017.

Note: Supplemental tables appear at www.ajtmh.org.

Acknowledgments: We would like to thank all the medical staff and research collaborators from the four study sites. Louie Mar A. Gangcuangco would like to acknowledge the Takeda Science

Foundation for the research fellowship grant supporting this project. The Nagasaki Institute of Tropical Medicine would like to thank Mediadora Sanie and Regina Berba from Philippine General Hospital for their participation in the 2010 Nagasaki Institute of Tropical Medicine Research Forum and their contribution to the Philippine General Hospital human immunodeficiency virus (HIV) cohort. We dedicate this manuscript to the HIV patients enrolled in our study.

Authors' addresses: Louie Mar A. Gangcuangco, Department of Clinical Tropical Medicine, Institute of Tropical Medicine (NEKKEN), Graduate School of Biomedical Science, Nagasaki University, Nagasaki, Japan, and Department of Internal Medicine, Bridgeport Hospital-Yale New Haven Health, Bridgeport, CT, E-mail: louiemarmd@gmail.com. Ikumi Sawada and Koya Ariyoshi, Department of Clinical Tropical Medicine, Institute of Tropical Medicine (NEKKEN), Graduate School of Biomedical Science, Nagasaki University, Nagasaki, Japan, E-mails: sawada.ikumi@gmail.com and kari@nagasaki-u.ac.jp. Naho Tsuchiya, Department of Clinical Tropical Medicine, Institute of Tropical Medicine (NEKKEN), Graduate School of Biomedical Science, Nagasaki University, Nagasaki, Japan, and Department of Preventive Medicine and Epidemiology, Tohoku Medical Megabank Organization, Tohoku University, Miyagi, Japan, E-mail: nahot@megabank.tohoku.ac.jp. Cuong D. Do and Thanh Thuy T. Pham, Department of Infectious Diseases, Bach Mai Hospital, Hanoi, Vietnam, E-mails: doduy.cuong@gmail.com and phamthanhthuy714@gmail.com. Archawin Rojanawiwat, Department of Medical Sciences, Ministry of Public Health, Nonthaburi, Thailand, E-mail: archawin191@yahoo.com. Marissa Alejandria and Katerina Leyritana, Section of Infectious Diseases, Department of Medicine, Philippine General Hospital, University of the Philippines, Manila, Philippines, E-mails: mmalejandria@gmail.com and kateleyritana@yahoo.com. Yoshiyuki Yokomaku, Department of Infectious Diseases and Immunology, National Hospital Organization Nagoya Medical Center, Nagoya, Japan, E-mail: yokomaku@nnh.hosp.go.jp. Panita Pathipvanich, Day Care Center, Lampang Hospital, Lampang, Thailand, E-mail: panitapa@yahoo.com.

REFERENCES

- Blaxhult A, Kirk O, Pedersen C, Dietrich M, Barton SE, Gatell JM, Mulcahy F, Hirschel B, Mocroft A, Lundgren JD, 2000. Regional differences in presentation of AIDS in Europe. *Epidemiol Infect* 125: 143–151.
- Nucci M, Queiroz-Telles F, Tobon AM, Restrepo A, Colombo AL, 2010. Epidemiology of opportunistic fungal infections in Latin America. *Clin Infect Dis* 51: 561–570.
- Handford CD, Rackal JM, Tynan AM, Rzeznikiewicz D, Glazier RH, 2012. The association of hospital, clinic and provider volume with HIV/AIDS care and mortality: systematic review and meta-analysis. *AIDS Care* 24: 267–282.
- Kaslow RA, Carrington M, Apple R, Park L, Munoz A, Saah AJ, Goedert JJ, Winkler C, O'Brien SJ, Rinaldo C, Detels R, Blattner W, Phair J, Erlich H, Mann DL, 1996. Influence of combinations of human major histocompatibility complex genes on the course of HIV-1 infection. *Nat Med* 2: 405–411.
- Pant Pai N, Shivkumar S, Cajas JM, 2012. Does genetic diversity of HIV-1 non-B subtypes differentially impact disease progression in treatment-naïve HIV-1-infected individuals? A systematic review of evidence: 1996–2010. *J Acquir Immune Defic Syndr* 59: 382–388.
- Ruxrungtham K, Brown T, Phanuphak P, 2004. HIV/AIDS in Asia. *Lancet* 364: 69–82.
- Yang CH, Huang YF, Hsiao CF, Yeh YL, Liou HR, Hung CC, Yang SY, 2008. Trends of mortality and causes of death among HIV-infected patients in Taiwan, 1984–2005. *HIV Med* 9: 535–543.
- AIDS Surveillance Committee, Ministry of Health Labour and Welfare, 2014. *HIV/AIDS Surveillance in 2014*. Available at: http://api-net.jfap.or.jp/status/2014/14nenpo/14nenpo_menu.html. Accessed April 27, 2016.
- Dickson N, McAllister S, Sharples K, Paul C, 2012. Late presentation of HIV infection among adults in New Zealand: 2005–2010. *HIV Med* 13: 182–189.

10. Kiertiburanakul S, Boonyarattaphun K, Atamasirikul K, Sungkanuparph S, 2008. Clinical presentations of newly diagnosed HIV-infected patients at a university hospital in Bangkok, Thailand. *J Int Assoc Physicians AIDS Care (Chic)* 7: 82–87.
11. Kong BN, Harwell JI, Suos P, Lynen L, Mohiuddin S, Reinert S, Pugatch D, 2007. Opportunistic infections and HIV clinical disease stage among patients presenting for care in Phnom Penh, Cambodia. *Southeast Asian J Trop Med Public Health* 38: 62–68.
12. Zhou J, Kumarasamy N, Ditangco R, Kamarulzaman A, Lee CK, Li PC, Paton NI, Phanuphak P, Pujari S, Vibhagool A, Wong WW, Zhang F, Chuah J, Frost KR, Cooper DA, Law MG, 2005. The TREAT Asia HIV Observational Database: baseline and retrospective data. *J Acquir Immune Defic Syndr* 38: 174–179.
13. Chariyalertsak S, Sirisanthana T, Saengwonloey O, Nelson KE, 2001. Clinical presentation and risk behaviors of patients with acquired immunodeficiency syndrome in Thailand, 1994–1998: regional variation and temporal trends. *Clin Infect Dis* 32: 955–962.
14. Yasuoka A, Tsukamoto M, Kurihara S, Takami Y, Shiki N, 2012. *A National Survey of HIV-Related Opportunistic Infections in 2010*. AIDS Research Project. Tokyo, Japan: Ministry of Health, Labour and Welfare, 16–34.
15. Louie JK, Chi NH, Thao LTT, Quang VM, Campbell J, Chau NV, Rutherford GW, Farrar JJ, Parry CM, 2004. Opportunistic infections in hospitalized HIV-infected adults in Ho Chi Minh City, Vietnam: a cross-sectional study. *Int J STD AIDS* 15: 758–761.
16. Klotz SA, Nguyen HC, Van Pham T, Nguyen LT, Ngo DT, Vu SN, 2007. Clinical features of HIV/AIDS patients presenting to an inner city clinic in Ho Chi Minh City, Vietnam. *Int J STD AIDS* 18: 482–485.
17. Section Bureau of Epidemiology, Department of Disease Control, Ministry of Public Health, 2008. *Epidemiological Information*. Bangkok, Thailand: Ministry of Public Health.
18. World Health Organization, 2007. *WHO Case Definitions of HIV for Surveillance and Revised Clinical Staging and Immunological Classification of HIV-Related Diseases in Adults and Children*. Available at: <http://www.who.int/hiv/pub/guidelines/hivstaging/en/>. Accessed May 4, 2015.
19. Ministry of Health Labour and Welfare, 2011. *Anti HIV Treatment Guideline*. Available at: <http://www.haart-support.jp/guideline.htm>. Accessed April 1, 2015.
20. Ministry of Public Health, 2000. *National Guidelines for the Clinical Management of HIV Infection in Children and Adults*, 6th Edition. Bangkok, Thailand: Ministry of Public Health.
21. Vietnam Ministry of Health, 2009. *Guidelines for HIV/AIDS Diagnosis and Treatment*. Available at: http://www.who.int/hiv/pub/guidelines/vietnam_art.pdf. Accessed June 4, 2015.
22. Philippine Society for Microbiology and Infectious Diseases, 2006. *Clinical Practice Guidelines for the Diagnosis, Treatment, Prevention and Control of Tuberculosis in Adult Filipinos: 2006 Update*. Available at: http://www.psmid.org.ph/pdf/Introduction_Edited_4.pdf. Accessed June 4, 2015.
23. Vanittanakom N, Cooper CR Jr, Fisher MC, Sirisanthana T, 2006. *Penicillium marneffei* infection and recent advances in the epidemiology and molecular biology aspects. *Clin Microbiol Rev* 19: 95–110.
24. Vietnam Authority of HIV/AIDS Control, 2014. *An Annual Update on the HIV Epidemic in Vietnam*. Available at: <http://www.aidsdatahub.org/annual-update-hiv-epidemic-viet-nam-vietnam-authority-hiv-aids-control-vaac-2014>. Accessed February 2, 2016.
25. Farr AC, Wilson DP, 2010. An HIV epidemic is ready to emerge in the Philippines. *J Int AIDS Soc* 13: 16.
26. HIV and AIDS Data Hub for Asia-Pacific, 2015. *Country Profiles*. Available at: <http://www.aidsdatahub.org>. Accessed May 11, 2015.
27. Trinh QM, Nguyen HL, Nguyen VN, Nguyen TV, Sintchenko V, Marais BJ, 2015. Tuberculosis and HIV co-infection-focus on the Asia-Pacific region. *Int J Infect Dis* 32: 170–178.
28. World Health Organization, 2012. *Global Tuberculosis Control: WHO Report 2011*. Available at: http://apps.who.int/iris/bitstream/10665/44728/1/9789241564380_eng.pdf. Accessed February 2, 2016.
29. Janssens JP, Rieder HL, 2008. An ecological analysis of incidence of tuberculosis and per capita gross domestic product. *Eur Respir J* 32: 1415–1416.
30. Kufa T, Mabuto T, Muchiri E, Charalambous S, Rosillon D, Churchyard G, Harris RC, 2014. Incidence of HIV-associated tuberculosis among individuals taking combination anti-retroviral therapy: a systematic review and meta-analysis. *PLoS One* 9: e111209.
31. Narasimhan P, Wood J, Macintyre CR, Mathai D, 2013. Risk factors for tuberculosis. *Pulm Med* 2013: 828939.
32. Deiss RG, Rodwell TC, Garfein RS, 2009. Tuberculosis and illicit drug use: review and update. *Clin Infect Dis* 48: 72–82.
33. Friedman H, Newton C, Klein TW, 2003. Microbial infections, immunomodulation, and drugs of abuse. *Clin Microbiol Rev* 16: 209–219.
34. Lowe DM, Rangaka MX, Gordon F, James CD, Miller RF, 2013. *Pneumocystis jirovecii* pneumonia in tropical and low and middle income countries: a systematic review and meta-regression. *PLoS One* 8: e69969.
35. Smulian AG, Sullivan DW, Linke MJ, Halsey NA, Quinn TC, MacPhail AP, Hernandez-Avila MA, Hong ST, Walzer PD, 1993. Geographic variation in the humoral response to *Pneumocystis carinii*. *J Infect Dis* 167: 1243–1247.
36. An P, Penugonda S, Thorball CW, Barthol, Goedert JJ, Donfield S, Buchbinder S, Binns-Roemer E, Kirk GD, Zhang W, Fellay J, Yu XF, Winkler CA, 2016. Role of APOBEC3F gene variation in HIV-1 disease progression and *Pneumocystis* pneumonia. *PLoS Genet* 12: e1005921.
37. Ashton LJ, Stewart GJ, Biti R, Law M, Cooper DA, Kaldor JM, 2002. Heterozygosity for CCR5-Delta32 but not CCR2b-64i protects against certain intracellular pathogens. *HIV Med* 3: 91–96.
38. Saijo S, Fujikado N, Furuta T, Chung SH, Kotaki H, Seki K, Sudo K, Akira S, Adachi Y, Ohno N, Kinjo T, Nakamura K, Kawakami K, Iwakura Y, 2007. Dectin-1 is required for host defense against *Pneumocystis carinii* but not against *Candida albicans*. *Nat Immunol* 8: 39–46.
39. Yanagisawa K, Ogawa Y, Uchiumi H, Gohda F, Mawatari M, Ishizaki T, Mitsui T, Yokohama A, Handa H, Tsukamoto N, Nojima Y, 2015. Gene polymorphisms of mannose-binding lectin confer susceptibility to *Pneumocystis* pneumonia in HIV-infected patients. *J Infect Chemother* 21: 769–775.
40. An P, Li R, Wang JM, Yoshimura T, Takahashi M, Samudralal R, O'Brien SJ, Phair J, Goedert JJ, Kirk GD, Troyer JL, Sezgin E, Buchbinder SP, Donfield S, Nelson GW, Winkler CA, 2011. Role of exonic variation in chemokine receptor genes on AIDS: CCR2 F167Y association with *Pneumocystis* pneumonia. *PLoS Genet* 7: e1002328.
41. Forthal DN, Landucci G, Bream J, Jacobson LP, Phan TB, Montoya B, 2007. FcγRIIIa genotype predicts progression of HIV infection. *J Immunol* 179: 7916–7923.
42. Schoffelen AF, van Lelyveld SF, Barth RE, Gras L, de Wolf F, Netea MG, Hoepelman AI, 2013. Lower incidence of *Pneumocystis jirovecii* pneumonia among Africans in The Netherlands host or environmental factors? *AIDS* 27: 1179–1184.
43. Kamei K, Sano A, Kikuchi K, Makimura K, Niimi M, Suzuki K, Uehara Y, Okabe N, Nishimura K, Miyaji M, 2003. The trend of imported mycoses in Japan. *J Infect Chemother* 9: 16–20.
44. Kuroki M, Phichaichumpon C, Yasuoka A, Chiranairadul P, Chosa T, Sirinirund P, Miyazaki T, Kakeya H, Higashiyama Y, Miyazaki Y, Ishida Y, Kohno S, 2004. Environmental isolation of *Cryptococcus neoformans* from an endemic region of HIV-associated cryptococcosis in Thailand. *Yeast* 21: 809–812.
45. Chaiwarith R, Vongsanim S, Supparatpinyo K, 2014. Cryptococcal meningitis in HIV-infected patients at Chiang Mai University Hospital: a retrospective study. *Southeast Asian J Trop Med Public Health* 45: 636–646.
46. Kiertiburanakul S, Likhitpongwi S, Ratanasiri S, Sungkanuparph S, 2007. Malignancies in HIV-infected Thai patients. *HIV Med* 8: 322–323.

47. Rojanawiwat A, Miura T, Thaisri H, Pathipvanich P, Umnajsirisuk S, Koibuchi T, Vongsheree S, Iwamoto A, Ariyoshi K, Sawanpanyalert P, 2005. Frequent detection of Epstein-Barr virus and cytomegalovirus but not JC virus DNA in cerebrospinal fluid samples from human immunodeficiency virus-infected patients in northern Thailand. *J Clin Microbiol* 43: 3484–3486.
48. Ebrahim SH, Peterman TA, Zaidi AA, Hamers FF, 1997. Geography of AIDS-associated Kaposi's sarcoma in Europe. *AIDS* 11: 1739–1745.
49. Rohner E, Wyss N, Heg Z, Faralli Z, Mbulaiteye SM, Novak U, Zwahlen M, Egger M, Bohlius J, 2016. HIV and human herpesvirus 8 co-infection across the globe: systematic review and meta-analysis. *Int J Cancer* 138: 45–54.
50. Sugiura W, 2009. Subtypes of HIV prevalent in Japan, 2003–2007. *Infect Agents Surv Rep* 30: 234–235.
51. Paladin FJ, Monzon OT, Tsuchie H, Aplasca MR, Learn GH Jr, Kurimura T, 1998. Genetic subtypes of HIV-1 in the Philippines. *AIDS* 12: 291–300.
52. Schwem B, Itable J, Ching P, Ganchua S, Alejandria M, Lim J, Destura R, Salvana EM, 2014. *The Changing Molecular Epidemiology of HIV in the Philippines: Major Shift from Subtype B to CRF01_AE. ID Week 2014*. Philadelphia, PA: Infectious Diseases Society of America.
53. Hemelaar J, Gouws E, Ghys PD, Osmanov S, 2006. Global and regional distribution of HIV-1 genetic subtypes and recombinants in 2004. *AIDS* 20: W13–W23.