Published in final edited form as:

Psychiatry Res. 2014 April 30; 222(1-2): 107–110. doi:10.1016/j.psychres.2014.02.003.

Face processing in depersonalization: An fMRI study of the unfamiliar self

Sarah Ketay^{a,*}, Holly K. Hamilton^b, Brian W. Haas^c, and Daphne Simeon^d

^aDepartment of Psychology, University of Hartford, West Hartford, CT, USA

^bDepartment of Psychology, University of California at Los Angeles, Los Angeles, CA, USA

^cDepartment of Psychology, University of Georgia, Athens, Georgia, USA

^dDepartment of Psychiatry, Mount Sinai School of Medicine, New York, NY, USA

Abstract

Depersonalization disorder (DPD) is characterized by a core sense of unfamiliarity. Nine DPD participants and 10 healthy controls underwent functional magnetic resonance imaging while viewing self and unfamiliar faces. Compared with control subjects, the DPD group exhibited significantly greater activation in several brain regions in response to self vs. stranger faces. Implications are discussed.

Keywords

Depersonalization; Self; Unfamiliarity

1. Introduction

Two core clinical features of DSM-IV-TR depersonalization disorder (DPD) are a subjective sense of hypoemotionality and unfamiliarity, despite intact reality testing. *Corticolimbic disconnections* (Sierra and Berrios, 1998) involving heightened prefrontal cortical activation and reciprocal limbic inhibition may account for the hypoemotionality and has largely been the focus of neuroimaging studies (Medford, 2012). Knowledge of the neural correlates of unfamiliarity in DPD (feeling separate from oneself, unreality of the self), on the other hand, is limited. One positron emission tomography study suggested that disruptions in posterior sensory associative cortical functioning may mediate multimodal perceptual disturbances leading to altered self-perception (Simeon et al., 2000).

The self-face paradigm has been used to examine the neural basis of the self. A review of neuroimaging studies indicates a right-dominated yet distributed network of self-processing

Corresponding Author: Sarah Ketay, Ph.D., Department of Psychology, University of Hartford, 200 Bloomfield Avenue, West Hartford, CT 06117, USA, Telephone: 860.768.5906, Fax: 860.768.5292, ketay@hartford.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

with consistent involvement of the left fusiform gyrus, bilateral middle and inferior frontal gyri, and parietal cortical regions (Devue and Bredart, 2011). Further studies reveal the superior parietal lobe, right inferior temporal gyrus and the occipital face area, consisting of the superior parietal lobule and the inferior occipital gyrus, are associated with self-face recognition in healthy populations (Uddin et al., 2005; Apps et al., 2012). There is also evidence for a left-hemisphere-dominated network for self-face recognition when viewing a self-face transitioning to a familiar face (Turk et al., 2002). Activation in the medial prefrontal cortex (MedPFC) is consistently reported in neuroimaging studies of self vs. other judgments, yet the precise role of this region is unclear. Greater activity in ventral MedPFC was shown during trait judgments of self vs. other (Kelley et al., 2002), and MedPFC regions involved in making judgments for self and other may overlap (Ochsner et al., 2005). Activation has been found in the right prefrontal cortex during explicit discrimination between self and other (Platek et al., 2006). Furthermore, the right anterior cingulate and bilateral prefrontal cortices were found to activate during active vs. passive self-face processing (Sugiura et al., 2000).

To date, there are limited neuroimaging studies examining self-recognition in patient populations. One study highlights the neural systems underlying shared representation of self and other in children with autism (Uddin et al., 2008). In response to a self-recognition task involving morphed images, the right frontal region was activated in typically developing children for self and other faces. This region was activated for self faces, but not other faces, suggesting that children with autism lack a shared neural representation for self and others.

Interpreting and integrating these activations has proved difficult due to varying control stimuli (e.g., familiar, unfamiliar, famous, romantic partner face, trait judgments) and experimental tasks (e.g., photograph content, stimuli presentation, implicit vs. explicit tasks). Most neuroimaging research on self to date involves supraliminal or explicit tasks. Recent research on self-processing at multiple levels of awareness, using both subliminal and supraliminal self-face stimuli, shows that self-related stimuli enjoy the self-advantage of a more robust neural response even when processed subliminally (Geng et al., 2012).

The present functional magnetic resonance imaging (fMRI) study is the first to examine unfamiliarity of self in DPD during implicit self vs. other (stranger) face processing. We hypothesized that DPD participants would demonstrate altered activation in “self” brain regions compared with healthy controls reflecting altered implicit self-processing, in the presence of intact supraliminal self-recognition.

2. Methods

2.1. Participants

The study was approved by the institutional review board at Mount Sinai School of Medicine. Participants were medically and neurologically healthy right-handed adults aged 18 – 55. The DPD participants, $n=9$, were assessed by structured interviews (SCID-I/P, First et al., 2002; SCID-II, First et al., 1997) for Axis I and II disorders, and SCID-D (Steinberg, 1994) for dissociative disorders. Control participants, $n=10$, scored <10 on the Dissociative

Experiences Scale (Bernstein and Putnam, 1986) and had no lifetime Axis I or II disorders. All participants provided written informed consent.

2.2. Task

Participants underwent fMRI while viewing photographs of their own faces and a stranger's face. Participants' photographs were facing forward and at a 45° angle with neutral expression and cropped above the shoulders using Adobe Photoshop. Stranger photos, matched for gender, race, age, facial expression, and angle, were found via internet search. Two 6-min runs were conducted, each presenting twenty 18-s blocks consisting of six self and stranger facial stimuli in pseudo-random order. Participants judged whether faces were facing the camera via a button press.

2.3. MRI acquisition

Neuroimaging data were collected using a 3T GE Siemens Allegra MR scanner. See Supplementary Materials for data-acquisition parameters.

2.4. Image processing and analyses

Images were processed as follows: (1) realigned correct for head motion and coregistered to their respective T2 image; (2) spatially normalized to a standard template (MNI: Montreal Neurological Institute) using a 12-parameter affine registration algorithm; and (3) smoothed with an 8×8×8 mm Gaussian kernel. Experimental condition effects were estimated according to the general linear model using a canonical hemodynamic response function. Fixed effects models (Friston, 1994) were used at the individual subject level of analysis, and random effects models (Holmes and Friston, 1998) were used for group level analyses. On an individual subject level, *t*-contrast images (fixed effects) were calculated based on the difference between the self and stranger conditions (self > stranger). To determine areas that were significantly different between groups, a two-sample *t*-test (random effects) was performed. All fMRI analyses were corrected for multiple comparisons. For whole-brain analyses, we corrected for multiple comparisons using AFNI's 3dClustSim based on 1000 Monte Carlo simulations; with the use of a combined height ($P < 0.001$) and spatial-extent (93 voxels) threshold, the significance level was established at $P < 0.05$ corrected. Imaging analyses were performed with SPM 5 (Wellcome Department of Imaging Neuroscience, London, UK) in MATLAB 7.0 (Mathworks, Natick, MA).

Additionally, we conducted exploratory Pearson's correlation analyses between the Cambridge Depersonalization Scale (total CDS score) (Sierra and Berrios, 2000) and reactivity to self vs. stranger within the areas of interest using a combined $P=0.005$ (uncorrected), 20-voxel extent threshold (Lieberman and Cunningham, 2009).

3. Results

The two groups did not significantly differ in gender ($\chi^2=1.82$, $P=0.188$) or age (DPD: $M=33.2$, $SD=11.2$, HC: $M=31.9$, $SD=11.8$, $t(15)=0.54$, $P=0.466$). The two groups did not differ in reaction time when viewing self, $t(15)=1.66$, $P=0.18$, or stranger, $t(15)=1.77$, $P=0.10$, or in response accuracy for self, $t(14)=-1.3$, $P=0.21$, or stranger, $t(14)=1.40$, $P=0.18$.

Compared with controls, DPD participants exhibited greater response to self vs. stranger processing within two large clusters (Fig. 1A). One cluster was located within the right anterior cingulate (124 voxels) (ACC, BA 24; $z = 4.45$; peak voxel: MNI: 2, 24, 16). The second cluster extended into three brain regions: left MedPFC (469 voxels) (L. MedPFC, BA 6/9; $z = 4.26$; peak voxel: MNI: -18, 32, 34), right medial prefrontal cortex (R. MedPFC; BA 6/9; $z = 3.92$; peak voxel: MNI: 2, 30, 36) and left middle frontal gyrus (L. MFG; BA 6; $z = 5.15$; peak voxel: MNI: -12, 16, 46). Standardized contrast estimates extracted from the peak voxel for each brain region are presented in Fig. 1B. There were no areas of significantly greater response to self vs. stranger processing in controls compared with the DPD group.

Within the DPD group, higher CDS scores were positively correlated with reactivity to self vs. stranger faces in the left MedPFC (peak voxel: MNI: -12, 10, 54; 63 voxels; $t = 7.13$; $P < 0.001$) and the left Middle FG (peak voxel: MNI: -24, 38, 22; 119 voxels; $t = 7.10$; $P < 0.001$).

4. Discussion

DPD participants exhibited greater response to self vs. stranger processing within the right ACC, bilateral MedPFC, and left middle frontal gyrus, and depersonalization severity was significantly associated with activation in the latter two regions. Broadly, findings are consistent with a reported positive correlation between depersonalization severity and right frontal cortex and ACC activation in healthy individuals with intravenous tetrahydrocannabinol-induced depersonalization (Mathew et al., 1999). Findings are also consistent with increased ACC and right prefrontal and frontal activation in PTSD patients experiencing an imagery-induced dissociative state (Lanius et al., 2002). Explanations of our findings not related to self-processing cannot be excluded. The two groups could be demonstrating different brain activation secondary to different cognitive task demands. However, the comparable reaction times and recognition accuracy in the two groups makes this explanation less likely. Furthermore, the positive relationship between depersonalization severity and brain activation in the MedPFC also points to a deficit in implicit self-processing.

Another alternative interpretation of the findings related to ACC activation could involve a conflict between conscious and unconscious processing of self in DPD. Although individuals with DPD have no difficulty recognizing the self-face at the conscious level, this could be coupled with an unconscious response of “not-knowing” the self, thus activating conflict detection and resolution in the ACC (Botvinick et al. 2004), as has been shown in another dissociative condition, conversion disorder (van Beilen et al., 2010).

Finally, the two core features of depersonalization, hypoemotionality and unfamiliarity, may be interrelated. Neural pathways for perceptual recognition are likely linked to parallel pathways for assigning emotional content to percepts. As such, abnormal self-face processing may underlie the lack of emotional resonance. The association between hypoemotionality and unfamiliarity in depersonalization warrants further investigation.

There are limitations to this initial study, which necessitate and motivate further investigation, including the small sample size, the task used, and the control condition. The use of a famous, versus a stranger, face as a control would address familiarity as a confounding factor. The subtraction method presumes that self-face minus stranger indicates self-face activation; however, related cognitive processes may be activated during control.

Understanding the neural underpinnings of unfamiliarity holds the potential to inform novel treatment approaches for DPD. Target sites for repetitive transcranial magnetic stimulation treatment have been selected based on prior neuroimaging findings (Mantovani et al., 2011). Further investigation of the neural basis of the depersonalized self-experience is warranted.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by grant #UL1TR000067 from the National Center for Research Resources, National Institutes of Health

References

- Apps MA, Tajadura-Jiménez A, Turley G, Tsakiris M. The different faces of one's self: an fMRI study into the recognition of current and past self-facial appearances. *Neuroimage*. 2012; 63(3):1720–1729. [PubMed: 22940117]
- Bernstein EM, Putnam FW. Development, reliability, and validity of a dissociation scale. *Journal of Nervous & Mental Disease*. 1986; 174:727–735. [PubMed: 3783140]
- Botvinick MM, Cohen JD, Carter CS. Conflict monitoring and anterior cingulate cortex: an update. *Trends in Cognitive Science*. 2004; 8:539–546.
- Devue C, Brédart S. The neural correlates of visual self-recognition. *Consciousness and Cognition*. 2011; 20:40–51. [PubMed: 20880722]
- First, MB., Gibbon, M., Spitzer, RL., Williams, JBW., Benjamin, LS. Structured Clinical Interview for DSM-IV Axis II Personality Disorders (SCID-II). American Psychiatric Press; Washington, DC: 1997.
- First, MB., Spitzer, RL., Gibbon, M., Williams, JBW. Structured Clinical Interview for DSM-IV (SCID-I) - Research Version. Biometrics Research Department, New York State Psychiatric Institute; New York, NY: 2002.
- Friston, KJ. Statistical parametric mapping. In: Thatcher, RW, Hallett, M, Zeffiro, T, John, ER., Huerta, M., editors. *Functional Neuroimaging: Technical Foundations*. Academic Press; San Diego: 1994. p. 79-93.
- Geng H, Zhang S, Li Q, Tao R, Xu S. Dissociations of subliminal and supraliminal self-face from other-face processing: behavioral and ERP evidence. *Neuropsychologia*. 2012; 50:2933–2942. [PubMed: 22898645]
- Holmes AP, Friston KJ. Generalisability, random effects and population inference. *NeuroImage*. 1998; 7:S754.
- Kelley WM, Macrae CN, Wyland CL, Caglar S, Inati S, Heatherton TF. Finding the self? An event-related fMRI study. *Journal of Cognitive Neuroscience*. 2002; 14:785–794. [PubMed: 12167262]
- Lanius RA, Williamson PC, Boksman K, Densmore M, Gupta M, Neufeld RW, Gati JS, Menon R. Brain activation during script-driven imagery induced dissociative responses in PTSD: a functional MRI investigation. *Biological Psychiatry*. 2002; 52:305–311. [PubMed: 12208637]
- Lieberman MD, Cunningham WA. Type I and Type II error concerns in fMRI research: re-balancing the scale. *Social Cognitive and Affective Neuroscience*. 2009; 4:423–428. [PubMed: 20035017]

- Mantovani A, Simeon D, Urban N, Bulow P, Allart A, Lisanby S. Temporo-parietal junction stimulation in the treatment of DPD. *Psychiatry Research*. 2011; 186:138–140. [PubMed: 20837362]
- Mathew RJ, Wilson WH, Chiu NY, Turkington TG, Degrado TR, Coleman RE. Regional cerebral blood flow and depersonalization after tetrahydrocannabinol administration. *Acta Psychiatrica Scandinavica*. 1999; 100:67–75. [PubMed: 10442442]
- Medford N. Emotion and the unreal self: Depersonalization disorder and deaffectualization. *Emotion Review*. 2012; 4:2139–2144.
- Ochsner KN, Beer JS, Robertson ER, Cooper JC, Kihlstrom JF, D'Esposito M, Gabrieli JDE. The neural correlates of direct and reflected self-knowledge. *NeuroImage*. 2005; 28:797–814. [PubMed: 16290016]
- Platek SM, Loughhead JW, Gur RC, Busch S, Ruparel K, Phend N, Panyavin IS, Langleben DD. Neural substrates for functionally discriminating self-face from personally familiar faces. *Human Brain Mapping*. 2006; 27:91–98. [PubMed: 16035037]
- Sierra M, Berrios GE. Depersonalization: neurobiological perspectives. *Biological Psychiatry*. 1998; 44:898–908. [PubMed: 9807645]
- Sierra M, Berrios G. The Cambridge Depersonalisation Scale: a new instrument for the measurement of depersonalization. *Psychiatry Research*. 2000; 93:153–164. [PubMed: 10725532]
- Simeon D, Guralnik O, Hazlett E, Spiegel-Cohen J, Hollander E, Buchsbaum MS. Feeling unreal: a PET study of depersonalization disorder. *The American Journal of Psychiatry*. 2000; 157:1782–1788. [PubMed: 11058475]
- Steinberg, M. *The Structured Clinical Interview for DSM-IV Dissociative Disorders-Revised (SCID-D)*. American Psychiatric Press; Washington, DC: 1994.
- Sugiura M, Kawashima R, Nakamura K, Okada K, Kato T, Nakamura A, Hatano K, Itoh K, Kojima S, Fukuda H. Passive and active recognition of one's own face. *NeuroImage*. 2000; 11:36–48. [PubMed: 10686115]
- Turk DJ, Heatherton TF, Kelley WM, Funnell MG, Gazzaniga MS, Macrae CN. Mike or me? Self-recognition in a split-brain patient. *Nature Neuroscience*. 2002; 5:841–842. [PubMed: 12195428]
- Uddin LQ, Davies MS, Scott AA, Zaidel E, Bookheimer SY, Iacoboni M, Dapretto M. Neural basis of self and other representation in autism: an FMRI study of self-face recognition. *PLoS One*. 2008; 3(10):e3526. [PubMed: 18958161]
- Uddin LQ, Kaplan JT, Molnar-Szakacs I, Zaidel E, Iacoboni M. Self-face recognition activates a frontoparietal “mirror” network in the right hemisphere: an event-related fMRI study. *Neuroimage*. 2005; 24(3):926–935.
- van Beilen M, Vogt B, Leenders K. Increased activation in cingulate cortex in conversion disorder: what does it mean? *Journal of Neurological Sciences*. 2010; 289:155–158.

Figure 1.

Areas of significantly greater response to self versus stranger processing in depersonalization personality disorder (DPD) as compared to healthy controls (HC). Areas of significant activation (A), using a whole brain analysis, are overlaid on a standardized template brain, in neurological orientation, on axial slices ranging between $z = 18$ to $z = 44$ (MNI). Plots (B) present mean contrast estimates for the comparison between self versus stranger processing, for each group. Error bars represent standard error from the mean. L = Left, R = Right; MedPFC = medial prefrontal cortex, ACC = anterior cingulate cortex, MFG = middle frontal gyrus.