Short Communication: Resolution of Tenofovir Disoproxil Fumarate Induced Fanconi Syndrome with Switch to Tenofovir Alafenamide Fumarate in a HIV-1 and Hepatitis B Coinfected Patient

Maile Young Karris

Abstract

Fanconi syndrome is a rare adverse effect of tenofovir disoproxil fumarate (TDF). Tenofovir alafenamide fumarate (TAF) is a novel prodrug with less nephrotoxicity. We report resolution of Fanconi syndrome in a HIV and hepatitis B coinfecting patient switched from TDF to TAF.

Keywords: HIV, hepatitis B, tenofovir alafenamide, Fanconi syndrome

Introduction

TENOFOVIR DISOPROXIL FUMARATE (TDF) is a prodrug of tenofovir (TFV) that is approved for the chronic management of HIV and Hepatitis B virus (HBV). In most persons with HIV and HBV coinfection, TDF is not only highly efficacious but also very well tolerated.² In persons living with HIV and HBV, TDF has been associated with a small increased risk of nephrotoxicity that in rare circumstances progresses to a severe renal tubulopathy that is characterized by phosphaturia and subsequent hypophosphatemia, renal glucosuria, aminoaciduria, tubular proteinuria, and proximal renal tubular acidosis (Fanconi syndrome).¹ TDF mediated nephrotoxicity is thought to be due to accumulation of TDF in proximal tubules of the kidneys resulting in local mitochondrial toxicity.³ This effect appears exacerbated by increased plasma TFV concentrations.⁴

In persons living with HIV, the new prodrug tenofovir alafenamide fumarate (TAF) demonstrates an equivalent *in vivo* potency to TDF at 30-fold lower the dose resulting in 91% lower plasma TFV concentrations.⁵ This results in a more favorable safety and tolerability profile of TAF containing regimens compared to TDF.⁶ Pozniak *et al.* recently published a study evaluating HIV infected persons with mild to moderate renal impairment (estimated creatinine clearance or CrCl 30–69 ml/min) who switched from variable anti-retroviral therapy (ART) regimens (65% on TDF, 22% on abacavir, and 5% on a nucleos(t)ide-free regimen) to daily TAF coformulated with cobicistat, elvitegravir, and em-

tricitabine (EVG/COBI/FTC/TAF).⁷ In the participants enrolled, there was minimal change in estimated glomerular filtration rate (eGFR), but significant improvements in proteinuria, albuminuria, and bone mineral density. Despite the growing body of literature supporting the safer profile of TAF compared to TDF in HIV, less is known about the safety and efficacy of TAF in persons coinfecting with HIV and HBV or in those with history of TDF-associated renal disease in real-world experiences. In this study we describe an experience with TAF in a patient with HIV and HBV who developed TDF associated renal tubulopathy.

Case Report

A 61-year-old man with HIV and HBV coinfection and chronic kidney disease stage 3 was transferred to our clinic in April 2014 on ritonavir-boosted fosamprenavir and lamivudine/abacavir with an undetectable HIV viral load and CD4 T cell count 320 cells/ μ L (36%). He was diagnosed with HIV in 1984 and started on zidovudine monotherapy followed by didanosine+dideoxycytidine and other regimens that he could not specifically recall. In this first visit, he reported the recent discontinuation of TDF due to kidney injury. He was treated with TDF in 2008 and recalled it caused kidney injury then as well. Initial laboratory evaluation revealed serum creatinine (Cr) of 1.81 mg/dl (normal range 0.67–1.17 mg/dl) and eGFR of 41.5 ml/min (by Cockcroft–Gault equation). Urinalysis was positive for 1+ protein, 2+ glucose (in the setting of normal serum glucose), and HBV DNA 278 IU/ml.

TABLE 1. LABORATORY TESTS AND MEDICATIONS FROM APRIL 2014 TO JUNE 2016

Laboratory tests	March 2014	April 2014	May 2014	August 2014	September 2014	January 2015	August 2015	September 2015	November 2015	December 2015	March 2016	June 2016	Medications
													FPV/r+3TC/ABC TDF
													FTC/TAF
													DRV/r+DTG
Creatinine (0.67–1.17 mg/dl)													
eGFR ^a ml/min by Cockcroft–Gault equation		1.81 41.5	1.84 40.8	1.66 44.9	1.91 39.1	1.74 40	2.1 33	1.43 60	1.93 42	1.66 42	1.74 40	1.82 41	
Alkaline phos. (40–129 U/liter)													
ALT (0–41 U/liter)		57 30	57 27	104 251	53 42	54 39	54 22	61 18	56 37	49 31	71 36	51 17	
AST (0–40 U/liter)		29	26	119	29	24	23	23	27	31	27	17	
Phosphorous (2.7–4.5 mg/dl)				3.9		3				2.4	4	3.3	
Hepatitis B DNA		278		>3 × 10 ⁶	1 × 10 ⁵	132	<20	3920	<20	<20	UD	UD	
HIV viral load ^b copies/ml		UD		UD		UD	<20	<20	<20	<20	UD	UD	
Dipstick protein		1		1	1	1	1	tr	1	1	neg	neg	
Dipstick glucose		2		neg	neg	small	neg	tr	neg	neg	neg	neg	
Urine Protein: Cr		0.57	0.55	0.21		0.29				0.27	0.18	0.19	
Urine β ₂ : Cr (0–300 μg/g Cr)			49,950									15,822	

Time on medications indicated by shading. Normal laboratory reference ranges in parenthesis. Blank values indicate tests not performed at that interval.

^aeGFR calculated by Cockcroft–Gault equation.

^bLaboratory reports HIV viral load as undetectable when there is no signal and <20 when a positive signal exists but cannot be quantified.

3TC, lamivudine; ABC, abacavir; ALT, alanine aminotransferase; AST, aspartate aminotransferase; DRV, darunavir; DTG, dolutegravir; eGFR, estimated glomerular filtration rate; FPV, fosamprenavir; FTC, emtricitabine; Neg, negative; r, ritonavir; TAF, tenofovir disoproxil fumarate; tr, trace; UD, undetectable.

Liver enzymes were normal. HBV genotype suggested resistance to lamivudine and entecavir, but sensitivity to TFV and adefovir [drug resistance mutations L180M, S202G, and M204V of the HBV polymerase region (Pol/RT)].

Nephrology evaluation confirmed that his presentation was consistent with Fanconi syndrome likely due to TDF after ruling out other possible causes and recommended avoidance of future TDF use. He was also not deemed a candidate for adefovir given chronic renal dysfunction and the ongoing risk of renal tubulopathy and hepatology consultation expressed concern that interferon therapy may result in hepatic decompensation. Thus the patient remained off TDF for the next 5 months with close monitoring. Off TDF his kidney function slightly improved to Cr 1.67 mg/dl (GFR 45.6 ml/min) with urinalysis demonstrating 1+ protein but no glycosuria. However his HBV DNA increased to over 3×10^6 IU/ml accompanied by a rise in alanine aminotransferase (ALT) from 29 to 251 U/liter and aspartate aminotransferase (AST) from 19 to 119 U/liter. Hepatology evaluation confirmed cirrhosis with HBV flare and determined he was not a candidate for liver transplant due to intermittent methamphetamine use and a lack of social support. Given the risk of death due to HBV-associated liver disease, TDF coformulated with emtricitabine (FTC/TDF) was resumed at a renally adjusted interval (thrice a week) with a plan to eventually transition to chronic hemodialysis.

At this point, compassionate use of TAF was pursued. On April 1, 2015 the U.S. Food and Drug Administration approved a single patient investigational new drug application (No. 126033). The patient gained access to TAF in the form of FTC/TAF September 2015 after approval from the local institutional review board.

While awaiting access to TAF, a GenoSure Archive (Monogram Biosciences, San Francisco, CA) was performed that demonstrated class-wide nucleos(t)ide reverse transcriptase

inhibitor drug resistance. He was switched to ritonavir-boosted darunavir (DRV/r), with the addition of dolutegravir (DTG) and continued on FTC/TDF thrice a week. As expected on TDF, his kidney dysfunction slowly progressed with Cr 2.1 mg/dl (GFR 35 ml/min) accompanied by low grade proteinuria and glycosuria, but liver function tests normalized in December 2014 and HBV became undetectable in August 2015 (Table 1 and Fig. 1). On August 31, 2015 the patient stopped all HIV medications, including FTC/TDF, in response to acutely worsening kidney function and in anticipation of imminent FTC/TAF start. Laboratory tests ordered while off all antiretrovirals, including TDF and just before start of TAF, revealed improved kidney function (Cr 1.43 mg/dl and GFR 61 ml/min with detectable trace proteinuria and glycosuria) but detectable HBV (3,920 IU/ml). He restarted DRV/r+DTG and initiated daily FTC/TAF (200 mg of FTC and 10 mg of TAF) on September 20, 2015.

On F/TAF, he demonstrated a slight improvement in his GFR (40–46 ml/min) and 9 months after switching had improvement of proteinuria (by urinalysis and quantitative protein to creatinine and β_2 microglobulin to creatinine ratios) and resolution of glycosuria. Of note urine protein to creatinine and β_2 microglobulin to creatinine ratios were used in concert with urinalysis as a more accurate assay for proteinuria and renal tubular damage. He did not complain of any notable adverse effects with TAF initiation. HBV became quickly undetectable and he did not experience another HBV flare.

Discussion

This case represents the first published experience of use of TAF in a HIV and HBV coinfecting patient with TDF induced renal tubulopathy. FTC/TAF was well tolerated and did result in both HIV and HBV virologic control. As consistent with TAF use in persons with chronic kidney disease, the

FIG. 1. Graphical representation of the patients' laboratory tests and timing of antiretrovirals. Months are represented by number (i.e., March is 3). 3TC, lamivudine; ABC, abacavir; ALT, alanine aminotransferase; AST, aspartate aminotransferase; Cr, creatinine; DRV, darunavir; DTG, dolutegravir; eGFR, estimated glomerular filtration rate calculated by Cockcroft–Gault equation; FPV, fosamprenavir; FTC, emtricitabine; Glu, glucose; HBV, hepatitis B virus; Pro, protein; r, ritonavir; TAF, tenofovir alafenamide; TDF, tenofovir disoproxil fumarate.

eGFR did not significantly change with switch to TAF, but tubular damage did improve as evident by glycosuria (by urinalysis) and proteinuria (by urinalysis and protein to creatinine ratios).⁷

TAF is believed to have an improved safety profile due to (1) equivalent efficacy at significantly lower plasma levels of TAF and subsequently plasma TFV and (2) unlike TDF, TAF does not appear to be a substrate of organic anion transporters that are thought to contribute to the accumulation of TFV in the proximal tubules.⁸ In a recent study evaluating virologically suppressed patients with renal impairment, a small proportion of patients (2% [5/242]) discontinued EVG/COBI/FTC/TAF due to decreased creatinine clearance, but none had evidence of proximal tubulopathy. However, these cases are confounded as all had a secondary reason for progression such as uncontrolled hypertension. Interestingly, in this study, two participants with a history of TDF induced Fanconi syndrome demonstrated stable eGFR and had reductions in proteinuria.⁹ This study, our experience, and previously reported case reports of TAF use in the setting of Fanconi syndrome suggests that TAF may have a particularly beneficial role in persons with TDF-induced renal tubular disease.^{10,11} However, one limitation in the translatability of this patient's experience is that he had access to FTC/TAF dosed at 200/10 mg due to concurrent dosing with r/DRV. Whether the FDA-approved dosage of FTC/TAF at 200/25 mg provides similar benefit to persons with tubular nephropathy when combined with a boosted protease inhibitor remains unknown.

The ability of TAF to control HBV replication also supports the use of this antiviral in persons coinfecting with HIV and HBV. In fact, in our patient, the use of TAF was purely to control HBV as he had essentially maintained virologic control of HIV with DRV/r+DTG (due to widespread nucleos(t)ide resistance on GenoSure Archive). TAF at 25 mg has demonstrated similar efficacy to TDF in mono-infected HBV with significantly less bone and renal toxicity up to week 48 of use.¹² One ongoing clinical trial is evaluating the impact of simplification to EVG/COBI/FTC/TAF from TDF-based ART regimens in HIV and HBV coinfecting persons. Similar to our experience, persons that transitioned to EVG/COBI/FTC/TAF maintained HBV viral suppression, improved eGFR, and bone turnover markers.¹³

Conclusion

In clinical trials of persons living with HIV, TAF demonstrates efficacy and a favorable safety and tolerability profile even in persons with mild to moderate kidney dysfunction. Ongoing research will elucidate whether long-term use of TAF in HBV mono and HBV/HIV coinfecting persons will continue to be efficacious, safe, and well tolerated. However, in one patient coinfecting with HIV and HBV, we demonstrate that switching from TDF to TAF did result in improved markers of renal tubulopathy and sustained virologic control of HIV and HBV.

Acknowledgments

The investigator extends her sincere gratitude to the healthcare providers involved in the care of this patient specifically Dr. Eric Shigeno at the Desert AIDS Project, Palm Springs, CA, who assisted in the cocare of this patient,

as well as Drs. Amber Sanchez and Scott Mullaney in the Division of Nephrology and Dr. Irene Vodkin in the Division of Abdominal Transplant and Hepatology at the University of California San Diego, CA. The investigator also is appreciative to Dr. Martin Rhee and colleagues at Gilead Sciences, Foster City, CA, who provided early access to F/TAF through their compassionate care program.

Financial Support

Support for this publication came, in part, by the National Institutes of Health, R01 HD083042 and R01 MH110057. In addition, this publication was supported by the University of California, San Diego Center for AIDS Research (CFAR), an NIH funded program (P30 AI036214), which is supported by the following NIH Institutes and Centers: NIAID, NCI, NIMH, NIDA, NICHD, NHLBI, NIA, NIGMS, and NIDDK. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Author Disclosure Statement

Dr. Karris receives research funding to the institution from Gilead Sciences and has served on an advisory board.

References

1. Gracey DM, Snelling P, McKenzie P, Strasser SI: Tenofovir-associated Fanconi syndrome in patients with chronic hepatitis B mono-infection. *Antivir Ther* 2013;18:945–948.
2. de Vries-Sluijs TE, Reijnders JG, Hansen BE, Zaaier HL, Prins JM, Pas SD, Schutten M, Hoepelman AI, Richter C, Mulder JW, de Man RA, Janssen HL, van der Ende ME: Long-term therapy with tenofovir is effective for patients co-infected with human immunodeficiency virus and hepatitis B virus. *Gastroenterology* 2010;139:1934–1941.
3. Herlitz LC, Mohan S, Stokes MB, Radhakrishnan J, D'Agati VD, Markowitz GS: Tenofovir nephrotoxicity: Acute tubular necrosis with distinctive clinical, pathological, and mitochondrial abnormalities. *Kidney Int* 2010;78:1171–1177.
4. Rodriguez-Novoa S, Labarga P, D'Avolio A, Barreiro P, Albalade M, Vispo E, Solera C, Siccardi M, Bonora S, Di Perri G, Soriano V: Impairment in kidney tubular function in patients receiving tenofovir is associated with higher tenofovir plasma concentrations. *AIDS* 2010;24:1064–1066.
5. Markowitz M, Zolopa A, Squires K, Ruane P, Coakley D, Kearney B, Zhong L, Wulfsohn M, Miller MD, Lee WA: Phase I/II study of the pharmacokinetics, safety and anti-retroviral activity of tenofovir alafenamide, a new prodrug of the HIV reverse transcriptase inhibitor tenofovir, in HIV-infected adults. *J Antimicrob Chemother* 2014;69:1362–1369.
6. Sax PE, Wohl D, Yin MT, Post F, DeJesus E, Saag M, Pozniak A, Thompson M, Podzamczar D, Molina JM, Oka S, Koenig E, Trottier B, Andrade-Villanueva J, Crofoot G, Custodio JM, Plummer A, Zhong L, Cao H, Martin H, Callebaut C, Cheng AK, Fordyce MW, McCallister S: Tenofovir alafenamide versus tenofovir disoproxil fumarate, coformulated with elvitegravir, cobicistat, and emtricitabine, for initial treatment of HIV-1 infection: Two randomised, double-blind, phase 3, non-inferiority trials. *Lancet* 2015;385:2606–2615.
7. Pozniak A, Arribas JR, Gathe J, Gupta SK, Post FA, Bloch M, Avihingsanon A, Crofoot G, Benson P, Lichtenstein K, Ramgopal M, Chetchotisakd P, Custodio JM, Abram ME, Wei X, Cheng A, McCallister S, SenGupta D, Fordyce

- MW: Switching to tenofovir alafenamide, coformulated with elvitegravir, cobicistat, and emtricitabine, in HIV-infected patients with renal impairment: 48-Week results from a single-arm, multicenter, open-label phase 3 study. *J Acquir Immune Defic Syndr* 2016;71:530–537.
8. Bam RA, Yant SR, Cihlar T: Tenofovir alafenamide is not a substrate for renal organic anion transporters (OATs) and does not exhibit OAT-dependent cytotoxicity. *Antivir Ther* 2014;19:687–692.
 9. Post FA, Tebas P, Clarke A, Cotte L, Short WR, Abram ME, Jiang S, Cheng A, Das M, Fordyce M: Brief report: Switching to tenofovir alafenamide, co-formulated with elvitegravir, cobicistat, and emtricitabine, in HIV-infected adults with renal impairment: 96 week results from a single-arm, multicenter, open-label phase 3 study. *J Acquir Immune Defic Syndr* 2017;74:180–184.
 10. Mikula JM, Manion MM, Maldarelli F, Suarez LM, Norman-Wheeler JF, Ober AG, Dewar RL, Kopp JB, Lane HC, Pau AK: Tenofovir alafenamide as part of a salvage regimen in a patient with multi-drug resistant HIV and tenofovir DF-associated renal tubulopathy. *Antivir Ther* 2016;21:553–558.
 11. Garcia M, Le Moal G, Godet C, Beraud G, Chagneau-Derrode C, Roblot F: First case report of renal improvement on tenofovir alafenamide in an HIV/hepatitis B virus-coinfected patient with adefovir-induced Fanconi's syndrome. *AIDS* 2016;30:1487–1488.
 12. Buti M, Seto WK, Chan H, Chuang W, Stepanova T, Hui A, Lim Y, Mehta R, Janssen H, Acharya S, Flaherty J, Massetto B, Cathcart A, Dinh P, Subramanian G, McHutchinson J, Pan C, Brunetto M, Izumi N, Marcellin P: A phase 3 study of tenofovir alafenamide compared with tenofovir disoproxil fumarate in patients with HBeAg-negative, chronic hepatitis B: Week 48 efficacy and safety results. *EASL*, April 22–24, 2016, Barcelona, Spain, 2016.
 13. Gallant J, Brunetta J, Crofoot G, Benson P, Mills A, Brinson C, Cheng A, Garner W, Fordyce M, Das M, McCallister S: Efficacy and safety of switching to simpler single-tablet of elvitegravir/cobicistat/emtricitabine/tenofovir alafenamide (E/C/F/TAF) in HIV-1/hepatitis B coinfecting adults in North America and Japan (NCT02071082) week 48 results. *IAS 2015*, July 19–23, Vancouver, British Columbia, 2015.

Address correspondence to:

Maile Young Karris

Department of Medicine

University of California San Diego

200 West Arbor Drive Mail Code 8208

San Diego, CA 92103-8208

E-mail: m1young@ucsd.edu