Published in final edited form as:

Clin Nutr. 2018 February ; 37(1): 139–143. doi:10.1016/j.clnu.2016.11.003.

Adherence to healthy lifestyle factors and risk of death in men with diabetes mellitus: the Physicians' Health Study

Yash R Patel, MD MPH¹, Taraka V. Gadiraju, MD¹, J. Michael Gaziano, MD MPH^{1,2}, and Luc Djoussé, MD MPH ScD^{1,2}

¹Division of Aging, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA

²Massachusetts Veterans Epidemiology and Research Information Center (MAVERIC) and Geriatric Research, Education, and Clinical Research Center (GRECC), Boston Veterans Affairs Healthcare System, Boston, MA

Abstract

Background & aims—The relationship between healthy lifestyle factors and mortality in people with type 2 diabetes is unclear. The purpose of this study was to examine whether healthy lifestyle factors are associated with mortality in people with type 2 diabetes.

Methods—We prospectively studied 1,163 men with type 2 diabetes from the Physicians' Health Study. Lifestyle factors consisted of currently not smoking, moderate drinking (1–2 drinks/day), vigorous exercise (1+/week), BMI <25 kg/m², and being in the top 2 quintiles of the alternate healthy eating index-2010 (AHEI-2010). Multivariate cox regression models were used to estimate hazard ratios (95% confidence intervals) of mortality.

Results—At baseline, average age was 69 years and mean follow up was 9 years. About 22% of study participants had 1 healthy lifestyle factor, 37% had two, 29% had three, and 12% had four or more healthy lifestyle factors. An inverse relationship was found between the number of lifestyle factors and total mortality. Compared with participants who had 1 healthy lifestyle factor, the risk of death was 42% (95% CI; 19%–58%) lower for those with two healthy lifestyle factors, 41% (95% CI; 18%–58%) lower for those with three, and 44% (95% CI; 12%–64%) lower for those with 4 or more healthy lifestyle factors.

Conclusion—Adherence to modifiable healthy lifestyle factors is associated with a lower risk of death among adult men with type 2 diabetes. Our study emphasizes the importance of educating individuals with diabetes to adhere to healthy lifestyle factors.

Corresponding author: Yash Patel, MD MPH, Division of Aging, Brigham and Women's Hospital, 1620 Tremont St. 3rd floor, Boston MA 02120, dryashpatel21@gmail.com, Tel. (617) 525-7591, Fax. (617) 525-7739.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Disclosures: None by any authors

Keywords

lifestyle; epidemiology; mortality; diabetes mellitus

Introduction

Type 2 Diabetes (DM) is a growing health problem in United States and throughout the world¹. In 2012, 29.1 million Americans had DM, which comprises 9.3% of the US population, with 1.7 million new cases diagnosed in year 2012¹. The total cost of DM care in the United States was \$245 billion in year 2012² and an estimated 1.5 million deaths worldwide were directly related by DM³.

Lifestyle risk modification has been a cornerstone of DM prevention as shown in Diabetes Prevention Study^{4, 5}. This has also been demonstrated in prospective cohort study of Physicians' Health Study, Cardiovascular Health Study, and the Nurses' Health Study, where modifiable lifestyle factors including regular exercise, healthy diet, moderate drinking, normal body weight, and not smoking were jointly associated with a lower residual lifetime risk of DM^{6–8}. In patients with impaired glucose tolerance, lifestyle interventions have been shown to reduce all-cause mortality in Da Qing Diabetes Prevention Study⁹. These healthy lifestyle factors have also been associated with lower risk of all-cause mortality in healthy subjects^{10, 11}.

However, data on the relation of individual and combined lifestyle factors on mortality in people with DM are scarce^{12, 13}. The answer to this question is particularly important, as the impact of lifestyle factors on outcomes might differ between people with DM and the general population free of DM. We sought to test the hypothesis that healthy lifestyle factors are inversely associated with incidence of death in subjects with DM.

Materials and methods

Study design

Data from the Physicians' Health Study (PHS) were used in these analyses. A detailed description of the PHS has been published previously^{14–16}. In brief, PHS was a randomized controlled trial that used a 2×2 factorial design of low-dose aspirin and β -carotene in the primary prevention of cardiovascular disease and cancer. Each participant gave a written informed consent, and the Institutional Review Board at the Brigham and Women's Hospital approved the study protocol. From the 22,071 participants recruited into the PHS, we had 1,163 men with prevalent diabetes at the time of PHS II enrollment (1997). Validation of self-reported diagnosis of DM in this population has been reported⁸.

Ascertainment of lifestyle factors

We focused on modifiable lifestyle factors that have been shown to influence the risk of mortality in DM. These included physical activity, smoking, alcohol consumption, body mass index, and dietary habits. At baseline, each subject provided information on smoking (never, former, and current smoker); alcohol intake (servings per month, week, and day), and

exercise (how often do you exercise vigorously enough to work up sweat? Possible answers included rarely/never, 1–3/month, 1/week, 2–4/week, 5–6/week, and daily). Self-reported baseline weight and height were used to calculate body mass index (weight in kilograms divided by height in meter squared). We used a 128-item food frequency questionnaire (whose validity has been published¹⁷) to assess diet in PHS. For each individual food item, participants were asked to report their average consumption during the past year. Possible answers were never or less than once per month; 1–3/month; 1/week; 2–4/week; 5–6/week; 1/day; 2–3/day; 4–5/day; and 6+/day. We used the alternate healthy eating index (AHEI) AHEI-2010 to assess dietary habits¹⁸. The components of AHEI-2010 diet score include vegetables, fruits, nuts/soy, white to red meat ratio, cereal fiber, trans fat, polyunsaturated to saturated fat ratio, multivitamin use, and alcohol intake.

Definition of lifestyle groups

To investigate the association between healthy lifestyle factors and risk of mortality in DM, each lifestyle factor was dichotomized as follows: normal weight (BMI < 25 kg/m²) vs. overweight/obese (BMI ≥ 25 kg/m²); non-current smoking vs. current smoking; vigorous exercise (≥ 1 day per week) vs. infrequent/no exercise (< 1 day per week). A score of 1 is each assigned to subjects who have BMI < 25 kg/m², are non-current smoking, and are exercising ≥ 1 days per week. AHEI-2010 diet score was divided into quintiles; subjects in the upper 2 quintiles of AHEI-2010 score were considered as adhering to healthy dietary pattern and were given a score of 1. For moderate drinking, a score of 1 was assigned to self-report of up to 1–2 drinks per day as defined by US dietary guidelines and 0 otherwise. Each subject could have a minimum of 0 and a maximum of 5 healthy lifestyle factors. Since only 17 men were in categories with 5 healthy lifestyle factors, we collapsed the upper 2 categories to obtain stable estimates (referred to as 4+ groups). Thus, study participants were categorized according to the number of desirable lifestyle factors (0–1, 2, 3, and 4+).

Ascertainment of deaths

A questionnaire was mailed to participants every 6 months during the first year and annually thereafter to collect data on intervention compliance and new medical diagnoses, including death. Participants who did not return the questionnaires within 5–6 weeks were sent a follow-up questionnaire. Up to 4 questionnaires were sent to nonrespondents. Last, subjects were called if they were still nonrespondents. A letter with a return postcard was also mailed to participants at the 6 month time point between annual mailings, which was to be completed if any major medical problems (e.g., death) occurred. An endpoints committee confirmed death after review of the medical records. The deaths were categorized by type as due to cardiovascular disease, coronary artery disease, stroke, or cancer and based on review of autopsy reports, death certificates, medical records, or family/next of kin report^{14, 16, 19, 20}. Details on endpoint validation in the PHS have been published^{16, 19, 20}.

Statistical analyses

Means and percentages of baseline characteristics of the study participants are presented according to the number of healthy lifestyle factors met by participants. We used Cox proportional hazard models to estimate hazard ratios (HRs) and 95% confidence intervals (CIs). We fitted crude, age-adjusted, and a multivariable models that controlled for age,

history of cancer, coronary heart disease, hypertension, and congestive heart failure. P for linear trend was obtained by treating the number of healthy lifestyle factors met as ordinal variable in the Cox regression models. In secondary analyses, we examined the association of meeting each healthy factor while adjusting for others in the same regression model. Since adherence to healthy diet showed a strong inverse relation with mortality, we examined the relation of AHEI (quintiles) with total mortality by fitting Cox regression as described above. Assumptions of Cox proportional hazard models were checked using product terms of exposure and log (person-time) and were met (all $p > 0.05$).

All analyses were performed using SAS version 9.3 (SAS Institute Inc, Cary, North Carolina), and the alpha level was set at 0.05. All P values were 2-sided.

Results

During a median follow up of 9 years, 248 men died. Table 1 shows baseline characteristics of the participants according to the number of lifestyle factors met. Twenty-two percent of study participants met 1 lifestyle factor, thirty-seven percent met two lifestyle factors, twenty-nine percent met three lifestyle factors, and about twelve percent met four or more lifestyle factors.

In this study, there was an inverse relation between number of healthy lifestyle factors and mortality (Table 2). Compared with participants with <1 healthy lifestyle factor, male diabetic physicians with two, three, and four or more healthy lifestyle factors had a 42 % (95% CI 19%, 58%), 41% (95%CI 18%, 58%), and 44% (95% CI 12%, 64%) lower risk for mortality, respectively ($p=0.005$ for linear trend, Table 2), adjusting for age, history of cancer, coronary heart disease, hypertension, and congestive heart failure. AHEI-2010 was the only statistically significant predictor of mortality among the five healthy lifestyle factors examined (adjusted HR = 0.59; 95% CI 0.44, 0.79, Table 3, comparing subjects in the top 2 quintiles to those in the bottom 3 quintiles of the AHEI-2010; adjusting for age, history of cancer, coronary heart disease, hypertension, and congestive heart failure).

In a secondary analysis, there was an inverse relation of AHEI with mortality (P for trend 0.0003, Table 4), adjusting for age, history of cancer, coronary heart disease, hypertension, and congestive heart failure.

Discussion

In this prospective cohort of male physicians with DM, we found an inverse relation between number of healthy lifestyle factors met and all-cause mortality. Studies focusing on individual lifestyle factors and mortality exist, including inverse association with Mediterranean diet²¹, physical activity²², moderate alcohol consumption²³, healthy eating (fruits, vegetables, and legumes²⁴, whole-grain, cereal fiber, bran, and germ intake²⁵), and positive association with smoking cigarettes²⁶. However, only limited studies have focused on combination of lifestyle factors in adults with DM. Nothlings et al. conducted a prospective cohort study of 1263 German adults with DM and followed them for an average of 7.8 years to study healthy lifestyle behaviors - no smoking, BMI < 30 kg/m², physical activity for ≥ 3.5 h/week, dietary habits, and moderate alcohol consumption, on mortality

in people with DM¹³. Compared with individuals with no favorable factors, the risk of death was 34% lower for those with one favorable risk factor, 49% lower for those with two risk factors, and 63% lower for subjects with three or more risk factors. Similarly, Lin et al. studied lifestyle-related factors (smoking, alcohol use, exercise, and carbohydrate intake) on all-cause and cause-specific mortality in 5686 patients with DM: after a median follow-up of 4.0 years, higher all-cause mortality was observed with smoking (HR 1.31, 95% CI 1.01–1.71), alcohol drinking (HR 1.43, 95% CI 1.03–2.00), and not exercising (HR 1.63, 95% CI 1.34–1.99) while excess carbohydrate intake (>65%) was not associated with all-cause mortality (HR 1.39, 95% CI 0.92–2.09)¹². Our study results are further supported by recently published MOLI-SANI study²¹ where traditional Mediterranean diet consistent with moderate alcohol drinking, high intake of cereals, vegetables, and reduced consumption of dairy and meat products, was associated with lower risk of both total and cardiovascular mortality in individuals with DM. Furthermore, healthy lifestyle factors were also associated with a lower risk of total mortality in people with impaired glucose tolerance in the Finnish Diabetes Prevention Study and DaQing Diabetes Prevention Study^{9, 27}.

In our study, healthy diet appeared to have the strongest relation with mortality among individuals with DM. This could be explained by higher AHEI score being associated with lower risk factors of cardiovascular disease in DM including waist circumference and lower density lipoprotein cholesterol²⁸. From a recent meta-analysis of 1,020,642 subjects, Schwingschackl et al. found that high AHEI score is associated with a significant lower risk of all-cause mortality, cardiovascular disease, and type DM²⁹.

Strengths and limitations of this study

The strengths of our study include the lengthy follow-up period, the standardized methods for data collection, comprehensive assessment of dietary habits, and the robustness of the findings in multivariate analyses. Our study has several limitations. First, given the PHS enrolled mostly Caucasian male physicians, the results from the study limit generalizability to the general population. Second, unmeasured confounding factors might partially or completely explain our findings. Third, diagnosis of DM was done by self-reporting, and despite validation of the diagnosis of DM by 2 independent physicians through review of medical records, we cannot exclude the possibility of DM misclassification. Fourth, our sample size was relatively small to examine the effect of other individual lifestyle factors including smoking, moderate exercise, and vigorous exercise, on mortality.

Conclusion

In this prospective study, we have shown that adherence to healthy lifestyle factors is inversely related to death among men with DM. Confirmation of these results in other populations is needed.

Acknowledgments

We are indebted to the participants in the PHS for their outstanding commitment and cooperation and to the entire PHS staff for their expert and unfailing assistance.

Sources of Funding: The Physicians' Health Study is supported by grants CA-34944, CA-40360, and CA-097193 from the National Cancer Institute and grants R21 HL088081, HL-26490, and HL-34595 from the National Heart, Lung, and Blood Institute, Bethesda, MD.

References

1. Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes research and clinical practice*. 2014; 103(2):137–149. e-pub ahead of print 2014/03/19. DOI: 10.1016/j.diabres.2013.11.002 [PubMed: 24630390]
2. The cost of diabetes. American Diabetes Association. (Accessed February 19, 2016, at <http://www.diabetes.org/advocacy/news-events/cost-of-diabetes.html?referrer=https://www.google.com/>). In
3. Diabetes. World Health Organization; (Accessed February 19, 2016, at <http://www.who.int/mediacentre/factsheets/fs312/en/>). In
4. The Diabetes Prevention Program (DPP): description of lifestyle intervention. *Diabetes care*. 2002; 25(12):2165–2171. e-pub ahead of print 2002/11/28. [PubMed: 12453955]
5. Knowler WC, Barrett-Connor E, Fowler SE, Hamman RF, Lachin JM, Walker EA, et al. Reduction in the incidence of type 2 diabetes with lifestyle intervention or metformin. *The New England journal of medicine*. 2002; 346(6):393–403. e-pub ahead of print 2002/02/08. DOI: 10.1056/NEJMoa012512 [PubMed: 11832527]
6. Hu FB, Manson JE, Stampfer MJ, Colditz G, Liu S, Solomon CG, et al. Diet, lifestyle, and the risk of type 2 diabetes mellitus in women. *The New England journal of medicine*. 2001; 345(11):790–797. e-pub ahead of print 2001/09/15. DOI: 10.1056/NEJMoa010492 [PubMed: 11556298]
7. Mozaffarian D, Kamineni A, Carnethon M, Djousse L, Mukamal KJ, Siscovick D. Lifestyle risk factors and new-onset diabetes mellitus in older adults: the cardiovascular health study. *Archives of internal medicine*. 2009; 169(8):798–807. e-pub ahead of print 2009/04/29. DOI: 10.1001/archinternmed.2009.21 [PubMed: 19398692]
8. Djousse L, Driver JA, Gaziano JM, Buring JE, Lee IM. Association between modifiable lifestyle factors and residual lifetime risk of diabetes. *Nutrition, metabolism, and cardiovascular diseases : NMCD*. 2013; 23(1):17–22. e-pub ahead of print 2011/10/11. DOI: 10.1016/j.numecd.2011.08.002
9. Li G, Zhang P, Wang J, An Y, Gong Q, Gregg EW, et al. Cardiovascular mortality, all-cause mortality, and diabetes incidence after lifestyle intervention for people with impaired glucose tolerance in the Da Qing Diabetes Prevention Study: a 23-year follow-up study. *The lancet Diabetes & endocrinology*. 2014; 2(6):474–480. e-pub ahead of print 2014/04/16. DOI: 10.1016/S2213-8587(14)70057-9 [PubMed: 24731674]
10. Prinelli F, Yannakoulia M, Anastasiou CA, Adorni F, Di Santo SG, Musicco M, et al. Mediterranean diet and other lifestyle factors in relation to 20-year all-cause mortality: a cohort study in an Italian population. *The British journal of nutrition*. 2015; 113(6):1003–1011. e-pub ahead of print 2015/03/10. DOI: 10.1017/S0007114515000318 [PubMed: 25746109]
11. Petersen KE, Johnsen NF, Olsen A, Albieri V, Olsen LK, Dragsted LO, et al. The combined impact of adherence to five lifestyle factors on all-cause, cancer and cardiovascular mortality: a prospective cohort study among Danish men and women. *The British journal of nutrition*. 2015; 113(5):849–858. e-pub ahead of print 2015/02/19. DOI: 10.1017/S0007114515000070 [PubMed: 25690300]
12. Lin CC, Li CI, Liu CS, Lin WY, Fuh MM, Yang SY, et al. Impact of lifestyle-related factors on all-cause and cause-specific mortality in patients with type 2 diabetes: the Taichung Diabetes Study. *Diabetes care*. 2012; 35(1):105–112. e-pub ahead of print 2011/11/30. DOI: 10.2337/dc11-0930 [PubMed: 22124717]
13. Nothlings U, Ford ES, Kroger J, Boeing H. Lifestyle factors and mortality among adults with diabetes: findings from the European Prospective Investigation into Cancer and Nutrition-Potsdam study*. *Journal of diabetes*. 2010; 2(2):112–117. e-pub ahead of print 2010/10/07. DOI: 10.1111/j.1753-0407.2010.00069.x [PubMed: 20923493]

14. Djousse L, Gaziano JM. Egg consumption in relation to cardiovascular disease and mortality: the Physicians' Health Study. *The American journal of clinical nutrition*. 2008; 87(4):964–969. e-pub ahead of print 2008/04/11. [PubMed: 18400720]
15. Christen WG, Gaziano JM, Hennekens CH. Design of Physicians' Health Study II—a randomized trial of beta-carotene, vitamins E and C, and multivitamins, in prevention of cancer, cardiovascular disease, and eye disease, and review of results of completed trials. *Annals of epidemiology*. 2000; 10(2):125–134. e-pub ahead of print 2000/02/26. [PubMed: 10691066]
16. Muntwyler J, Hennekens CH, Manson JE, Buring JE, Gaziano JM. Vitamin supplement use in a low-risk population of US male physicians and subsequent cardiovascular mortality. *Archives of internal medicine*. 2002; 162(13):1472–1476. e-pub ahead of print 2002/07/02. [PubMed: 12090883]
17. Willett WC, Sampson L, Stampfer MJ, Rosner B, Bain C, Witschi J, et al. Reproducibility and validity of a semiquantitative food frequency questionnaire. *American journal of epidemiology*. 1985; 122(1):51–65. e-pub ahead of print 1985/07/01. [PubMed: 4014201]
18. Chiuve SE, Fung TT, Rimm EB, Hu FB, McCullough ML, Wang M, et al. Alternative dietary indices both strongly predict risk of chronic disease. *The Journal of nutrition*. 2012; 142(6):1009–1018. e-pub ahead of print 2012/04/20. DOI: 10.3945/jn.111.157222 [PubMed: 22513989]
19. Albert CM, Campos H, Stampfer MJ, Ridker PM, Manson JE, Willett WC, et al. Blood levels of long-chain n-3 fatty acids and the risk of sudden death. *The New England journal of medicine*. 2002; 346(15):1113–1118. e-pub ahead of print 2002/04/12. DOI: 10.1056/NEJMoa012918 [PubMed: 11948270]
20. Bowman TS, Gaziano JM, Kase CS, Sesso HD, Kurth T. Blood pressure measures and risk of total, ischemic, and hemorrhagic stroke in men. *Neurology*. 2006; 67(5):820–823. e-pub ahead of print 2006/09/13. DOI: 10.1212/01.wnl.0000233981.26176.e1 [PubMed: 16966544]
21. Bonaccio M, Di Castelnuovo A, Costanzo S, Persichillo M, De Curtis A, Donati MB, et al. Adherence to the traditional Mediterranean diet and mortality in subjects with diabetes. Prospective results from the MOLI-SANI study. *European journal of preventive cardiology*. 2016; 23(4):400–407. e-pub ahead of print 2015/02/05. DOI: 10.1177/2047487315569409 [PubMed: 25648935]
22. Trichopoulou A, Psaltopoulou T, Orfanos P, Trichopoulos D. Diet and physical activity in relation to overall mortality amongst adult diabetics in a general population cohort. *Journal of internal medicine*. 2006; 259(6):583–591. e-pub ahead of print 2006/05/18. DOI: 10.1111/j.1365-2796.2006.01638.x [PubMed: 16704559]
23. Koppes LL, Dekker JM, Hendriks HF, Bouter LM, Heine RJ. Meta-analysis of the relationship between alcohol consumption and coronary heart disease and mortality in type 2 diabetic patients. *Diabetologia*. 2006; 49(4):648–652. e-pub ahead of print 2006/02/08. DOI: 10.1007/s00125-005-0127-x [PubMed: 16463045]
24. Nothlings U, Schulze MB, Weikert C, Boeing H, van der Schouw YT, Bamia C, et al. Intake of vegetables, legumes, and fruit, and risk for all-cause, cardiovascular, and cancer mortality in a European diabetic population. *The Journal of nutrition*. 2008; 138(4):775–781. e-pub ahead of print 2008/03/22. [PubMed: 18356334]
25. He M, van Dam RM, Rimm E, Hu FB, Qi L. Whole-grain, cereal fiber, bran, and germ intake and the risks of all-cause and cardiovascular disease-specific mortality among women with type 2 diabetes mellitus. *Circulation*. 2010; 121(20):2162–2168. e-pub ahead of print 2010/05/12. DOI: 10.1161/CIRCULATIONAHA.109.907360 [PubMed: 20458012]
26. Al-Delaimy WK, Willett WC, Manson JE, Speizer FE, Hu FB. Smoking and mortality among women with type 2 diabetes: The Nurses' Health Study cohort. *Diabetes care*. 2001; 24(12):2043–2048. e-pub ahead of print 2001/11/28. [PubMed: 11723080]
27. Uusitupa M, Peltonen M, Lindstrom J, Aunola S, Ilanne-Parikka P, Keinänen-Kiukaanniemi S, et al. Ten-year mortality and cardiovascular morbidity in the Finnish Diabetes Prevention Study—secondary analysis of the randomized trial. *PloS one*. 2009; 4(5):e5656. e-pub ahead of print 2009/05/30. doi: 10.1371/journal.pone.0005656 [PubMed: 19479072]
28. Wu PY, Huang CL, Lei WS, Yang SH. Alternative health eating index and the Dietary Guidelines from American Diabetes Association both may reduce the risk of cardiovascular disease in type 2

- diabetes patients. *Journal of human nutrition and dietetics : the official journal of the British Dietetic Association*. 2015; e-pub ahead of print 2015/05/27. doi: 10.1111/jhn.12317
29. Schwingshackl L, Hoffmann G. Diet quality as assessed by the Healthy Eating Index, the Alternate Healthy Eating Index, the Dietary Approaches to Stop Hypertension score, and health outcomes: a systematic review and meta-analysis of cohort studies. *Journal of the Academy of Nutrition and Dietetics*. 2015; 115(5):780–800. e785. e-pub ahead of print 2015/02/15. DOI: 10.1016/j.jand.2014.12.009 [PubMed: 25680825]

Table 1

Baseline characteristics of the 1,163 male physicians with prevalent DM according to the number of lifestyle factors met*

	Number of lifestyle factors met				P value
	1 (n=258)	2 (n=421)	3 (n=341)	4+ (n=143)	
Age (years)	67.5 ± 7.9	68.8 ± 8.3	70.1 ± 8.2	69.6 ± 8.1	0.001
BMI (kg/m ²)	29.7 ± 4.1	27.7 ± 3.8	26.3 ± 4.0	23.8 ± 2.7	<0.0001
Energy intake (Kcal)	1591 ± 489	1641 ± 522	1711 ± 521	1907 ± 630	<0.0001
AHEI-2010 score	38.5 ± 8.4	44.6 ± 10.2	50.0 ± 11.1	57.0 ± 7.9	<0.0001
Fruits & vegetables (serv/d)	4.1 ± 2.0	4.7 ± 2.2	5.6 ± 2.9	6.7 ± 3.2	<0.0001
Whole grain (serv/d)	1.2 ± 1.3	1.6 ± 1.5	1.9 ± 1.8	2.9 ± 2.6	<0.0001
Atrial fibrillation (%)	12.0	12.4	12.0	15.4	0.75
Cancer (%)	11.2	10.7	12.9	10.5	0.78
Coronary heart disease (%)	18.6	27.3	30.2	28.0	0.01
Congestive heart failure (%)	7.0	5.7	8.8	5.6	0.34
Hypertension (%)	72.1	68.7	71.3	64.3	0.62

* Lifestyle factors includes not smoking, moderate drinking (1–2 drinks/day), vigorous exercise (1+/week), BMI <25 kg/m², and being in the top 2 quintiles of the alternate healthy eating index-2010.

Data are presented as mean±SD or %.

Abbreviations: AHEI-2010, alternate healthy index – 2010; BMI, body mass index, DM; type 2 diabetes mellitus.

Table 2

HR (95%CI) for death by number of lifestyle factors met in 1,163 male physicians with DM

No. of lifestyle factors	% of deaths	Hazard ratios (95%CI)		
		Crude	Age - adjusted model	Multivariable model ^a
<1	29	REF	REF	REF
2	33	0.70 (0.51–0.97)	0.58 (0.42–0.80)	0.58 (0.42–0.81)
3	28	0.79 (0.57–1.11)	0.59 (0.43–0.83)	0.59 (0.42–0.82)
4+	10	0.72 (0.46–1.13)	0.56 (0.35–0.87)	0.56 (0.36–0.88)
P for linear trend		0.17	0.005	0.005

^aHazard ratio (HR) with 95% confidence interval (CI) from a model including age, history of cancer, coronary heart disease, hypertension, and congestive heart failure.

Abbreviations: DM; type 2 diabetes mellitus.

Table 3

HR (95%CI) for death by individual lifestyle factors met in 1,163 male physicians with DM

Individual lifestyle factors	Hazard ratios (95%CI)		
	Crude	Age-adjusted model	Multivariable model ^a
Not smoking (n=1107)	1.19 (0.65–2.18)	0.86 (0.47–1.59)	0.91 (0.49–1.68)
Moderate drinking (n=202)	1.08 (0.79–1.50)	0.99 (0.72–1.37)	0.96 (0.70–1.33)
Vigorous exercise (n=601)	0.90 (0.70–1.15)	0.86 (0.67–1.11)	0.84 (0.65–1.08)
BMI<25 kg/m² (n=347)	1.17 (0.90–1.54)	0.95 (0.72–1.24)	1.01 (0.77–1.32)
AHEI-2010 (n=435)	0.64 (0.48–0.85)	0.59 (0.45–0.79)	0.59 (0.44–0.79)

^aHazard ratio (HR) with 95% confidence interval (CI) from a model including age, history of cancer, coronary heart disease, hypertension, and congestive heart failure. HRs are mutually adjusted across lifestyle factors.

Abbreviations: AHEI-2010, alternate healthy index – 2010; BMI, body mass index, DM; type 2 diabetes mellitus.

Table 4

HR (95%CI) for death by quintiles of AHEI-2010 in 1,163 male physicians with DM

Quintiles of AHEI-2010	No. of deaths	Hazard ratios (95%CI)		
		Crude	Age - adjusted model	Multivariable model ^a
1	61	REF	REF	REF
2	58	1.05 (0.73–1.50)	0.97 (0.68–1.39)	0.99 (0.69–1.43)
3	65	0.98 (0.69–1.38)	0.88 (0.62–1.24)	0.86 (0.61–1.23)
4	38	0.74 (0.49–1.11)	0.58 (0.39–0.88)	0.61 (0.40–0.92)
5	26	0.54 (0.34–0.86)	0.52 (0.33–0.82)	0.50 (0.31–0.80)
P linear trend		0.0033	0.0003	0.0003

^aHazard ratio (HR) with 95% confidence interval (CI) from a model including age, individual lifestyle factors (body mass index, smoking, alcohol, and exercise), and history of cancer, coronary heart disease, hypertension, and congestive heart failure.

Abbreviations: AHEI-2010, alternate healthy index – 2010, DM; type 2 diabetes mellitus.