Published in final edited form as:

Prev Med. 2017 July ; 100: 229–234. doi:10.1016/j.ypmed.2017.04.026.

The Stability of Children's Weight Status Over Time, and the Role of Television, Physical Activity, and Diet in Elementary School

Sandra L. Jackson, PhD^a and Solveig A. Cunningham, PhD^b

^aNutrition and Health Sciences Program, Graduate Division of Biological and Biomedical Sciences, Emory University, 1518 Clifton Road; Atlanta, GA 30322, USA, sandra.jackson@outlook.com ^bHubert Department of Global Health, Rollins School of Public Health, Emory University, 1518 Clifton Road; Atlanta, GA 30322, USA, sargese@emory.edu

Abstract

Weight-related behaviors such as sedentary activity, physical activity, and diet have been the focus of efforts to prevent and reduce the occurrence of obesity and overweight in children, but few longitudinal studies have examined the effects of weight-related behaviors on changes in weight status over time in children. This study examines the effects of weight-related behaviors on subsequent changes in weight during childhood. We used the Early Childhood Longitudinal Study Kindergarten Cohort (ECLS-K), a nationally representative prospective cohort of children in the United States. Data, including anthropometric measures, were collected six times across 1998–2007 (analytic sample =4,938). We employed an autoregressive cross-lagged model in a structural equation model framework to assess the effects of behavioral factors -intake of fruit, vegetables, fast food and soda, television viewing, and physical activity - on weight stability over time. BMI z-scores were highly stable throughout childhood: the standardized parameter estimates of BMI z-scores on subsequent period BMI z-scores ranged from 0.79 to 0.86. BMI z-was least stable between Kindergarten and 1st grade but became highly stable between 3rd and 5th grades. After accounting for prior weight, behavioral factors had little effect on subsequent weight. The most important behavioral factor was TV viewing in the 1st and 3rd grades; an additional hour of daily TV viewing was associated with 0.04 higher BMI z-scores. It is important to prevent excessive weight gain early in childhood, as weight patterns are long-lasting; the most important behavioral factor may be limiting children's screen time.

Corresponding author: Solveig A. Cunningham; Hubert Department of Global Health, Emory University, 1518 Clifton Road; Atlanta, GA 30322, USA; sargese@emory.edu; tel: 404-727-6486; fax: 404-727-4590.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest

The authors declare they have no conflict of interest.

Keywords

Child health; obesity; overweight; health behaviors; trajectories; United States

Introduction

Childhood obesity has become a major public health concern, with recent estimates in the United States showing 18% of children between the ages of 6 and 11 were obese (Ogden et al., 2014). Obesity in children is associated with a number of serious health problems, including asthma and sleep-disordered breathing, menstrual abnormalities, early onset of type 2 diabetes, and metabolic disorders (Institute of Medicine, 2005), as well as psychological risks such as depression and anxiety (Storch et al., 2007).

Evidence suggests that weight trajectories are established early in childhood (Franzese et al., 1998; Hesketh et al., 2004) and obesity in childhood is difficult to reverse (Kamath et al., 2008) and often persists into adulthood (Singh et al., 2008). For example, body mass index (BMI) at school entry is a strong predictor of obesity in adolescence (Nielsen et al., 2007). Studies beginning even earlier in life have shown that infants with high BMI are at higher risk of obesity later in childhood and adulthood compared with other children (Baird et al., 2005; Vogels et al., 2006). Many mechanisms have been suggested to explain the early development of weight trajectory, such as genetic and epigenetic factors, breastfeeding and early feeding practices, and family and environmental factors (Carnell et al., 2012; Skelton et al., 2011).

Although prior weight is a strong predictor of subsequent weight, not all children who become overweight or obese remain at the upper end of weight distribution throughout childhood (Huang et al., 2011). For example, in one study, about 13% of children who were overweight at age 7 became normal weight by age 11 (Wright et al., 2010). Weight-related behaviors, such as screen time, physical activity, and dietary choices, may play a role in the development or reversal of obesity (Carson et al., 2016; Danner, 2008; Janssen and Leblanc, 2010; Malik et al., 2006; Thibault et al., 2010). To some degree, behavior may overcome underlying predisposition to obesity, including genetic vulnerability. For example, evidence suggests that physical activity (Ahmad et al., 2013) and dietary restraint (Vogels et al., 2006) can attenuate genetic risk of obesity. Numerous interventions have been developed to target changing these weight-related behaviors among school-aged children, but, so far, their effectiveness has been limited (Kamath et al., 2008). The persistence of obesity suggests the importance of investigating early causes and possible avenues of early intervention.

Given the emphasis placed on behavior change in childhood obesity prevention interventions and policy recommendations (Waters et al., 2011; World Health Organization, 2016), the aim of this study was to explore the stability of weight and the effects of weight-related behaviors in determining children's weight status in elementary and middle school. We also aimed to investigate whether – contrary to current thinking – associations between presumed weight-related behaviors and weight status may be partly driven by reverse causality: i.e., a cross-sectional association between weight status and television viewing may be due in part to children who are already obese choosing to watch more television, rather than the

accepted wisdom that watching more television leads to obesity (Carson et al., 2016). To examine such relationships, we used data from the Early Childhood Longitudinal Study (ECLS-K), a large and nationally representative longitudinal survey of children followed from elementary to middle school. We estimated an autoregressive cross-lagged panel model in a structural equation-modeling (SEM) framework that allowed us to examine changes in children's BMI z-score in relation to weight-related behaviors across time (Biesanz, 2012). By controlling for the previous period's measures, we aimed to assess whether behaviors believed to be related to weight, such as television viewing, physical activity, and dietary intake, predict changes in weight, or whether weight itself predicts changes in these behaviors.

Data and Methods

The Early Childhood Longitudinal Study, Kindergarten Class of 1998–99 (ECLS-K), developed by the National Center for Education Statistics, followed a nationally representative cohort of children from kindergarten into eighth grade. During Kindergarten and grades 1, 3, 5, and 8, the survey included one-on-one direct child assessments, interviews with parents, teachers, and principals, and, in grades 5 and 8, interviews with children (Tourangeau et al., 2009). The ECLS-K is the only national representative longitudinal survey with objectively measured height and weight in this age group; these measures were collected twice per wave by trained assessors (U.S. Department of Education, 2004), thereby avoiding the bias of self-reported weight and height (Bogaert et al., 2003).

Variables

Height and weight were collected at each round in the ECLS-K. Using the 2000 CDC Growth Charts, we calculated standardized BMI using sex and age in months (Vidmar et al., 2013). We used the standard cutoffs of 95th and 85th percentiles to identify obese, overweight, and normal weight status.

We calculated hours of television viewing per day based on numerous indicators across waves of data collection. In the Kindergarten wave, parents were asked how many hours per day the child watched television during weekdays and during the weekend. In 1st, 3rd, and 5th grades, parents were asked about how much television the child watched per day during particular segments of the day during weekdays, and separately on Saturdays and Sundays. In 8th grade, children were asked how many hours of television they watched per day during weekdays and on weekends. During this wave, 58 children reported implausible viewing amounts; we top-coded the number of hours of television per day for these individuals at 18 hours. For Kindergarten, 1st grade, 3rd grade, and 5th grade, there were 2, 3, 3, and 1 individuals with top-coded values, respectively.

In Kindergarten, grade 3, and grade 5, parents were asked: "In a typical week, on how many days does (CHILD) get exercise that causes rapid breathing, perspiration, and a rapid heartbeat for 20 continuous minutes or more?" In 8th grade, a similar question was asked of the children instead of the parents: "On how many of the past 7 days did you exercise or participate in physical activity for at least 20 minutes that made you sweat and breathe hard,

such as basketball, soccer, running, swimming laps, fast bicycling, fast dancing, or similar aerobic exercise?"

In the 5th and 8th grades, children were asked how many times in the past week they consumed certain foods; we grouped foods together as healthy choices (fruits, green salads, carrots, and other vegetables) and as less healthy choices (fast food, soda and juice drinks). For green salad, for example, children were asked: "During the past 7 days, how many times did you eat green salad?" The responses were coded as: "I did not eat green salad during the past 7 days" (0), "1 to 3 times during the past 7 days" (2/7), "4 to 6 times during the past 7 days" (5/7), "1 time per day" (1), "2 times per day" (2), "3 times per day" (3), "4 or more times per day" (4). Although children were also asked about purchases of snacks, sweets, and drinks at school, due to the large proportion of students who were unable to purchase these items at school (ranging from 61% in grade 5 to 23% in grade 8), we did not include these indicators.

When estimating the full model, we also included relevant covariates that may be associated with weight or with weight-related behaviors, including gender (male vs. female), race/ethnicity (non-Hispanic white, non-Hispanic black, Hispanic, and other), and socioeconomic status. Socioeconomic status at each grade was available as a composite continuous measure provided by the ECLS-K based on parent education, occupation, and household income at the time of interview.

Analysis

We used ECLS-K survey weights to maintain the nationally representative nature of the data across all waves of data collection, and produced survey-adjusted results. There were 4,938 children with valid longitudinal sample weights and non-missing data for all variables at all waves.

After descriptive analyses, we constructed a cross-lagged longitudinal model to investigate the stability of children's weight over time and (1) whether weight-related behaviors were associated with subsequent BMI after accounting for the previous period's BMI or (2) whether BMI was associated with subsequent weight-related behaviors. Our model, using a structural equation modeling framework, featured observed variables for physical activity, television and BMI z-score and latent variables for dietary quality based on observed measures for fruit, vegetables, soda, and fast food consumption. At each time point, BMI and weight-related behaviors were allowed to co-vary. We used robust maximum likelihood, also called pseudo-maximum likelihood (PML), to account for the complex survey design in the ECLS-K (Oberski, 2014). PML scales the chi-square model statistic and derived model fit measures and provides robust standard errors. We evaluated overall model fit, which was acceptable according to the root mean squared error of approximation (RMSEA) and the comparative fit index (CFI) (West et al., 2012) (Browne et al., 1993). The measurement models for healthy (fruit and vegetables) and unhealthy (sweetened beverages and fast food) dietary behavior had acceptable fits to the data. We used Stata 13.1 (StataCorp, 2013) to produce descriptive statistics and R (3.1.1) (Team, 2014) and R packages lavaan (0.5–17) (Rosseel), survey (3.30-3) (Lumley, 2014), and lavaan.survey (1.1) (Oberski) to produce the survey-adjusted analytic models.

Results

In this cohort representative of children who entered Kindergarten in 1998 or first grade in 1999, 49% were boys; 60% non-Hispanic white, 14% non-Hispanic black, 19% Hispanic, and 7% other race/ethnicity (Table 1). The percent of children at or above the cut-offs for obesity ranged from a low of 12.5% in Kindergarten to a high of 21.7% in 5th grade.

In fifth grade, the majority of children had soda (85%) and fast food (72%) at least once a week (Table 2). Approximately 62% of children reported eating less than one serving of fruit each day, 91% of children reported less than one serving of green salad per day, 88% of children reported less than one serving of carrots per day, and 74% reported less than one serving of other vegetables per day. These proportions remained unchanged in eighth grade.

Parents reported that their children watched an average of two hours of television daily in Kindergarten, 1st grade, 3rd grade, and 5th grade; children reported watching over 3 hours of television daily in 8th grade.

Children engaged in at least 20 minutes of vigorous physical activity on approximately 4 days per week in Kindergarten, 3rd grade, and 5th grade, according to their parents; when children were asked for themselves in 8th grade, they reported engaging in vigorous physical activity nearly 5 days per week.

BMI was highly stable throughout childhood (Figures 1a and 1b) after accounting for weight-related behaviors, gender, race, and socioeconomic status. The standardized parameter estimates for autoregressive relationships were all significant. In particular, the standardized parameter estimates for BMI are substantial, ranging from 0.79 to 0.86 ($p < 0.001$). These values indicate that the relative rank order of BMI z-scores stayed largely the same across time, even if individuals were increasing in weight. BMI z-score was most stable between 3rd and 5th grades, and least stable between Kindergarten and 1st grade. The overall variance explained by the model for 1st, 3rd, 5th and 8th grade BMI z-score was 63.5%, 65.2%, 76.1% and 74.5%, respectively.

After accounting for previous BMI z-score, behavioral factors including food and drink consumption, TV viewing, and physical activity had little impact on a child's BMI z-score between Kindergarten and 8th grade. For example, of the weight-related behaviors examined, only television viewing was significantly associated with subsequent BMI z-score. In 1st and 3rd grades, hours of daily television viewing was a significant predictor of BMI z-score in the 3rd (0.04, $p = 0.020$) and 5th (0.04, $p = 0.016$) grades, respectively. This indicates that a 1-hour increase in daily TV viewing was associated with a 0.04 increase in BMI z-score in the subsequent grade, while a change in BMI z-score of >0.5 is considered clinically meaningful (Reinehr and Andler, 2004). Across all ages, physical activity and dietary choices were not significantly associated with subsequent BMI z-score.

In addition to testing whether weight-related behaviors influenced subsequent BMI z-score, we also tested whether BMI z-score influenced subsequent weight-related behaviors, to examine the potential role of reverse causality. Although the effects were small, BMI z-score appeared to impact subsequent physical activity and television viewing. BMI z-score in 1st

grade was a significant predictor of physical activity in the 3rd grade (-0.05 , $p = 0.026$), with higher BMI z-score associated with lower subsequent physical activity. BMI z-score in 3rd grade was a significant predictor of TV in 5th grade (0.06 , $p = 0.001$), with greater BMI z-score associated with subsequently greater television viewing.

Discussion

This study found that children's BMI z-scores are highly stable throughout elementary and middle school. It also found that health behaviors that are central to recommendations for preventing and reversing childhood obesity - television viewing, physical activity, and dietary intake - were largely not significant predictors of subsequent BMI z-score, after controlling for prior weight status. The one exception is screentime -hours of television viewing in the first and third grades were positively associated with slightly higher subsequent BMI z-scores. Simultaneously, we found that higher BMI z-score was associated with subsequently more television viewing and less exercise, indicating that overweight in children may negatively affect behavior.

Prior BMI was a much stronger predictor of subsequent weight than any weight-related behaviors. This is consistent with other studies that have cast doubt on the impact of lifestyle factors on BMI trajectory (Twisk et al., 1998), as well as prior research indicating the importance of early BMI in the establishment of obesity in childhood (Franzese et al., 1998; Hesketh et al., 2004). Influences on weight that may be set in place prior to school entry include factors such as genetic traits (Skelton et al., 2011) predispositions such as cognitive restraint, disinhibition, hunger, food cue responsiveness, eating rate, satiety responsiveness, and preference for fatty food (Carnell et al., 2012), environmental exposures such as gut microbiota (Skelton et al., 2011), endocrine disruptors (Karoutsou and Polymeris, 2012), or other chemicals and pollutants (La Merrill and Birnbaum, 2011), intrauterine and neonatal environments including maternal under-nutrition (Ravelli et al., 1976), maternal over-nutrition (Skelton et al., 2011), diabetes (Skelton et al., 2011), breastfeeding (Skelton et al., 2011), and parents' feeding styles (Carnell et al., 2012).

While early-life factors may be of primary importance in the establishment and maintenance of weight trajectories, some of children's behaviors may affect changes in weight. Particularly, TV viewing was associated with slight subsequent BMI increases. This may be due to an obesogenic effect of sedentary time (i.e., fewer calories are expended while seated compared to other activities that may involve standing, walking, or playing); it may be also be linked to the increased caloric intake that occurs during screen time (Rey-Lopez et al., 2008), prompted by lack of attention to portion sizes and advertising cues to consume high-calorie foods. Although the association between television viewing and BMI was too small to be considered clinically meaningful (Reinehr and Andler, 2004), it may be that this and other behaviors have a greater impact on weight change among children who are already at-risk for obesity than among leaner children (Mitchell et al., 2013). For example, children with low social competence are more likely to become overweight or obese (Jackson and Cunningham, 2015), which may be driven by degree of preference for sedentary, solitary activity such as television viewing. Exploring effects of potential weight-related behaviors among population subgroups may be a fruitful direction for future research. In addition,

while the present study was only able to examine television viewing, children today are exposed to a greater variety of electronic media, and the combined effects of television viewing and screen time on other devices such as computers, tablets, and cell phones could be explored.

We also noted that obesity may drive subsequent behavior changes, such that children with higher BMI z-score were more likely to subsequently watch more television and engage in less physical activity. We could conceive of a cycle linking weight and behaviors across childhood in ways that reinforce obesity, in which children's behaviors such as television viewing affect their weight; as they become heavier, physical activity becomes more tiresome and uncomfortable and therefore less frequent (Tucker et al., 2013), leading to further weight gain. In addition, obese children are often subject to weight-related teasing and exclusion, which has been shown to discourage physical activity and encourage sedentary time (Barkley et al., 2012; Storch et al., 2007). This pattern is consistent with a longitudinal study conducted among Australian children, which demonstrated a bi-directional relationship between television and BMI, such that children who watched more television were more likely to gain weight, and heavier children were more likely to watch TV (Fuller-Tyszkiewicz et al., 2012).

This study has limitations. Some variables, such as television viewing and exercise, were parent-reported in grades K through 5, and reported by children in grade 8. The increases in television viewing and exercise reported from fifth to eighth grade, and the simultaneous decreases in parameter estimates, could indicate measurement error due the differences between parent-reported vs. self-reported activities. Such measurement error would only introduce bias if it occurred differently for children who were heavier compared with lighter. A second consideration is that a three-year time lag, such as between fifth and eighth grades, is a long time in the context of child growth and development. Still, during elementary school, children are already developing their sense of identity and establishing self-perceptions (as, for example, a person who is good at sports), and many related perceptions and behaviors are likely to carry through the transition to middle school (Huston and Ripke, 2006). Some of the food measures deviated from the assumption of normality, which may bias parameter and standard error estimates downwards; robust standard errors were used to partially address this issue (Finney and DiStefano, 2006). Another limitation is that measures of food consumption were not available prior to grades five and eight, so any effect of dietary habits in earlier grades was not captured. Future studies would ideally incorporate validated measures of food consumption such as 24-hr food recalls, objective measures of physical activity such as accelerometers, and assessments of additional sources of screen time such as computers, tablets, and cell phones.

Conclusions

Overall, weight-related behaviors in elementary school appear to have little effect on weight during elementary and middle school, and are largely overshadowed by the effect of prior weight. This research may have implications for child obesity interventions, which have generally failed to achieve substantial impact on BMI (Kamath et al., 2008). Of the weight-related behaviors we examined, television viewing was the only significant predictor of

subsequent weight change, and may merit the most attention as an obesity prevention strategy. However, our finding that weight becomes highly stable by the time children reach school-age, with little influence from weight-related behaviors, may indicate that obesity prevention programs should be aimed to reach children younger than age 5. Targeting efforts toward younger children and identifying promising areas for early-life intervention may be key for developing effective policy to reduce childhood obesity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by grant no. R03 HD061509-01A from the Eunice Kennedy Shriver National Institute of Child Health & Human Development and by Emory University's University Research Committee.

The content is solely the responsibility of the authors and does not necessarily represent the official views of the Eunice Kennedy Shriver National Institute of Child Health & Human Development or the National Institutes of Health.

The authors thank Behzad Kianian for assistance with data analysis.

References

- Ahmad S, Rukh G, Varga TV, Ali A, Kurbasic A, Shungin D, Ericson U, Koivula RW, Chu AY, et al. Gene $\times$ physical activity interactions in obesity: combined analysis of 111,421 individuals of European ancestry. *PLoS genetics*. 2013; 9:e1003607. [PubMed: 23935507]
- Baird J, Fisher D, Lucas P, Kleijnen J, Roberts H, Law C. Being big or growing fast: systematic review of size and growth in infancy and later obesity. *BMJ (Clinical research ed.)*. 2005; 331:929.
- Barkley JE, Salvy SJ, Roemmich JN. The effect of simulated ostracism on physical activity behavior in children. *Pediatrics*. 2012; 129:e659–66. [PubMed: 22311997]
- Biesanz, JC. Autoregressive longitudinal models. In: Hoyle, RH., editor. *Handbook of Structural Equation Modeling*. Guilford, New York: 2012. p. 459-71.
- Bogaert N, Steinbeck KS, Baur LA, Brock K, Bermingham MA. Food, activity and family--environmental vs biochemical predictors of weight gain in children. *Eur J Clin Nutr*. 2003; 57:1242–9. [PubMed: 14506484]
- Browne, MW., Cudeck, R., Bollen, KA. *Alternative ways of assessing model fit*. Sage Focus Editions; 1993.
- Carnell S, Kim Y, Pryor K. Fat brains, greedy genes, and parent power: a biobehavioural risk model of child and adult obesity. *International review of psychiatry (Abingdon, England)*. 2012; 24:189–99.
- Carson V, Hunter S, Kuzik N, Gray CE, Poitras VJ, Chaput JP, Saunders TJ, Katzmarzyk PT, Okely AD, et al. Systematic review of sedentary behaviour and health indicators in school-aged children and youth: an update. *Applied physiology, nutrition, and metabolism = Physiologie appliquee, nutrition et metabolisme*. 2016; 41:S240–65.
- Danner FW. A national longitudinal study of the association between hours of TV viewing and the trajectory of BMI growth among US children. *Journal of pediatric psychology*. 2008; 33:1100–7. [PubMed: 18390579]
- Finney SJ, DiStefano C. Non-normal and categorical data in structural equation modeling, *Structural equation modeling: A second course*. 2006:269–314.
- Franzese A, Valerio G, Argenziano A, Esposito-Del Puente A, Iannucci MP, Caputo G, Alfonsi L, Contaldo F, Rubino A. Onset of obesity in children through the recall of parents: relationship to parental obesity and life events. *J Pediatr Endocrinol Metab*. 1998; 11:63–7. [PubMed: 9642631]

- Fuller-Tyszkiewicz M, Skouteris H, Hardy LL, Halse C. The associations between TV viewing, food intake, and BMI. A prospective analysis of data from the Longitudinal Study of Australian Children. *Appetite*. 2012; 59:945–8. [PubMed: 23000277]
- Hesketh K, Wake M, Waters E, Carlin J, Crawford D. Stability of body mass index in Australian children: a prospective cohort study across the middle childhood years. *Public health nutrition*. 2004; 7:303–9. [PubMed: 15003138]
- Huang R-C, de Klerk NH, Smith A, Kendall GE, Landau LI, Mori TA, Newnham JP, Stanley FJ, Oddy WH, et al. Lifecourse Childhood Adiposity Trajectories Associated With Adolescent Insulin Resistance. *Diabetes Care*. 2011
- Huston, AC., Ripke, MN. Developmental contexts in middle childhood: bridges to adolescence and adulthood. Cambridge University Press; 2006.
- Institute of Medicine. Preventing Childhood Obesity - Health in the Balance. The National Academies Press; Washington, DC: 2005.
- Jackson SL, Cunningham SA. Social Competence and Obesity in Elementary School. *American journal of public health*. 2015; 105:153–58. [PubMed: 25393191]
- Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. *The international journal of behavioral nutrition and physical activity*. 2010; 7:40. [PubMed: 20459784]
- Kamath CC, Vickers KS, Ehrlich A, McGovern L, Johnson J, Singhal V, Paulo R, Hettinger A, Erwin PJ, et al. Clinical review: behavioral interventions to prevent childhood obesity: a systematic review and metaanalyses of randomized trials. *The Journal of clinical endocrinology and metabolism*. 2008; 93:4606–15. [PubMed: 18782880]
- Karoutsou E, Polymeris A. Environmental endocrine disruptors and obesity. *Endocrine regulations*. 2012; 46:37–46. [PubMed: 22329821]
- La Merrill, M., Birnbaum, LS. The Mount Sinai journal of medicine. Vol. 78. New York: 2011. Childhood obesity and environmental chemicals; p. 22-48.
- Lumley T. Survey: analysis of complex survey samples. R package version 3.30-3. 2014
- Malik VS, Schulze MB, Hu FB. Intake of sugar-sweetened beverages and weight gain: a systematic review. *Am J Clin Nutr*. 2006; 84:274–88. [PubMed: 16895873]
- Mitchell JA, Pate RR, Beets MW, Nader PR. Time spent in sedentary behavior and changes in childhood BMI: a longitudinal study from ages 9 to 15 years. *Int J Obes (Lond)*. 2013; 37:54–60. [PubMed: 22430304]
- Niclasen BV, Petzold MG, Schnohr C. Overweight and obesity at school entry as predictor of overweight in adolescence in an Arctic child population. *Eur J Public Health*. 2007; 17:17–20. [PubMed: 17071950]
- Oberski D. lavaan.survey: An R package for complex survey analysis of structural equation models. *Journal of Statistical Software*. 2014; 57:1–27. [PubMed: 25400517]
- Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of childhood and adult obesity in the United States, 2011–2012. *JAMA : the journal of the American Medical Association*. 2014; 311:806–14. [PubMed: 24570244]
- Ravelli GP, Stein ZA, Susser MW. Obesity in young men after famine exposure in utero and early infancy. *N Engl J Med*. 1976; 295:349–53. [PubMed: 934222]
- Reinehr T, Andler W. Changes in the atherogenic risk factor profile according to degree of weight loss. *Archives of disease in childhood*. 2004; 89:419–22. [PubMed: 15102630]
- Rey-Lopez JP, Vicente-Rodriguez G, Biosca M, Moreno LA. Sedentary behaviour and obesity development in children and adolescents. *Nutr Metab Cardiovasc Dis*. 2008; 18:242–51. [PubMed: 18083016]
- Rosseel Y. lavaan: An R package for structural equation modeling. *Journal of Statistical Software*. 48:1–36.
- Singh AS, Mulder C, Twisk JW, van Mechelen W, Chinapaw MJ. Tracking of childhood overweight into adulthood: a systematic review of the literature. *Obesity reviews : an official journal of the International Association for the Study of Obesity*. 2008; 9:474–88. [PubMed: 18331423]
- Skelton JA, Irby MB, Grzywacz JG, Miller G. Etiologies of obesity in children: nature and nurture. *Pediatr Clin North Am*. 2011; 58:1333–54. ix. [PubMed: 22093854]

- StataCorp. Stata Statistical Software: Release 13. College Station, TX: 2013.
- Storch EA, Milsom VA, DeBraganza N, Lewin AB, Geffken GR, Silverstein JH. Peer victimization, psychosocial adjustment, and physical activity in overweight and at-risk-for-overweight youth. *J. Pediatr. Psychol.* 2007; 32:80–89. [PubMed: 16601255]
- Team, RC. Foundation for Statistical Computing. Vienna, Austria: 2014. R: A Language and Environment for Statistical Computing.
- Thibault H, Contrand B, Saubusse E, Baine M, Maurice-Tison S. Risk factors for overweight and obesity in French adolescents: physical activity, sedentary behavior and parental characteristics. *Nutrition (Burbank, Los Angeles County, Calif.)*. 2010; 26:192–200.
- Tourangeau, K., Nord, C., Lê, T., Sorongon, AG., Najarian, M. National Center for Education Statistics. Washington, DC: 2009. ECLS-K: Combined User's Manual for the ECLS-K Eighth-Grade Data Files and Electronic Codebooks (NCES 2009-004).
- Tucker JM, Tucker LA, LeCheminant J, Bailey B. Obesity increases risk of declining physical activity over time in women: a prospective cohort study. *Obesity*. 2013; 21:E715–E20. [PubMed: 23512799]
- Twisk JW, Kemper HC, van Mechelen W, Post GB, van Lenthe FJ. Body fatness: longitudinal relationship of body mass index and the sum of skinfolds with other risk factors for coronary heart disease. *Int J Obes Relat Metab Disord*. 1998; 22:915–22. [PubMed: 9756252]
- U.S. Department of Education. User's manual for the ECLS-K Base Year Public-Use Data Files and Electronic Codebook: NCES 2001-029 (revised). Author; Washington, DC: 2004.
- Vidmar SI, Cole TJ, Pan H. Standardizing anthropometric measures in children and adolescents with functions for egen: Update. *Stata Journal*. 2013; 13:366–78.
- Vogels N, Posthumus DL, Mariman EC, Bouwman F, Kester AD, Rump P, Hornstra G, Westerterp-Plantenga MS. Determinants of overweight in a cohort of Dutch children. *Am J Clin Nutr*. 2006; 84:717–24. [PubMed: 17023696]
- Waters E, de Silva-Sanigorski A, Hall BJ, Brown T, Campbell KJ, Gao Y, Armstrong R, Prosser L, Summerbell CD. Interventions for preventing obesity in children. *The Cochrane database of systematic reviews*:Cd001871. 2011
- West S, Taylor A, Wu W. Model fit and model selection in structural equation modeling, *Handbook of structural equation modeling*. 2012:209–31.
- World Health Organization. Report of the Commission on Ending Childhood Obesity. Geneva, Switzerland: 2016.
- Wright CM, Emmett PM, Ness AR, Reilly JJ, Sherriff A. Tracking of obesity and body fatness through mid-childhood. *Archives of disease in childhood*. 2010; 95:612–7. [PubMed: 20522467]

Highlights

- Child obesity interventions often focus on television, physical activity, and diet.
- Most health-related behaviors had little impact on weight trajectories.
- Screen time was associated with subsequent weight gain.
- Children's BMI z-score were highly stable across middle childhood.
- Higher BMI z-score was associated with subsequent adoption of unhealthy behaviors.

Figure 1.

Relationships between BMI, Diet, and Activities across Elementary School: Standardized Parameter Estimates from a Cross-lagged Autoregressive Model

Abbreviations: TV_p: Hourly television viewing per day, reported by the parent; TV_C: Hourly television viewing per day, reported by the child; Exer_p: Number of days with 20 minutes exercise, reported by the parent; Exer_C: Number of days with 20 minutes exercise, reported by the child; BMI: BMI z-score; FV_C: fruit and vegetable intake latent variable (intake reported by the child); SF_C: Soda and fast food intake latent variable (intake reported by the child)

Data source: Early Childhood Longitudinal Study, Kindergarten Class of 1998–1999, U.S. cohort followed 1998–2007 (n=4,938)

Table 1

Baseline Characteristics of Kindergarteners

Characteristic	Mean or %	SE
Gender		
Male	48.8	0.9
Female	51.2	0.9
Race/ethnicity		
Non-Hispanic White	60.3	1.8
Non-Hispanic Black	14.3	1.1
Hispanic	18.8	1.6
Other	6.6	0.7
Weight Status in Kindergarten		
Normal	73	0.9
Overweight	14.5	0.7
Obese	12.5	0.7
BMI Z-score in Kindergarten	0.415	0.03

Survey-adjusted results. Data: Early Childhood

Data source: Early Childhood Longitudinal Study, Kindergarten Class of 1998–1999, U.S. cohort followed 1998–2007(n=4,938)

Table 2**Food Consumption in Fifth and Eighth Grade**

Characteristic	5 th grade		8 th grade	
	Percent	SE	Percent	SE
Soda				
Did not eat during past 7 days	15.2	0.88	14.55	0.77
1 to 3 times per week	39.94	1.03	42.96	1.05
4 to 6 times per week	17.27	0.66	16.3	0.81
1 time per day	10.09	0.49	10.93	0.62
2 times per day	7.72	0.6	8.24	0.58
3 times per day	3.15	0.35	4.04	0.37
4 or more times per day	6.63	0.63	2.98	0.41
Fast Food				
Did not eat during past 7 days	28.27	1.06	28.57	1.32
1 to 3 times per week	51.75	1.33	54.71	1.22
4 to 6 times per week	10.35	0.74	9.38	0.59
1 time per day	4.89	0.51	3.62	0.45
2 times per day	2.11	0.39	2.13	0.42
3 times per day	0.66	0.19	1.07	0.35
4 or more times per day	1.96	0.36	0.52	0.14
Fruits				
Did not eat during past 7 days	8.64	0.52	7.95	0.61
1 to 3 times per week	31.63	1.09	27.07	0.92
4 to 6 times per week	21.48	0.97	24.86	1.04
1 time per day	13.82	0.81	15.67	0.79
2 times per day	11.92	0.8	14.42	0.78
3 times per day	5.68	0.39	5.36	0.47
4 or more times per day	6.84	0.46	4.67	0.6
Green Salads				
Did not eat during past 7 days	50.68	1.18	36.81	1.03
1 to 3 times per week	32.81	1.21	40.42	1.12
4 to 6 times per week	7.33	0.47	11.4	0.63
1 time per day	6.15	0.44	7.36	0.57
2 times per day	1.41	0.15	2.32	0.36
3 times per day	0.48	0.11	0.79	0.23
4 or more times per day	1.15	0.25	0.9	0.3
Carrots				
Did not eat during past 7 days	46.21	1.03	51.41	1.3
1 to 3 times per week	32.2	1.1	35.47	1.15
4 to 6 times per week	9.35	0.55	6.77	0.46
1 time per day	5.93	0.46	3.87	0.39
2 times per day	1.97	0.22	1.11	0.22

Characteristic	5 th grade		8 th grade	
	Percent	SE	Percent	SE
3 times per day	1.66	0.27	0.7	0.14
4 or more times per day	2.68	0.37	0.67	0.17
Other Vegetables				
Did not eat during past 7 days	17.3	0.89	12.09	0.78
1 to 3 times per week	36.83	1.05	36.34	1.06
4 to 6 times per week	20.24	0.83	25.01	0.9
1 time per day	12.07	0.71	14.46	0.72
2 times per day	6.66	0.6	7.76	0.54
3 times per day	3.02	0.37	2.69	0.38
4 or more times per day	3.88	0.47	1.66	0.31

Survey-adjusted results.

Data source: Early Childhood Longitudinal Study, Kindergarten Class of 1998–1999, U.S. cohort followed 1998–2007 (n=4,938)

Table 3

Television and Physical Activity Behavior and Obesity Status

Characteristic	Kindergarten Mean (SE)	1 st grade Mean (SE)	3 rd grade Mean (SE)	5 th grade Mean (SE)	8 th grade Mean (SE)
TV Per Day, hours (All)	2.00 (0.041)	2.14 (0.035)	2.16 (0.028)	2.28 (0.031)	3.32 (0.065)
Normal	1.95 (0.037)	2.08 (0.034)	2.08 (0.031)	2.15 (0.031)	3.23 (0.089)
Overweight	2.01 (0.080)	2.30 (0.139)	2.23 (0.060)	2.33 (0.085)	3.30 (0.116)
Obese	2.28 (0.085)	2.33 (0.069)	2.39 (0.057)	2.61 (0.065)	3.64 (0.116)
Exercise, days per week	3.95 (0.059)	N/A	3.99 (0.038)	3.78 (0.048)	4.65 (0.050)
Normal	3.96 (0.063)	N/A	4.05 (0.048)	3.88 (0.061)	4.72 (0.054)
Overweight	3.94 (0.138)	N/A	4.12 (0.096)	3.86 (0.099)	4.61 (0.090)
Obese	3.94 (0.128)	N/A	3.67 (0.096)	3.46 (0.090)	4.44 (0.117)

Survey-adjusted results.

Weight status categories were determined by percentile associated with BMI-Z-score (obese 95%, overweight 85–94%)

Data source: Early Childhood Longitudinal Study, Kindergarten Class of 1998–1999, U.S. cohort followed 1998–2007 (n=4,938)