CORRESPONDENCE

Chronic Abdominal Wall Pain—a Poorly Recognized Clinical Problem

by Prof. Dr. med. Herbert Koop, Dr. Simona Koprđova, and Dr. med. Christine Schürmann in issue 4/2016

Intercostal Nerve Course Different

I would like to thank the authors for their very interesting article (1). Since, according to the literature, abdominal pain originates from the abdominal wall in 2% to 30% of cases, attention should once again be paid the intercostal nerves. However, the actual course of the intercostal nerves in this area differs significantly from that depicted in the figure.

The intercostal nerves run, coming from dorsal cranial and descending in a ventral caudal direction between the internal oblique and the transverse abdominal muscle. They penetrate the rectus sheath obliquely and then lie within the rectus sheath lateral to the rectus abdominis muscle. Their direction does not change during this passage. Since the intercostal nerves are enveloped by a tunnel of firm connective tissue during their oblique path through the rectus sheath, this is the site where nerve compression may occur.

Within the rectus sheath, the intercostal nerves first run between the rectus abdominis muscle and the posterior lamina of the rectus sheath. In most cases, they split into a medial and a lateral branch before they enter the rectus abdominis muscle from dorsal at an angle, then pass through it obliquely while innervating it, then exit it on its anterior surface to finally penetrate through the anterior lamina of the rectus sheath. I think compression at the exit points are less relevant. If this happened, skin nerves would then also be compressed at other sites of the body, resulting in superficial pain.

However, this course of the intercostal nerve does not make any difference to the symptoms described in the article. Nevertheless, I think it is of help to present the anatomical background information correctly.

DOI: 10.3238/arztebl.2016.0503a

REFERENCES

1. Koop H, Koprđova S, Schürmann C: Chronic abdominal wall pain—a poorly recognized clinical problem. *Dtsch Arztebl Int* 2016; 113: 51–7.

Dr. med. Cornelius Lemke
Institut für Anatomie I
Jena, Germany
Cornelius.Lemke@med.uni-jena.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Chronic Pain in the Rectus Abdominis Muscle

The authors mention several treatment options in their article: local anesthetic injection, where indicated in combination with corticosteroids; neurolysis with phenol; surgical neurectomy (1). These recommendations are based on reports in the literature that pain relief can be achieved by injecting lidocaine into “the points of maximal abdominal pain beneath the anterior fascia of the rec-

tus abdominis muscle”. In a double-blind, randomized study, 22 patients underwent neurectomy, while another 22 patients underwent sham surgery. In the neurectomy group, 16 patients were free of pain after the procedure, in contrast to 4 patients in the sham surgery group. Complications included 5 hematomas, 1 infection and 1 case with increased pain (2). After 8 years, 71 of the 181 patients still experienced pain (3).

This category of abdominal wall pain comprises chronic pain in the rectus abdominis muscle through which the rami cutanei anteriores abdominales pass to reach the anterior lamina of the rectus sheath, whereas at other locations of the abdominal wall pain, such as the linea semilunaris, linea alba and in the subcostal region, nerves of this type are not present. In these regions, one cannot expect to achieve the same results with neurectomy that one would want to achieve at the rectus abdominis muscle. According to standard criteria, abdominal wall pain is classed as a chronic pain disorder (ICD diagnosis code F45.41). In the review of Koop et al. (1), a cohort study, reporting on classification, diagnosis and cognitive behavioral therapy in 55 patients with abdominal wall pain (4), is not mentioned. Even though it was not possible to conduct this study as a controlled trial, and consequently the observed response to treatment is not considered to be evidence-based, it is unreasonable to recommend to patients with chronic abdominal wall pain that they should undergo an excision of nerve branches in the abdominal wall. The “success” rate of neurectomy is discussed in reference (3). Chronicity of pain leads to the generation of a pain memory in the central nervous system which signals pain independently of peripheral lesions. Thus, surgical treatment is not only ineffective, it may even aggravate the pain. In these patients, behavioral therapy is indicated, including multimodal treatment approaches. Without the discussion of this alternative concept of diagnosis and treatment, this review could appear misleading.

DOI: 10.3238/arztebl.2016.0503b

REFERENCES

1. Koop H, Koprđova S, Schürmann C: Chronic abdominal wall pain—a poorly recognized clinical problem. *Dtsch Arztebl Int* 2016; 113: 51–7.
2. Boelens OB, van Assen T, Houterman S, Scheltinga MR, Roumen RM: A double-blind, randomized, controlled trial on surgery for chronic abdominal pain due to anterior cutaneous nerve entrapment syndrome. *Ann Surg* 2013; 257: 845–9.
3. van Assen T, Boelens OB, van Eerten PV, Scheltinga MR, Roumen MR: Surgical options after a failed neurectomy in anterior cutaneous nerve entrapment syndrome. *World J Surg* 2014; 38: 3105–11.
4. Feurle GE: Abdominal wall pain—classification, diagnosis and treatment suggestions. *Wien Klin Wochenschr* 2007; 119: 633–8.

Prof. Dr. med. Gerhard E. Feurle
DRK Krankenhaus Neuwied, Germany
g.e.feurle@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists.

Differentiation of Etiology Missing

Agreeably, the article elaborates on abdominal pain symptoms which are from time to time encountered in clinical practice where they frequently present diagnostic and therapeutic challenges (1).

However, the—apparently relevant—neuropathic aspect of these symptoms is not differentiated in the article. Unfortunately, the recommended diagnostic local anesthetic injections do not contribute to a further differentiation of the etiology of the pain either. The invasive treatment options recommended in the article should be regarded as rather disadvantageous due to the always temporary nature of their effect and the risk of further chronification. Especially neurectomy and neurectomy can trigger clinically very challenging exacerbations of neuropathic pain and thus are no longer in line with today's treatment recommendations (2).

For this reason, the 2012 guideline "Pharmacological, non-interventional management of chronic neuropathic pain" excludes interventional therapy already in its title and it is not part of any other guideline either (3). With respect to isolated neuropathic pain syndromes, Dworkin et al. stress in their broad review that there is a lack of standardized study data in the literature, which primarily consists of case reports (4).

Thus, one cannot agree with the authors' statement that "fears of adverse effects [of injection therapy] are unfounded". I am critical about the statements that in children, some patients "needed multiple injection" and "surgical therapy was particularly effective", not least because of the frequently functional aspects of abdominal pain in pediatric patients. Adequate treatment options are based on non-surgical pain management, including, among others, anti-neuropathic medications and local treatments or TENS, where necessary embedded in a multi-modal overall plan.

DOI: 10.3238/arztebl.2016.0504a

REFERENCES

1. Koop H, Koprdova S, Schürmann C: Chronic abdominal wall pain—a poorly recognized clinical problem. *Dtsch Arztebl Int* 2016; 113: 51–7.
2. Tronnier VM, Rasche D: Neurodestruktive Verfahren in der Schmerztherapie. *Schmerz* 2009; 23: 531–43.
3. Diener H-C, Weimar C: Pharmakologische nichtinterventionelle Therapie chronisch neuropathischer Schmerzen. In: Diener HC, Maier C (eds.): Leitlinien der Diagnostik und Therapie in der Neurologie. Stuttgart: Thieme, 2012.
4. Dworkin RH, O'Connor AB, Kent J, et al.: Interventional management of neuropathic pain: NeuPSIG recommendations. *Pain* 2013; 154: 2249–61.

Dr. med. Thorsten Nickel
imland Klinik Rendsburg, Germany
thorsten.nickel@imland.de

Conflict of interest statement

Dr. Nickel has received reimbursement of conference fees and travel expenses from Grünenthal and Mundipharma.

In Reply:

The above letters from readers as well as the many comments and questions we have received directly are testimony to the great interest in this long-neglected clinical problem which has not been studied systematically to a greater extent as yet. We would like to thank Dr. C. Lemke for contributing more precise

anatomical details; however, he does not think that they make any difference to the diagnostic and therapeutic procedure.

Prof. G. Feurle sees the risk of excessive and frequently not indicated surgical treatment, referring to a case series study he conducted. However, his sample included a significant number of patients with irritable bowel syndrome or depression/anxiety disorders (1). Among these patients, abdominal wall pain is indeed present, but frequently covers a larger area. A history of an association between parietal abdominal wall pain and visceral symptoms (e.g. flatulence) or an influence of food intake or defecation should always raise doubts about the correctness of the diagnosis of abdominal wall pain as defined in our article; the same applies to patients with overt psychopathology. Dr. T. Nickel's comment follows a similar line of argument. He is concerned about a lack of consideration of neuropathic mechanisms, potentially contributing to the etiology of abdominal wall pain, and warns of the use of invasive treatment options. We had already discussed the differential diagnosis of neuropathic pain in detail elsewhere (2, 3). Crucial for diagnosing abdominal wall pain in the sense of a focal nerve alteration—more precisely described by the term "anterior cutaneous nerve entrapment syndrome"—is the clearly defined area of no more than 2 cm in diameter in which the pain can be reproducibly triggered. This pain does not cover larger areas and is not usually referred elsewhere (only occasionally in case of an alteration of the ilioinguinal / iliohypogastric nerve). Relevant data on the non-surgical pain management of this condition are not available; from our experience, prior non-surgical treatment was unsuccessful in many patients. In contrast, injection therapy with local anesthetics has eliminated all symptoms in many patients and thus we think it should be regarded as the first-line treatment.

The strategy for non-responders, where multiple injections brought only temporary relief and the pain is always at the same location, remains unclear; here—as discussed in our article—a great need for further research remains, which also applies to identifying the best possible treatment approach in children. To proceed to surgical treatment after one unsuccessful injection therapy (as pursued by the Dutch working group) is not recommended; the indication for neurectomy must always be based on a very critical evaluation of potential benefits and risks and the decision should never be taken lightly; here, a repeat differential diagnostic evaluation is considered an indispensable prerequisite.

DOI: 10.3238/arztebl.2016.0504b

REFERENCES

1. Feurle GE: Abdominal wall pain—classification, diagnosis and treatment suggestions. *Wien Klin Wochenschr* 2007; 119:633–638
2. Koop H, Schürmann C: Chronischer Bauchwandsschmerz. *Gastroenterol up2date* 2014; 10: 129–140
3. Koop H, Koprdova S, Schürmann C: Chronic abdominal wall pain—a poorly recognized clinical problem. *Dtsch Arztebl Int* 2016; 113: 51–7.

Prof. Dr. med. Herbert Koop
Dr. med. Simona Koprdova
Dr. med. Christine Schürmann

Klinik für Allgemeine Innere Medizin und Gastroenterologie, Helios Klinikum Berlin-Buch, Germany
prof.koop@t-online.de

Conflict of interest statement

The authors declare that no conflict of interest exists.