Electronic data protection: procedures need drastic improvement

EDITOR—Gulena et al are correct to challenge overzealous interpretation of data protection law in their questionnaire survey of patients' and visitors' views about having their names displayed in hospital.¹ Measures taken by doctors to safeguard patient confidentiality, especially electronic records, are, however, severely deficient and fuel the concerns of those entrusted to police data security.

We invited 32 surgical trainees to complete a questionnaire regarding their Data Protection Act registration and electronic data confidentiality practices.

Of 29 responders, 26 trainees regularly computerised and stored patients' data. One person was registered with the Data Protection Act. Only three of 14 desktops, eight of 19 laptops, and three of 14 handheld computers forced a password logon. Sixteen of 29 trainees used the same password for all machines, and 25 of 27 passwords were less than eight characters long. All desktops, 16 of 19 laptops, and five of 14 handhelds were routinely connected to the internet, and half of these had not had their online security settings adjusted. Of 29 trainees, 28 did not encrypt their sensitive data files. Ten trainees had sent patients' data unencrypted over the internet, using a non-secure server.

Electronic data confidentiality practices among surgical trainees are unsafe, and this is unlikely to be confined to our study population. A breach in confidentiality would have widespread implications for public trust. Clinicians have a legal obligation to implement adequate medical IT security.²⁻⁴ Existing passwords should be improved, and biometric logon systems (fingerprint scanning) considered. Files containing sensitive data should be routinely encrypted. Magnetic and solid state computer storage devices (including hard disks, floppy disks, CD Roms, DVD Roms, and memory sticks) should undergo routine physical or electronic data shredding after their final use.⁵ We are implementing medical information technology security training for this particu-

lar group of doctors and would urge others to initiate similar programmes before a serious breach occurs.

Damian Mole *research fellow, surgery*
Department of Surgery, Queen's University Belfast,
Belfast BT12 6BJ
damianmole@ntlworld.com

Colin Fox *information technology manager*
Giulio Napolitano *information technology and security manager*
Northern Ireland Cancer Registry, Department of Epidemiology and Public Health, Queen's University Belfast

Competing interests: None declared.

- 1 Gulena R, Luwemba S, Williams A, Jenkinson LR. Data protection gone too far: questionnaire survey of patients' and visitors' views about having their names displayed in hospital. *BMJ* 2004;329:1491. (18-25 December.)
- 2 General Medical Council. *Maintaining good medical practice*. London: GMC; 2003.
- 3 The Council of Europe. *The protection of medical data*. Recommendation No R (97) 5 and explanatory memorandum. English edition. Strasbourg: Council of Europe Publishing, 1997.
- 4 Information Commissioner's Office. *Do I need to notify?* Wilmslow: IC, 2005. www.informationcommissioner.gov.uk/eventual.aspx?id=2662 (accessed 24 Feb 2005).
- 5 Gutmann P. *Secure deletion of data from magnetic and solid-state memory*. www.cs.auckland.ac.nz/~pgut001/pubs/secure_del.html (accessed 24 Feb 2005).

Independent medical school is long overdue

EDITOR—The concept of the United Kingdom's first independent medical school at the University of Buckingham has created great excitement.¹ We believe that the time is right for this development and that it mirrors changes in healthcare delivery, education, and society. The news article by Cohen and Fister focused on two quotations that neither address the real issues nor further the debate on future education.

Buckingham is Britain's only independent, not for profit, university, with successful courses in business, finance, education, and law. More doctors are urgently needed globally. Can we achieve this without burdening the taxpayer? We believe we can, and increase the diversity among tomorrow's doctors at the same time.

The commentary highlights two issues: cost and speed. The cost to the student is little greater than that already charged to overseas students as the fast tracking keeps the total cost low. With continuing increases in fees for home students, the financial difference will reduce. We will provide bursaries and loans. We are looking for the most able and motivated and nothing else.

The course will entail as many hours' study as the conventional graduate course. The first year will consist of an innovative distance learning package with residential blocks, followed by three full time years. Professors Janet Grant and Michael Stewart from the Open University are members of this board and are experts in this area. We have already been in constructive discussion with the General Medical Council to meet its rigorous standards.

We need a new approach to encourage entry into medicine of socially aware, successful people with a diverse range of experience. An independent university is the ideal crucible for this.

Helen Nellis *chair*

Karol Sikora *dean elect*
karolsikora@hotmail.com
University of Buckingham Medical School
Development Board, Buckingham MK18 1EG

Competing interests: Both authors are members of the University of Buckingham Development Board.

- 1 Cohen D, Fister K. Rejecting political correctness. *BMJ* 2005;330:62. (8 January.)

Gastric rupture and laryngeal mask airway

Negative "hype" this case report might generate is worrying

EDITOR—Case reports have always been useful in increasing awareness of some issues not apparent when new procedures or techniques are introduced. Haslam et al describe gastric rupture associated with use of the laryngeal mask airway during cardiopulmonary resuscitation.¹ I believe that the use of the laryngeal mask airway has been pivotal in managing the airway by non-anaesthetists, especially in prehospital settings. Recent literature agrees that attempting to use orotracheal intubation in prehospital environments actually worsens outcomes compared with "simpler" methods such as bag-valve mask ventilation.

I am concerned that the "hype" that this case report might generate may deter people from using a device as excellent as the laryngeal mask airway because of concerns of producing gastric rupture and pneumoperitoneum.

This report should be used as a reminder that any tool can potentially produce harm, and that the benefits of this tool's prehospital use exceed the harm by far. Tidal volumes and insufflation pressures should always be monitored, no matter which airway device is

bmj.com

Letters appearing here are an edited selection of rapid responses originally posted on bmj.com

We ask for all letters to the editor to be submitted as rapid responses via bmj.com

For advice see: bmj.com/rapidresponses

used; and appropriate resuscitation bags to facilitate this should be made available.

Pedro F Ibarra *trauma, anaesthesia, and critical care consultant*
Clinica Reina Sofia, Bogota, Colombia 1
pfibarra@yahoo.com

Competing interests: None declared.

1 Haslam N, Campbell GC, Duggan JE. Gastric rupture associated with the use of the laryngeal mask airway during cardiopulmonary resuscitation. *BMJ* 2004;329:1225-6. (20 November.)

Laryngeal mask airway was not likely cause of gastric rupture

EDITOR—As some of the people responsible for setting resuscitation guidelines, we are interested in the case report of Haslam et al describing gastric rupture after use of the laryngeal mask airway during cardiopulmonary resuscitation.¹

There is a risk of gastric inflation with all airway and ventilation techniques during cardiopulmonary resuscitation. Positive pressure ventilation via a laryngeal mask airway can cause gastric inflation, particularly if the airway is not positioned correctly and if the inspiratory pressure exceeds 20 cm H₂O.²⁻⁴ Most gas leaking from the airway escapes into the pharynx to exit through the mouth.³

In the case described the laryngeal mask airway is not the most likely cause of gastric rupture. The patient was given seven minutes of cardiopulmonary resuscitation by a bystander: chest compressions alone can cause gastric rupture, particularly if delivered too low on the sternum or upper abdomen. Furthermore, mouth to mouth or bag-valve mask ventilation is much more likely to cause gastric inflation than ventilation with a laryngeal mask airway: incorrect alignment of the airway will cause gastric inflation because the mask or rescuer's mouth prevents gas escaping through the patient's mouth, and high airway pressures will be transmitted entirely to the oesophagus.⁴ For this reason, using an laryngeal mask airway instead of a face mask to ventilate the lungs during cardiopulmonary resuscitation reduces the incidence of regurgitation.⁵

The advice given in the advanced life support course manual to attempt to undertake continuous chest compressions, uninterrupted during ventilation through the laryngeal mask airway, is based on evidence that even a brief pause in chest compressions causes a significant reduction in coronary perfusion pressure. If excessive gas leakage occurs, chest compressions will have to be interrupted to enable ventilation of the lungs. The precise risk of gastric inflation when using the laryngeal mask airway under these circumstances is unknown and should be studied.

Jerry P Nolan *consultant in anaesthesia and intensive care medicine*
Royal United Hospital, Bath BA1 3NG
jerry.nolan@ukgateway.net

Mick Colquhoun *senior lecturer in prehospital care*
Wales School of Medicine, Cardiff University,
Cardiff CF14 4XX

Charles D Deakin *consultant anaesthetist*
Southampton General Hospital, Southampton,
Hampshire SO16 6YD

The other four authors of this letter are David A Gabbott, consultant anaesthetist, Department of Anaesthetics, Gloucestershire Hospitals NHS Trust, Gloucester GL1 3NN; Andrew Lockey, consultant in accident and emergency medicine, Calderdale Royal Hospital, Salterhebble, Halifax HX3 0PW; Sarah Mitchell, director, Resuscitation Council (UK), London WC1H 9HR; Jasmeet Soar, consultant in anaesthesia and intensive care medicine, Southmead Hospital, North Bristol NHS Trust, Bristol BS10 5NB.

Competing interests: The authors are members of the Resuscitation Council (UK).

- 1 Haslam N, Campbell GC, Duggan JE. Gastric rupture associated with the use of the laryngeal mask airway during cardiopulmonary resuscitation. *BMJ* 2004;329:1225-6. (20 November.)
- 2 Latorre F, Eberle B, Weiler N, Mienert R, Stanek A, Goedecke R, et al. Laryngeal mask airway position and the risk of gastric insufflation. *Anesth Analg* 1998;86:867-71.
- 3 Ho-Tai LM, Devitt JH, Noel AG, O'Donnell MP. Gas leak and gastric insufflation during controlled ventilation: face mask versus laryngeal mask airway. *Can J Anaesth* 1998;45:206-11.
- 4 Weiler N, Latorre F, Eberle B, Goedecke R, Heinrichs W. Respiratory mechanics, gastric insufflation pressure, and air leakage of the laryngeal mask airway. *Anesth Analg* 1997;84:1025-8.
- 5 Stone BJ, Chantler PJ, Baskett PJ. The incidence of regurgitation during cardiopulmonary resuscitation: a comparison between the bag valve mask and laryngeal mask airway. *Resuscitation* 1998;38:3-6.

Authors' reply

EDITOR—We wanted to highlight the risk of gastric inflation when the laryngeal mask airway is used for positive pressure ventilation and to draw attention to the inadequacy of the current Resuscitation Council (UK) guidelines for using the laryngeal mask airway in resuscitation.¹ We did not intend to generate any negative "hype" about its use by non-anaesthetists.

Limiting airway pressure to less than 20 cm H₂O to minimise the risk of gas leak when ventilating a patient using the laryngeal mask airway is important. Standard resuscitation equipment is, however, not equipped to achieve this easily. Therefore practitioners must be aware of the issues involved in the use of the device and have the appropriate degree of skill to use it safely. For this reason we take issue with some statements and recommendations given in current resuscitation guidelines for use of the laryngeal mask airway.

We accept that most gas leaking from around the laryngeal mask airway during lung inflation will indeed exit through the mouth, but this misses the point. Only a small proportion of the total gas leak need enter the stomach to cause substantial distension: 600 ml gas will enter the stomach in only 20 breaths with a tidal oesophageal leak of just 30 ml (<5% of recommended tidal volume).

Furthermore, continuous, uninterrupted chest compression (100 per minute) will cause a surge in airway pressure 2-3 times per inspiration (inspiratory time 1-2 s). This is incompatible with the need to limit airway pressure, and makes a pharyngeal gas leak almost certain to occur. The current guideline dismisses the risk of gastric inflation and states that the only reason to cease continuous chest compression is if gas leakage results in inadequate ventilation.¹

Practitioners trained to use the laryngeal mask airway should be taught that the presence of a pharyngeal gas leak during lung inflation is indicative of an oesophageal leak and should be actively sought for and corrected. They should be instructed, given there is no obvious leak through the mouth, to listen with a stethoscope over the anterior aspect of the neck to exclude pharyngeal gas leaks of lesser flow. We believe that if any gas leak is present cardiac compressions should be interrupted for ventilation (15:2).

Nathaniel Haslam *specialist registrar in anaesthesia*,
Freeman Hospital, Newcastle upon Tyne NE7 7DN
nhaslam@doctors.org.uk

John E Duggan *consultant anaesthetist*
Department of Anaesthesia, Wansbeck Hospital,
Ashington, Northumberland NE63 9JJ

Competing interests: None declared.

1 Resuscitation Council (UK). Adult advanced life support. Resuscitation guidelines 2000. www.resus.org.uk/pages/als.htm#ccav (accessed 29 Jan 2005).

Surgical management of metastatic inguinal lymphadenopathy

The term deep inguinal nodes should be abandoned

EDITOR—Swan et al present an interesting review,¹ but their technical recommendations are inconsistent with our experience of groin dissection over 20 years. Evidence for preferential use of straight oblique incisions is minimal.

Tonouchi et al studied only 25 procedures, with eight S-shaped incisions and 17 "straight obliques."² The learning curve and lack of information on the case mix make it difficult to draw conclusions. Our 95 groin dissections using a "lazy S" incision caused one case of wound ischaemia, one case of wound dehiscence, and five postoperative infections. These results compare favourably with studies in which straight oblique incisions were used. An appropriately placed lazy S incision allows excellent access to the apex of the femoral triangle without compromising healing.

Swan et al say that preservation of the long saphenous vein reduces lymphoedema. Zhang et al showed reduced complications in such patients.³ However, a 70% incidence of lymphoedema in the group receiving vein ligation is surprisingly high. Baas et al performed 151 groin dissections sacrificing saphenous vein and noted a 20% incidence of lymphoedema.⁴

Support for sartorius transposition was also disappointing. There was no reduction in wound morbidity by this approach in a prospective randomised study.⁵ We never use sartorius transposition. The technique dates from times when inguinal lymphadenectomy was performed as a "wide excision," requiring split skin grafting.

Anatomists and surgeons who perform inguinal node clearance dispute the presence of deep inguinal nodes. We believe

that skeletonisation of the femoral vessels clears the femoral triangle completely. The term "deep inguinal nodes" should be abandoned.

Simon C Gibson senior house officer in general and vascular surgery
simonc@bham.ac.uk

Stephen Kettlewell specialist registrar in general and vascular surgery

Dominique S Byrne consultant general and vascular surgeon

Alan J McKay consultant general and vascular surgeon

Department of General and Vascular Surgery,
Gartnavel General Hospital, Glasgow G12 0YN

Competing interests: None declared.

- Swan MC, Furniss D, Cassell OD. Surgical management of metastatic inguinal lymphadenopathy. *BMJ* 2004;329:1272-6. (27 November.)
- Tonouchi H, Ohmori Y, Kobayashi M, Konishi N, Tanaka K, Mohri Y, et al. Operative morbidity associated with groin dissections. *Surg Today* 2004;34:413-8.
- Zhang SH, Sood AK, Sorosky JJ, Anderson B, Buller RE. Preservation of the saphenous vein during inguinal lymphadenectomy decreases morbidity in patients with carcinoma of the vulva. *Cancer* 2000;89:1520-5.
- Baas PC, Koops HS, Hoekstra HJ, van Bruggen JJ, van der Weele LT, Oldhoff J. Groin dissection in the treatment of lower-extremity melanoma. *Arch Surg* 1992;127:281-6.
- Judson PL, Jonson AL, Paley PJ, Bliss RL, Murray KP, Downs Jr LS, et al. A prospective, randomized study analyzing sartorius transposition following inguinal-femoral lymphadenectomy. *Gynecol Oncol* 2004;95:226-30.

Sentinel node biopsy has cost implications for the NHS

EDITOR—In their review of the surgical management of metastatic inguinal lymphadenopathy Swan et al describe sentinel lymph node biopsy as a promising technique, which may refine the indications for inguinal lymphadenectomy and provide better prognostic information, and reduce morbidity.¹

Current UK melanoma guidelines on the role of sentinel nodes in melanoma are inadequate.² Sentinel node biopsy remains a contentious issue with current guidelines recommending it is offered in the context of a clinical trial. It is becoming an integral part of the management and staging of primary melanoma despite no proved survival benefit. To assess the cost to the NHS for sentinel node biopsy for melanoma, we reviewed 100 consecutive patients in our unit. We assessed the costs of inpatient stay, theatre time, histology, nuclear medicine, follow up, completion lymphadenectomies, and the management of complications. Our calculated total cost of sentinel node biopsy was £1550 per patient. If sentinel node biopsy was offered to the 3600 newly diagnosed intermediate and thick melanoma patients annually in the United Kingdom,³ the NHS would face an estimated annual cost of £5.6m. Increased provision will need to be made in every centre managing melanoma, or centres with existing facilities will need additional resources.

Developing a sentinel node service, which includes nuclear medicine staffing, is a substantial cost with no proved survival benefit. The financial implications for the NHS of sentinel node biopsy in the management of malignant melanoma warrant its inclusion in the service guidance for the management of skin tumours including melanoma that the National Institute for Clinical Excellence is preparing.⁴

Negin Shamsian registrar, plastic surgery
nshamsian@hotmail.com

Stephen Hamilton specialist registrar, plastic surgery

Apul Parikh registrar, plastic surgery

Peter E M Butler consultant plastic surgeon

Royal Free Hospital, London NW3 2QG

Competing interests: None declared.

- Swan MC, Furniss D, Cassell OD. Surgical management of metastatic inguinal lymphadenopathy. *BMJ* 2004;329:1272-6. (27 November.)
- Newton Bishop JA, Corrie PG, Evans J, Gore ME, Hall PN, Kirkham N, et al. UK guidelines for the management of cutaneous melanoma. *Br J Plast Surg* 2002;55:46-54.
- Office for National Statistics. Cancer Registration Statistics, England 2000. London: ONS, 2003.
- National Institute for Clinical Excellence. Clinical guidelines in progress: skin cancer including melanoma. <http://www.nice.org.uk/page.aspx?o=33927> (accessed 12 February 2005).

Authors' reply

EDITOR—Gibson et al's experience of groin dissection compares favourably with the published data. Our preference for an oblique incision is based on an understanding of the blood supply to the groin; the vascular branches lie parallel to the natural skin creases within Camper's fascia, thus an oblique incision minimises their disruption.¹

Furthermore, we find that access to the femoral triangle is straightforward by using the oblique incision. We perform a sartorius transposition in order to provide femoral vessel coverage in the event of wound dehiscence or necrosis, not to reduce wound morbidity itself. Although possibly overcautious, in light of the high rate of wound complications reported in the literature, we consider this a simple and prudent measure. Cognisant that the evidence supporting long saphenous vein preservation is limited, we were mindful to indicate that this technique may reduce post-operative lymphoedema; it is not routinely performed in our unit.

We concur with Gibson et al that femoral vessel skeletonisation clears the femoral triangle completely. This entails removal of the nodes that lie superficial to the deep fascia and also those that lie deep to this layer and medial to the femoral vein.

Shamsian et al hint at the paradox between the UK guidelines for the management of malignant melanoma²—which say that sentinel lymph node biopsy should be offered in the context of a clinical trial—and the American Joint Committee on Cancer's staging system,³ which includes the presence

of lymph node micrometastasis (positive on sentinel lymph node biopsy) as a staging criterion. The profound cost implications to the NHS of this procedure must be taken into account by the National Institute for Clinical Excellence when preparing its guidance.

There is evidence for a survival benefit of sentinel lymph node biopsy,⁴ although the publication in 2006 of a large scale, multicentre, selective lymphadenectomy trial should definitively resolve this issue.⁵

Marc C Swan Royal College of Surgeons of England
surgical research fellow
marc.swan@surgery.oxford.ac.uk

Dominic Furniss clinical research fellow

Oliver C S Cassell consultant surgeon

Department of Plastic and Reconstructive Surgery,
Radcliffe Infirmary, Oxford OX2 6HE

Competing interests: MCS and OCSC are investigators for the Oxford Tisseel (Baxter Healthcare, Newbury, United Kingdom) trial, a prospective randomised controlled trial to determine whether fibrin sealant can reduce post-operative complications following axillary and inguinal lymphadenectomy. They receive no financial remuneration for their work.

- Spratt J. Groin dissection. *J Surg Oncol* 2000;73:243-62.
- Roberts DL, Anstey AV, Barlow RJ, Cox NH, Newton Bishop JA, Corrie PG, et al. U.K. guidelines for the management of cutaneous melanoma. *Br J Dermatol* 2002;146:7-17.
- Balch CM, Buzaid AC, Soong SJ, Atkins MB, Cascinelli N, Coit DG, et al. Final version of the American Joint Committee on Cancer staging system for cutaneous melanoma. *J Clin Oncol* 2001;19:3635-48.
- Kretschmer L, Hilgers R, Mohrle M, Balda BR, Breuninger H, Konz B, et al. Patients with lymphatic metastasis of cutaneous malignant melanoma benefit from sentinel lymphadenectomy and early excision of their nodal disease. *Eur J Cancer* 2004;40:212-8.
- Morton DL, Thompson JF, Essner R, Elashoff R, Stern SL, Nieweg OE, et al. Validation of the accuracy of intraoperative lymphatic mapping and sentinel lymphadenectomy for early stage melanoma: a multicenter trial. Multicenter Selective Lymphadenectomy Trial Group. *Ann Surg* 1999;230:453-63.

Assessing drug safety

EDITOR—Eaton reports that more surveillance of drugs is needed to protect the public.¹ All drugs are dangerous, some may also be useful. If drug efficacy is easy to show, drug safety (low risk) is essentially unprovable: absence of demonstration of risk is not demonstration of absence. Assessment of risk and benefit is a matter of health politics and opinion: is it better to die of a heart attack or of colon cancer? But the patient's opinion is rarely sought. Most adverse reactions are related to the known pharmacological properties of older drugs and their interactions.² Doctors need to be taught pharmacology and the effects of drugs and be trained in taking drug histories,³ and patients trained in proper and safe drug use. Managing drug induced diseases is a clinical specialty.⁴

Drug safety or pharmacovigilance ("watchfulness in guarding against danger from drugs or providing for safety of drugs") concerns can be generated by preclinical, premarketing, or post-marketing data, from similar products or previous disease or background related issues.

Premarketing safety data are probably about as good as they'll get. Clinical trials cannot ensure drug safety in practice. Demonstrating risk equivalence in a clinical trial requires 100-250 events per study arm,⁴ which can induce uselessly large trials. Bigger premarketing studies increase the cost of drug development without increasing safety. Clinical trials are not real life. Post-marketing studies cannot be done before a drug is marketed.

Real risks need to be discovered and assessed in real use. Patients' and prescribers' imagination and risk in drug use and misuse is infinite. Post-marketing drug risk assessment relies on multiple simultaneous methods:

Spontaneous reporting ("yellow cards") is an alert system. It is designed to generate hypotheses on rare unexpected events. It is useless, counterproductive, and a sign of ignorance to expect it to quantify known signals or common reactions. These will be assessed by using a combination of pharmacoepidemiology, ad hoc studies, and resources such as, in the United Kingdom, the Drug Safety Research Unit in Southampton, the General Practice Research Database in London, the Medicine Monitoring Unit in Dundee, or similar resources worldwide. These are a crucial part of the rapid exploration of emerging signals and should be encouraged, developed and funded.

Politicians, regulators, academics, journalists, journal editors, prescribers, and patients need to be trained to understand pharmacovigilance, drug safety signals, assessment of risks and benefits, and risk management.

Nicholas Moore *professor of clinical pharmacology*
INSERM U657, Université Victor Segalen,
Bordeaux, France
nicholas.moore@pharmaco.u-bordeaux2.fr

Competing interests: NM has worked in drug safety for over 20 years and is convinced he is right, most of the time. The initial version of this paper was over 2000 words long. Shrinking it may have made it more syllable.

- 1 Eaton L. More surveillance of drugs is needed to protect public. *BMJ* 2004;329:1124-b. (13 November.)
- 2 Pirmohamed M, James S, Meakin S, Green C, Scott AK, Walley TJ, et al. Adverse drug reactions as cause of admission to hospital: prospective analysis of 18 820 patients. *BMJ* 2004;329:15-9.
- 3 Moore N, Masson H, Noblet C, Joannides R. What medicines do patients really take? A comparison of free form vs oriented questionnaires. *Post Marketing Surveill* 1993;7:355-62.
- 4 Moore N. The role of the clinical pharmacologist in the management of adverse drug reactions. *Drug Saf* 2001;24:1-7.
- 5 Moore N, Tubert-Bitter P, Fourrier A, Begaud B. A simple method to estimate sample sizes for safety equivalence studies using inverse sampling. *J Clin Epidemiol* 2003;56:433-5.

Europe also needs agency for postmarketing surveillance

EDITOR—With respect to the withdrawal of rofecoxib (Vioxx), several months of debate on drug safety have focused on US policy, but very little has been said about the European situation.¹ Some fundamental ques-

tions remain to be answered. For example, why did the manufacturer, and not the regulatory agencies, withdraw rofecoxib?

Both the US Food and Drug Administration (FDA) and the European Medicines Agency (EMA) had concluded, before this decision, that this product—and the other drugs belonging to the cyclo-oxygenase 2 inhibitor class—were essentially safe, and all that was needed was a few changes to the summary of the product characteristics, a document that is seldom read by prescribing doctors. This had happened before, for cerivastatin and other drugs, raising doubts about the regulatory agencies' efficiency in reaching decisions about benefit-risk ratios.

The American government has already decided to create an agency separate from the FDA, the Drug Safety Oversight Board, to monitor the toxicity of drugs after they have entered the market. In Europe, the European Commission has not reacted, probably because the EMA reports to the Directorate General for Enterprise and Industry.

Several reasons exist to support the establishment of national and European agencies, not only to keep an eye on spontaneous reports from prescribers but also to take an active part in searching for drug induced toxicity. In Europe this agency should report to the Directorate General for Health and Consumer Protection (SANCO) and should be independent from the EMA.

It is essential to establish a balance of power between approvals and withdrawals of drugs or restrictions of therapeutic indications to provide adequate information to doctors and the public and help counteract the pharmaceutical companies' monopoly of information.

Silvio Garattini *director*
Istituto di Ricerche Farmacologiche "Mario Negri,"
Via Eritrea, 62, I-20157 Milan
garattini@marionegri.it

Competing interests: None declared.

- 1 Lenzer J. FDA is incapable of protecting US "against another Vioxx." *BMJ* 2004;329:1253. (27 November.)

Even pharmaceutical companies are vulnerable to loss of faith

EDITOR—The scientific process depends on trust: we assume that authors acknowledge influences such as the involvement of pharmaceutical companies. In medicine, industry sways authors by funding research and by setting the norms for career advancement. Refereed publications are essential for promotion, so if industry can control the journals it has power over the medical profession. Increasingly, companies subcontract papers to apparently respectable experts. Such ghostwriting should have no place in journals claiming to report primary scientific results.

Refereed journals set the benchmark for scientific excellence, but this gold standard is becoming tarnished. Editors admit that

companies exert pressure, even on prestigious journals.¹ Ghostwritten drug studies, little more than marketing propaganda, are published by leading journals. Pharmaceutical companies also protect their markets by promoting biased studies of antioxidants, such as vitamins C and E, and other forms of preventive medicine that could compete with drug sales.²

Drug companies treat scientific publication as a branch of marketing. If a study in a prestigious journal will guarantee extra sales, they allocate millions of pounds to recruit and control the experiment. Researchers who claim not to be influenced by such apparent generosity are deluding themselves. This escalation of deceit in the medical journals threatens genuine scientific progress.

It is becoming hard to differentiate good studies from bad, real science from marketing. Knowing that drug companies need sick people in order to generate profits, many people no longer trust medical science. If the medical endeavour is seen as unreliable, then everyone will suffer. Even the pharmaceutical companies are vulnerable to such a loss of faith: unless they clean up their act, they will kill the goose that lays the golden egg.

Steve Hickey *honorary fellow*
Department of Biological Sciences, John Dalton
Building, Manchester Metropolitan University,
Manchester M1 5GD
radicalascorbate@yahoo.com

Competing interests: None declared.

- 1 Eaton L. Editor claims drug companies have a "parasitic" relationship with journals. *BMJ* 2005;330:9. (1 January.)
- 2 Hickey S, Roberts H. *Ascorbate: the science of vitamin C*. Napa, CA: Lulu Press, 2004.

Funding for GMC should come from taxpayers

EDITOR—"We want to put an end to the idea that the GMC [General Medical Council] is a representative body for doctors. It is not. Its primary role must be to protect patients," said the health secretary, John Reid.¹

Surely, if this is the duty of the GMC, and it does not purport to represent doctors, its costs should not be borne by the doctors but by the taxpayer. A licensing and registration body may be legitimately funded by doctors, but it is unreasonable to expect doctors to fund the extremely high costs of protecting patients from possible criminal elements.

An organisation whose primary function is to protect patients and citizens should be funded in the same way as the police and report its findings to the courts and the licensing body.

Brian J Penney *general practitioner*
Smith Falls, Ontario, Canada K7A 2Y8
BPenney1@cogeco.ca

Competing interests: BP pays GMC dues.

- 1 Dyer C. GMC must protect patients says government minister. *BMJ* 2005;330:271. (5 February.)