Published in final edited form as:

Nat Neurosci. 2016 November ; 19(11): 1418–1425. doi:10.1038/nn.4413.

Using model systems to understand errant plasticity mechanisms in psychiatric disorders

Bruno B. Averbeck¹ and **Matthew V. Chafee^{2,3}**

¹Laboratory of Neuropsychology, National Institute of Mental Health, National Institutes of Health, Bethesda, MD, 20892-4415, USA

²Department of Neuroscience, University of Minnesota Minneapolis, MN 55455, USA

³Brain Sciences Center, VA Medical Center, Minneapolis, MN 55417, USA

Abstract

In vivo model systems are a critical tool for gaining insight into the pathology underlying psychiatric disorders. Modern functional imaging tools allow study of brain correlates of behavior in clinical groups and genome-wide association studies are beginning to uncover the complex genetic architecture of psychiatric disorders. Several psychiatric disorders derive from pathological neural plasticity, and studying the mechanisms that underlie these processes, include reinforcement learning and spike-timing dependent plasticity, requires the use of animals. It will be particularly important to understand how individual differences in plasticity mechanisms at a cellular level confer resiliency on some but lead to disease in others.

Understanding the neurobiology of psychiatric disorders requires the use of animals. Functional magnetic resonance imaging (fMRI) in patient groups has opened a window into the human brain that provides insight into the networks that may be disrupted in various disorders and positron-emission tomography (PET) can show differences in neurotransmitter systems between patient groups and comparison participants¹. However, this work only suggests gross anatomical correlates of disorders and provides little insight into the neural mechanisms that are disrupted in patient groups. Some psychiatric disorders are likely due to disordered learning caused by pathology in cellular plasticity mechanisms. Insight into disordered plasticity mechanisms at the cellular and synaptic levels requires experiments in model systems.

While it is clear that animals are necessary to gain an understanding of the mechanistic underpinnings of psychiatric disorders, model systems can be used in many ways². One approach to using animals in psychiatric research, which has failed to yield progress, is to develop an animal model of the disease using a behavior thought to be relevant to the disease in conjunction with a manipulation that affects this behavior. The model is then used as a

Correspondence to: Bruno B. Averbeck, Ph.D. Laboratory of Neuropsychology, NIMH/NIH, Building 49 Room 1B80, 49 Convent Drive MSC 4415, Bethesda, MD 20892-4415, bruno.averbeck@nih.gov, Office: 1 (301) 594-1126.

Author contributions: BBA and MVC wrote the paper.

Financial disclosures: None.

screen in drug discovery by showing that a drug can ameliorate a behavioral deficit in the model^{3, 4}. In this approach the model's translational utility is assessed with face, construct and predictive validity⁵. Face validity refers to similarity between behavioral or neural features of a disease and the animal model. As there are no well-validated biomarkers of psychiatric disorders⁵ behaviors are usually used to achieve face validity in animal models. However, reducing psychiatric disorders to a single or even a few behaviors is problematic for multiple reasons, including heterogeneity among patients with a single diagnosis.

Construct validity refers to the technique used to generate the animal model. If the neuropathological process that gives rise to the disease was understood, and that process could be recapitulated in an animal model, this would provide strong construct validity. The advent of transgenic rodent models suggested an approach to generating a model system with good construct validity⁶. Specifically, if a candidate gene existed for a psychiatric disorder, the disease causing allele could be knocked into a rodent. However, it is now clear that psychiatric disorders result from both a large number of common alleles with low penetrance and a much smaller number of very rare alleles with high penetrance^{7, 8}. In schizophrenia, each mechanism contributes similarly to disease liability in the population⁹ and therefore any single gene accounts for a limited fraction of disease cases. Studying the highly penetrant, low frequency alleles¹⁰ may provide insight into general disease mechanisms. However, expression of a gene that leads to disease in the human brain often recapitulates only a subset of disease features in animal models¹¹, therapies that ameliorate deficits in animal models often fail to translate to humans¹² and genes that have clear phenotypes in animal models³ often have weak penetrance in human populations.

Predictive validity is the ability of the model to predict whether a new treatment will prove effective in the clinical setting. This is often assessed by determining whether current state-of-the-art treatments are effective in the animal model, although effective is often relative to a behavior that itself lacks sufficient validity. Overall, the validity framework appeared plausible but it has been ineffective: the clinical efficacy of antipsychotic medications has not improved over the past 50 years of drug development^{13, 14}.

Identifying mechanisms in model systems

Animal models can also be used to study the neural mechanisms that underlie behavior or behavioral dimensions^{3, 15, 16}. A detailed understanding of pathology in neural mechanisms relevant to psychiatric disorders can give rise to treatments. Going from an understanding of mechanism to treatment is a difficult step in itself, because tools currently available to treat psychiatric disorders are blunt instruments. For example, even in disorders caused by single gene mutations like Huntington's disease⁸, discovery of the gene has led to increased understanding of the underlying pathology but not improvements in treatment. In addition, for this approach to work there needs to be a close collaboration between clinicians and basic scientists, so that highly similar behaviors that are mediated by very similar neural circuits and dynamics within those circuits are studied across species.

Understanding mechanism can, however, lead to improvements in clinical practice. For example, the use of deep-brain-stimulation (DBS) to treat Parkinson's disease was

developed from an understanding of how the basal ganglia drives movements^{17, 18}. Specifically, increased activity in the excitatory subthalamic nucleus (STN) drives increased activity in the inhibitory globus-pallidus internal segment (GPi), and this leads to decreased activity in thalamo-cortical circuits¹⁹ which drive movements. Although current understanding of this system is more complex²⁰, this simple model does predict the effects of manipulations of this circuit on behavior²¹. Lesions of the STN were shown to ameliorate motor deficits in the macaque MPTP model of Parkinsonism²². Subsequent work in patients examined stimulation of either GPi or the STN and found that both could alleviate symptoms²³. While DBS is clinically effective, its specific mechanism of action is still not clear²⁴. Although this example shows that understanding mechanism can lead to improvements in treatment, the pathophysiology that underlies Parkinson's disease has been understood since the beginning of the 20th century²⁵. In psychiatric disorders we still lack understanding of the pathophysiology²⁶.

Neuropsychiatric diseases as pathological learning

Understanding pathogenesis in psychiatric disorders is difficult because it requires establishing causal relationships between events that take place at different levels of scale in the brain. Fully understanding the causal chain of events leading to pathology requires understanding how genetic mutations^{7, 27} produce changes in development^{28–30} that interact with the environment³¹ to alter the physiology of neurons and synapses³² changing the patterns of activity generated by networks^{33–35} leading to distortions of perception and cognition^{36, 37}. We know most about disease at the two extremes of this sequence. What is lacking is a mechanistic understanding of events at intervening levels of scale, where pathogenesis proceeds by changing how neurons and synapses work.

Evidence at genetic and behavioral levels suggests that psychiatric disorders may be disorders of learning. For example, many of the genetic loci associated with schizophrenia are related to synaptic plasticity⁷. In addition, at a behavioral level, drug addiction follows from a pathological association of drug cues and motivational systems and post-traumatic stress disorder develops when the fear and anxiety driven by a traumatic event does not extinguish. Furthermore, across several disorders, treatments currently being put forward from a mechanistic point of view target NMDA receptors, and these receptors are closely related to synaptic plasticity. What is missing, however, is an understanding of the pathology in cellular mechanisms that link genetic predispositions, when they exist, to the deficits in learning at a behavioral level. Disorders in dopamine driven reinforcement learning³⁸ and spiketiming dependent plasticity³⁹ may underlie pathology in, for example, addiction and schizophrenia. Therefore, close links between genetics, plasticity at a cellular level driven by one of these learning mechanisms, and behavior, need to be worked out. Below we consider where studying learning and plasticity in model systems has led to suggestions for novel therapeutics and also consider areas where additional research on these mechanisms is needed. Particularly, understanding how individual differences in plasticity mechanisms confer resiliency on some and lead to disease in others is just beginning in addiction, but has been little explored in other disorders.

Relevance of fear learning and extinction to anxiety disorders

Fear conditioning or more specifically threat conditioning⁴⁰ and its extinction is one of the better understood behaviors with relevance to psychiatric disorders in systems neuroscience. In the fear conditioning model, a neutral conditioned stimulus (CS) is paired with an unconditioned stimulus (US), often a foot shock⁴¹. Following a few pairings, the animal forms an association between the CS and the US, such that when the animal is brought back in a subsequent session, presentation of the CS leads to freezing. Considerable work has shown that circuits within the amygdala play a critical role in the association of the CS and the US^{42–44}. For example, before the association between the CS and the US, cells in the lateral nucleus (LA) of the amygdala respond only weakly to the CS. However, after the association, the CS drives a robust response in the LA neurons^{45, 46}. Furthermore, the increased response to the CS is largely driven by plasticity in the LA, rather than being inherited from other structures^{47, 48}. The LA projects directly to the central nucleus (CN) of the amygdala, as well as to the basal nucleus (BN) which also projects to the CN⁴⁹. Freezing, following presentation of the CS, is driven by descending outputs from the CN of the amygdala to the peri-aqueductal gray.

Fear extinction is the process of extinguishing the behavioral response to the CS. In the extinction paradigm, once a fear association has been formed, it is extinguished by exposing the animal to the CS in the absence of the US. Following a sufficient number of trials, the freezing response to the CS disappears. While extinction learning has often been thought of as unlearning or erasing the original fear association⁵⁰, it is now accepted that most extinction learning (but see⁵¹, and below) is the learning of a new inhibitory association between the CS and no-US, that competes with the original CS-US association^{52, 53}. This is based on several findings in extinction experiments. Specifically, under renewal, when the CS is extinguished outside of the original context, a return to the original context leads to a return of freezing. Under reinstatement, if the US is delivered by itself following extinction, the CS will subsequently drive freezing. Under recovery, the simple passage of time leads to the reemergence of the fear memory. In addition, after extinction, re-association of the CS and US in a subsequent conditioning session occurs more quickly, a process referred to as savings. Thus all of these conditions show that the original CS-US association was still present in some form following extinction training.

This work has also been translated to human subjects and clinical studies of anxiety disorders^{54–58}. Using fMRI in healthy human participants, it was shown that the neural circuitry underlying fear learning and extinction is shared across humans and rodents⁵⁹. This work has not, however, translated into a useful biomarker for anxiety. In addition, fear conditioning and extinction in animal models differ from the more complex clinical picture of anxiety disorders. For example, participants with diagnosed PTSD that resulted from a single discrete trauma are hyper-responsive to acoustic startle, consistent with the fear conditioning model, whereas patients with PTSD that resulted from cumulative trauma have decreased response to acoustic startle⁶⁰. Discrete trauma PTSD patients also respond more readily to exposure therapy than cumulative trauma patients⁶¹. In addition to this clinical distinction, fear is an emotional state that can only be assessed in human participants, and fear as an emotion differs from physiological responses to threat assessed in animals⁴⁰.

Meta-analyses suggest that patients with anxiety disorders have a modest increase in acquisition of fear associations, particularly a generalization to safety cues (i.e. CSs not associated with aversive USs) as well as a modest increase in expression of fear during extinction⁵⁴. Similar to the functional imaging data, however, behavioral studies of fear learning are not diagnostic of anxiety disorders because there is large overlap in behavior across groups. In addition, and perhaps most importantly, individual differences in synaptic plasticity at a cellular level that may underlie resilience in some but result in anxiety disorders in others are not known.

Exposure therapy, derived from fear extinction, is an important component of cognitive behavioral therapy in anxiety disorders^{57, 62}. Similar to extinction in rodents, extinction in anxiety disorders leads to temporary relief from symptoms, but often fear returns after the passage of time due to renewal or recovery⁶³. This has led to a search for ways to improve exposure therapy by enhancing learning. Two ideas for improvement have come from basic science studies of extinction learning⁶⁴. Specifically, the use of D-cycloserine^{65, 66}, and reconsolidation-extinction, to enhance extinction learning^{67, 68}.

Early experiments showed that blocking NMDA receptors in the amygdala, which interferes with plasticity mechanisms decreased extinction learning in rodents⁶⁹. It was subsequently shown that D-cycloserine, a non-competitive NMDA agonist, administered systemically or within the amygdala, could enhance extinction⁷⁰. Follow-up studies showed that D-cycloserine administered to human participants with acrophobia in combination with exposure therapy led to increased extinction learning⁷¹. Subsequent results have been mixed and current meta-analyses suggest that results are inconclusive but additional studies should be done to clarify effects and mechanism⁷². Better understanding of the cellular mechanisms that underlie extinction may lead to better treatments to improve extinction learning.

The second approach that has emerged from basic research to increase the effectiveness of exposure therapy is the reconsolidation-extinction paradigm. As discussed above, standard extinction protocols do not lead to erasure of fear memory. Rather they lead to the creation of a new CS-no US memory that inhibits the original CS-US association. However, when memories are retrieved they become labile and must be reconsolidated⁷³, a process which requires protein synthesis. This was first exploited in an extinction paradigm in which, during extinction, rats were given a single CS with either anisomycin, which blocks protein synthesis, or ACSF as a control. Freezing to the single CS did not differ by group. However, on the day after the CS/drug condition, presentation of the CS led to less freezing by the rats that had received anisomycin. Unfortunately, anisomycin is toxic in humans. Therefore, a behavioral procedure was developed in an attempt to achieve the same end. In this study, it was shown that if rats were given a single CS presentation, either 10 minutes or 1 hour before extinction training, but not 6 or 24 hours before extinction training, the return of fear by recovery, reinstatement or renewal was substantially decreased (Fig. 1)⁶⁷. This was taken as evidence that, following the reconsolidation-extinction paradigm, the fear memory was permanently attenuated. Subsequent studies suggested that this approach also worked in healthy human subjects, in a fear-learning paradigm⁶⁸. Considerable additional work has followed this finding^{51, 74}. Current meta-analyses suggest that reconsolidation extinction reduces the return of fear in healthy human participants but not in rodents⁷⁵. Efficacy of this

procedure has not yet, however, been examined in clinical groups. These results suggest that bottom up work in basic science focused on understanding the neural mechanisms that underlie a relevant behavior can drive treatments based on novel mechanisms.

Neural circuits underlying compulsive drug taking

The neural circuitry that underlies addiction has considerable overlap with the circuitry that underlies fear learning⁷⁶. Addiction can be considered a pathological association between cues and appetitive USs (i.e. drugs of abuse) just like fear is a pathological association between cues and aversive USs. The amygdala plays an important role in addiction⁷⁷. However, the dopamine system and the striatum are also heavily implicated. Many drugs of abuse either directly or indirectly increase dopamine signaling^{78, 79}. There is considerable evidence that dopamine codes reward prediction errors (i.e. the difference between the received and expected reward)⁸⁰, and that reward prediction errors are an important component of reinforcement learning, or associating stimuli with prediction of reward⁸¹. This theory predicts that when outcomes are more rewarding than expected dopamine is released and the estimate of reward associated with cues, actions or the current context are increased. In explicit versions of this model the dopamine signal drives synaptic plasticity on cortical-striatal synapses and it is these synapses that encode expected reward⁸². In addition, temporal-difference learning models predict that the first cue that predicts reward drives dopamine release⁸³. In a simple appetitive Pavlovian experiment, when a CS is first shown, it will not lead to dopamine release because the CS does not predict reward. If a reward (US) is then delivered following the CS, there will be a dopamine response to the reward. However, following learning, when the CS comes to predict the reward, the dopamine response will shift to the CS because it is the first cue that predicts reward. When the reward is subsequently delivered there will be no dopamine response because the reward has been predicted.

Drugs may drive pathological levels of learning because they drive dopamine release and therefore a constant reward prediction error⁸⁴. Under normal physiological conditions, after learning, the reward prediction would equal the received reward (for example, the sight of a food pellet is a cue that predicts the actual reward obtained when the food pellet is consumed) and the reward prediction error would be zero. With no reward prediction error there would be no dopamine release and no additional learning. However, with drugs of abuse that cause an artificial increase in dopamine levels, there would always be a positive reward prediction error, and therefore learning. This would cause value estimates to be driven to pathological levels. Although several general aspects of this model have been well-supported by data, some specific predictions of this model have not been supported by subsequent experiments⁸⁵.

This model has been used to account for dopamine responses in addiction. PET work has shown that in addicts, there is a large dopamine response in the striatum to drug cues, which are CSs⁸⁶. However, there is decreased dopamine response to the actual delivery of the drug, which is the US. In addition, it has been shown that the dopamine released to the drug cues causes craving in addicts⁸⁷.

In addition to dopamine, amygdala dependent processes have also been implicated in addiction^{77, 88}. Studies have shown that the amygdala underlies appetitive learning^{89, 90}, in addition to fear learning^{42, 44}. Recent circuit analyses suggest that it may be the basal-lateral amygdala projections to the ventral-striatum that are important for appetitive learning⁹⁰⁻⁹², which argues for an interpretation of amygdala contributions to addiction within the dopamine reward prediction error framework. Specifically, this would predict that dopamine is modifying amygdala-ventral striatal synapses, instead of cortical-striatal synapses. However, there is also work that shows plasticity specifically within the amygdala during appetitive learning^{90, 93}.

Drug self-administration, in which animals learn to press a lever to receive drug, has often been used to mechanistically understand drug taking that, in humans, may lead to addiction⁹⁴. However, selfadministration does not account for drug seeking or craving. More recent behavioral paradigms have required animals to complete a random interval schedule on a seeking lever before being presented with a different, taking lever, that delivers drugs on a fixed response schedule^{95, 96}. In variants of this task, rats are extensively trained using the chained seeking/taking schedules. In subsequent sessions the rats are probabilistically punished in some trials after completing the seeking schedule, instead of being presented with the taking lever. A subset of rats continues to respond compulsively on the seeking lever, obtaining the taking lever in some trials and punishment in others. A complimentary approach has used second order conditioned reinforcement schedules^{72, 73}. In these procedures rats are reinforced only with drug associated cues, which are able to maintain responding over long intervals. Both of these procedures better capture the compulsive seeking aspect of drug abuse than selfadministration, although they have not yet directly led to changes in treatment. In addition, there is some work on individual differences in rats that predispose animals to compulsive drug taking⁹⁷. And a few recent studies have examined differences in plasticity mechanisms that may underlie behavioral differences^{98, 99}.

Additional work on individual differences in plasticity may better reveal treatment targets, or at least lead to a better understanding of why some individuals becomes addicts while others do not, given the same drug exposure.

Two learning manipulations originally proposed to affect aversive learning, driven by studies in the amygdala, have also been explored in appetitive learning. Specifically, both retrieval-extinction behavioral procedures and pharmacological manipulations of reconsolidation have been examined. Similar to the effects previously explored in fear extinction, a retrieval-extinction procedure has been developed to extinguish the appetitive associations tied to drug cues in addicts¹⁰⁰. This study found in animals, that presenting appetitive cues 10 minutes or one hour before a subsequent extinction session led to decreased renewal, reinstatement (Fig. 2) and spontaneous recovery of conditioned place preference. Similarly in human drug addicts, extinction training of drug cues following retrieval led to less spontaneous recovery up to 184 days after extinction training. This suggests that retrieval-extinction training can be effective in appetitive learning as well. The appetitive extinction effects have also been found to be significant in a meta-analysis in rodents⁷⁵.

In addition to these behavioral procedures, post-retrieval pharmacological manipulations have also been used to alter learning. In these experiments, rats are conditioned to CSs

paired with drugs. Following the conditioning, they are given an extinction session that serves to recall the CS memory. Either just before or just after the extinction session, they are given various drugs including NMDA antagonists or beta-adrenergic antagonists, as both have been shown to interfere with reconsolidation. They are subsequently tested behaviorally. Meta-analyses have shown that both NMDA antagonists and beta-adrenergic antagonists can affect reconsolidation and enhance extinction¹⁰¹.

Model systems for understanding cortical network dysfunction in schizophrenia

Schizophrenia is a more complex disorder to study than anxiety or addiction. In anxiety there are better developed models of the circuitry that gives rise to defensive or avoidance behaviors and in addiction there is a reasonable understanding of the pathogenesis. The gap between understanding and disease is larger in schizophrenia. Nonetheless substantial advances have been made in understanding the neurobiology of the disease. Functional imaging^{34, 102}, magnetoencephalography¹⁰³ and EEG¹⁰⁴ have shown changes in brain activity and synchrony patterns^{105, 106} in patients. These studies provide important insight into pathophysiological signatures of schizophrenia. However, they cannot resolve brain activity in patients at the level of action potentials in neurons. That is a significant limitation because accumulating evidence suggests that one aspect of disease pathogenesis in schizophrenia may involve the molecular mechanisms of activity-dependent synaptic plasticity¹⁰⁷. Specifically, some risk mutations in schizophrenia involve a cluster of genes that play a role in NMDA receptor (NMDAR) mediated synaptic transmission¹⁰⁸. NMDA receptors play a crucial role in triggering activity-dependent synaptic plasticity¹⁰⁹, and the plasticity mechanism mediated by NMDARs is exquisitely sensitive to the precise timing of action potentials in pre- and postsynaptic neurons³⁹. Additional risk mutations involve genes that play a role in the immune system¹¹⁰, and new evidence suggests these genes support molecular signaling pathways that target synapses for elimination during development, a process which itself is activity-dependent¹¹¹. This suggests that schizophrenia pathogenesis could involve a disturbance in the linkage between activity patterns at the level of spiking neurons and the strength of synaptic connections in cortical networks. Support for this class of theories could significantly redirect strategies to identify novel treatments. However, because pathogenic mechanisms of this type depend crucially on the timing of action potentials in synaptically coupled neurons and the disruption of this timing in the disease state, testing them requires resolving electrical activity at the cellular level, which requires the use of animals.

To effectively model the interaction between activity patterns and synaptic plasticity mechanisms, it is necessary to first try to replicate pathophysiological activity patterns likely to occur in the human disease in the model system. This will be more likely if behavior is well matched between patients with schizophrenia and the model system. Consequently, the two species should perform the same behavioral task. In addition, a manipulation should be identified that causes the emergence of the same behavioral error pattern in the animal model that is evident in patients. Second, because neural activity patterns are constrained by patterns of anatomical connectivity in cortical networks, it is optimal if the neural system

mediating performance in the animal is as similar in organization as possible to the corresponding network in the human brain. As behavioral and neurophysiological deficits in schizophrenia often preferentially involve the prefrontal cortex^{112, 113}, this places particular weight on the necessity to optimize homology of prefrontal networks across species, and this criterion favors primate over rodent models^{114, 115}.

This raises the question of which behavioral features of schizophrenia provide the most promising targets for successful translation to animals. The clinical symptoms of schizophrenia, which include hallucinations, delusions, and flatness of affect¹¹⁶ present obvious challenges for translation to animals (not least of which is that these symptoms are typically evaluated clinically by talking to patients). However, patients with schizophrenia also exhibit deficits in cognitive functions that are measurable using automated behavioral tasks that can be translated to animal models. Understanding the neural basis of cognitive deficits rather than clinical symptoms of schizophrenia has therefore been the focus of translational research^{31, 117}. Several multi-investigator consortia have identified behavioral paradigms that most reliably measure specific cognitive deficits reflecting genetic risk in schizophrenia: including deficits in executive control, working memory, attention, and sensory gating^{118–121}. As with the other psychiatric disorders, reducing a disease as complex as schizophrenia to one or a few cognitive deficits in an animal model will likely lead to a model with limited validity and therefore results from these studies should be interpreted carefully.

Patients with schizophrenia, and to a lesser degree their first degree relatives, exhibit deficits in tasks that require spatial working memory¹²². A deficit in spatial working memory can be demonstrated in patients performing the oculomotor delayed response (ODR task)¹²³. In this task, subjects direct a saccadic eye movement to the location of a peripheral visual cue several seconds after it disappears. Working memory for the location of the cue is associated with the persistent activation of spatially-selective neurons in the dorsolateral prefrontal cortex in monkeys^{124, 125}. Dorsolateral prefrontal lesions¹²⁶, inactivation¹²⁷, or injection of D1 dopamine receptor antagonists into monkey prefrontal cortex¹²⁸ induce spatially restricted deficits in working memory. Importantly, performance on the ODR task is preserved after insult to prefrontal cortex if the target remains visible through the delay, providing a visual target for the saccade at the time it is executed. This establishes the specificity of the deficit for working memory. The ODR task was successfully back-translated from monkeys to patients with schizophrenia and first-degree relatives to demonstrate a specific impairment in spatial working memory¹²⁹. It has been repeatedly documented using functional imaging in patients that reduced performance on the ODR and other working memory tasks is associated with reduced activation of prefrontal cortex^{130, 131} (Fig. 3A).

Most efforts to model behavioral and neurophysiological features of schizophrenia in nonhuman primates have been based on administration of noncompetitive NMDA receptor (NMDAR) antagonists such as ketamine or phencyclidine¹³². Blocking NMDAR in human control subjects replicates positive and negative symptoms as well as some cognitive deficits of schizophrenia¹³³. In monkeys, local iontophoretic application of NMDAR antagonists disrupts the persistent activation of prefrontal neurons associated with spatial working

memory¹³⁴ (Fig. 3B). Other studies have shown that systemic administration of ketamine to monkeys induces errors in a rule-based antisaccade task similar to those produced by patients, and neural recording has characterized concomitant reduction in the encoding of task-related information by prefrontal neurons^{135–137}. Collectively these studies provide important insight into how prefrontal circuits computationally fail in response to loss of NMDAR function. However, some robust findings in patients are not replicated by the NMDAR model¹³⁸.

The AX-CPT (continuous performance task) is a task that measures specific deficits in executive control in patients with schizophrenia^{34, 139–141} and has been successfully translated to monkeys^{142, 143}. In this task, a cue stimulus modifies the response required to a subsequent probe. On some trials, the cue stored in working memory countermands a habitual or prepotent response to the probe, and it is specifically on this subset of trials that patients with schizophrenia commit the most errors (Fig. 3C)^{119, 139, 140}. Administration of ketamine to monkeys performing the AX-CPT induces the same error pattern (Fig. 3D)¹⁴². This is one example of a strong behavioral match between the error pattern of patients and an animal model.

However, the fact that a drug in monkeys is able to mimic the symptoms of a disease in humans is not evidence that the drug and disease work through the same neural mechanism. Testing causal theories at the cell and circuit levels requires neural recordings to evaluate the impact of mutations on neural function. A *Df(16)A* +/- mouse¹⁴⁴ has been engineered to mimic the 22q11.2 chromosomal microdeletion (DiGeorge syndrome) that confers a 30-fold increase in risk of developing schizophrenia and is among the strongest genetic risks for the disease¹⁴⁴. This mouse exhibits behavioral deficits similar to those seen in schizophrenia, including reduced sensory gating¹⁴⁵ and impaired spatial working memory^{145, 146}, as well as reduced gamma band synchrony between prefrontal cortex and hippocampus¹⁴⁶. Deletion of a single gene (*dgcr8*) in this region that is involved in microRNA processing confers much of the phenotype, and has been shown to produce deficits in short-term synaptic plasticity in slice recordings¹⁴⁷. MicroRNA are involved in the post-transcriptional regulation of other genes. Genome-wide association studies indicate that the large majority of SNPs increasing risk of schizophrenia, as with other psychiatric disorders, do not occur in coding regions of genes⁸, suggesting protein structure may be largely normal in the disease. Both observations point to the possibility that a defect in gene regulation could trigger pathogenesis in schizophrenia, potentially via activity-dependent gene regulatory pathways.

Conclusion

It has proven difficult to use animal models to screen for new drugs with therapeutic potential. While models have been generated that satisfy face, construct and predictive validity, they have not facilitated drug discovery. Animals are, however, the only available tool for mechanistic studies of the neural circuitry that drives behavior. More importantly, animals provide the only tool available to study learning driven neural plasticity at a cellular level, and pathology in these plastic processes likely underlies many psychiatric disorders. Understanding individual differences in plasticity mechanisms may further lead to insights into why given similar genetic backgrounds or behavioral experiences, some individuals

develop disorders while other are resilient. This understanding may then lead to better treatments.

Acknowledgments

This work was funded in part by the intramural research program of the National Institute of Mental Health.

References

- Slifstein M, et al. Deficits in prefrontal cortical and extrastriatal dopamine release in schizophrenia: a positron emission tomographic functional magnetic resonance imaging study. *JAMA Psychiatry*. 72:316–324.2015; [PubMed: 25651194]
- Insel TR. From animal models to model animals. *Biological psychiatry*. 62:1337–1339.2007; [PubMed: 18054535]
- Wong AH, Josselyn SA. Caution When Diagnosing Your Mouse With Schizophrenia: The Use and Misuse of Model Animals for Understanding Psychiatric Disorders. *Biological psychiatry*. 79:32–38.2016; [PubMed: 26058706]
- Cosgrove VE, Kelsoe JR, Suppes T. Toward a Valid Animal Model of Bipolar Disorder: How the Research Domain Criteria Help Bridge the Clinical-Basic Science Divide. *Biological psychiatry*. 79:62–70.2016; [PubMed: 26531027]
- Nestler EJ, Hyman SE. Animal models of neuropsychiatric disorders. *Nature Neuroscience*. 13:1161–1169.2010; [PubMed: 20877280]
- Campbell IL, Gold LH. Transgenic modeling of neuropsychiatric disorders. *Molecular psychiatry*. 1:105–120.1996; [PubMed: 9118320]
- Biological insights from 108 schizophrenia-associated genetic loci. *Nature*. 511:421–427.2014; [PubMed: 25056061]
- Insel TR, Collins FS. Psychiatry in the genomics era. *The American journal of psychiatry*. 160:616–620.2003; [PubMed: 12668345]
- Gratten J, Wray NR, Keller MC, Visscher PM. Large-scale genomics unveils the genetic architecture of psychiatric disorders. *Nature Neuroscience*. 17:782–790.2014; [PubMed: 24866044]
- Sullivan PF, Daly MJ, O'Donovan M. Genetic architectures of psychiatric disorders: the emerging picture and its implications. *Nat Rev Genet*. 13:537–551.2012; [PubMed: 22777127]
- Puzzo D, Gulisano W, Palmeri A, Arancio O. Rodent models for Alzheimer's disease drug discovery. *Expert Opin Drug Discov*. 10:703–711.2015; [PubMed: 25927677]
- Franco R, Cedazo-Minguez A. Successful therapies for Alzheimer's disease: why so many in animal models and none in humans? *Front Pharmacol*. 5:146.2014; [PubMed: 25009496]
- Insel TR. Rethinking schizophrenia. *Nature*. 468:187–193.2010; [PubMed: 21068826]
- Lieberman JA, Stroup TS. The NIMH-CATIE Schizophrenia Study: what did we learn? *The American journal of psychiatry*. 168:770–775.2011; [PubMed: 21813492]
- Fernando AB, Robbins TW. Animal models of neuropsychiatric disorders. *Annu Rev Clin Psychol*. 7:39–61.2011; [PubMed: 21219191]
- Pine DS, Leibenluft E. Biomarkers With a Mechanistic Focus. *JAMA Psychiatry*. 72:633–634.2015; [PubMed: 25992716]
- Albin RL, Young AB, Penney JB. The functional anatomy of basal ganglia disorders. *Trends Neurosci*. 12:366–375.1989; [PubMed: 2479133]
- DeLong MR. Primate models of movement disorders of basal ganglia origin. *Trends Neurosci*. 13:281–285.1990; [PubMed: 1695404]
- Alexander GE, DeLong MR, Strick PL. Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annu Rev Neurosci*. 9:357–381.1986; [PubMed: 3085570]
- Cui G, et al. Concurrent activation of striatal direct and indirect pathways during action initiation. *Nature*. 494:238–242.2013; [PubMed: 23354054]
- Kravitz AV, et al. Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature*. 466:622–626.2010; [PubMed: 20613723]

22. Bergman H, Wichmann T, DeLong MR. Reversal of experimental parkinsonism by lesions of the subthalamic nucleus. *Science*. 249:1436–1438.1990; [PubMed: 2402638]
23. Obeso JA, et al. Surgical treatment of Parkinson's disease. *Baillieres Clin Neurol*. 6:125–145.1997; [PubMed: 9426872]
24. Wichmann T, DeLong MR. Deep Brain Stimulation for Movement Disorders of Basal Ganglia Origin: Restoring Function or Functionality? *Neurotherapeutics*. 13:264–283.2016; [PubMed: 26956115]
25. Lees AJ. Unresolved issues relating to the shaking palsy on the celebration of James Parkinson's 250th birthday. *Movement disorders : official journal of the Movement Disorder Society*. 22(Suppl 17):S327–334.2007; [PubMed: 18175393]
26. Kapur S, Phillips AG, Insel TR. Why has it taken so long for biological psychiatry to develop clinical tests and what to do about it? *Molecular psychiatry*. 17:1174–1179.2012; [PubMed: 22869033]
27. Kirov G, et al. De novo CNV analysis implicates specific abnormalities of postsynaptic signalling complexes in the pathogenesis of schizophrenia. *Molecular psychiatry*. 17:142–153.2012; [PubMed: 22083728]
28. Marenco S, Weinberger DR. The neurodevelopmental hypothesis of schizophrenia: following a trail of evidence from cradle to grave. *Dev Psychopathol*. 12:501–527.2000; [PubMed: 11014750]
29. Owen MJ, O'Donovan MC, Thapar A, Craddock N. Neurodevelopmental hypothesis of schizophrenia. *The British journal of psychiatry : the journal of mental science*. 198:173–175.2011; [PubMed: 21357874]
30. MacDonald AW 3rd, Chafee MV. Translational and developmental perspective on N-methyl-D-aspartate synaptic deficits in schizophrenia. *Dev Psychopathol*. 18:853–876.2006; [PubMed: 17152404]
31. Burrows EL, Hannan AJ. Cognitive endophenotypes, gene-environment interactions and experience-dependent plasticity in animal models of schizophrenia. *Biological psychology*. 116:82–89.2016; [PubMed: 26687973]
32. Crabtree GW, Gogos JA. Synaptic plasticity, neural circuits, and the emerging role of altered short-term information processing in schizophrenia. *Front Synaptic Neurosci*. 6:28.2014; [PubMed: 25505409]
33. MacDonald AW 3rd, Carter CS. Event-related fMRI study of context processing in dorsolateral prefrontal cortex of patients with schizophrenia. *Journal of Abnormal Psychology*. 112:689–697.2003; [PubMed: 14674880]
34. MacDonald AW 3rd, et al. Specificity of prefrontal dysfunction and context processing deficits to schizophrenia in never-medicated patients with first-episode psychosis. *The American journal of psychiatry*. 162:475–484.2005; [PubMed: 15741464]
35. Yoon JH, et al. Association of dorsolateral prefrontal cortex dysfunction with disrupted coordinated brain activity in schizophrenia: relationship with impaired cognition, behavioral disorganization, and global function. *The American journal of psychiatry*. 165:1006–1014.2008; [PubMed: 18519527]
36. Millan MJ, et al. Cognitive dysfunction in psychiatric disorders: characteristics, causes and the quest for improved therapy. *Nat Rev Drug Discov*. 11:141–168.2012; [PubMed: 22293568]
37. Keefe RS, Harvey PD. Cognitive impairment in schizophrenia. *Handb Exp Pharmacol*. :11–37.2012 [PubMed: 23027411]
38. Dayan P, Niv Y. Reinforcement learning: the good, the bad and the ugly. *Current opinion in neurobiology*. 18:185–196.2008; [PubMed: 18708140]
39. Feldman DE. The spike-timing dependence of plasticity. *Neuron*. 75:556–571.2012; [PubMed: 22920249]
40. LeDoux JE. Coming to terms with fear. *Proceedings of the National Academy of Sciences of the United States of America*. 111:2871–2878.2014; [PubMed: 24501122]
41. Romanski LM, LeDoux JE. Equipotentiality of thalamo-amygdala and thalamo-cortico-amygdala circuits in auditory fear conditioning. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 12:4501–4509.1992; [PubMed: 1331362]
42. LeDoux JE. Emotion circuits in the brain. *Annual Review of Neuroscience*. 23:155–184.2000;

43. Herry C, Johansen JP. Encoding of fear learning and memory in distributed neuronal circuits. *Nature Neuroscience*. 17:1644–1654.2014; [PubMed: 25413091]
44. Davis, M. The role of the amygdala in conditioned and unconditioned fear and anxiety. In: Aggleton, JP, editor. *The Amygdala*. Oxford University Press; Oxford: 2000. 213–288.
45. Quirk GJ, Armony JL, LeDoux JE. Fear conditioning enhances different temporal components of tone-evoked spike trains in auditory cortex and lateral amygdala. *Neuron*. 19:613–624.1997; [PubMed: 9331352]
46. Quirk GJ, Repa C, LeDoux JE. Fear conditioning enhances short-latency auditory responses of lateral amygdala neurons: parallel recordings in the freely behaving rat. *Neuron*. 15:1029–1039.1995; [PubMed: 7576647]
47. Johansen JP, et al. Optical activation of lateral amygdala pyramidal cells instructs associative fear learning. *Proceedings of the National Academy of Sciences of the United States of America*. 107:12692–12697.2010; [PubMed: 20615999]
48. Nabavi S, et al. Engineering a memory with LTD and LTP. *Nature*. 511:348–352.2014; [PubMed: 24896183]
49. Duvarci S, Pare D. Amygdala microcircuits controlling learned fear. *Neuron*. 82:966–980.2014; [PubMed: 24908482]
50. Rescorla, RA, Wagner, AR. A theory of Pavlovian conditioning: Variations in the effectiveness of reinforcement and nonreinforcement. In: Black, AH, WF, P, editors. *Classical conditioning II: Current research and theory*. Appleton-Century-Crofts; New York: 1972. 64–99.
51. Clem RL, Schiller D. New Learning and Unlearning: Strangers or Accomplices in Threat Memory Attenuation? *Trends in Neurosciences*. 39:340–351.2016; [PubMed: 27079843]
52. Bouton ME. Context, ambiguity, and unlearning: sources of relapse after behavioral extinction. *Biological psychiatry*. 52:976–986.2002; [PubMed: 12437938]
53. Bouton ME. Context, time, and memory retrieval in the interference paradigms of Pavlovian learning. *Psychol Bull*. 114:80–99.1993; [PubMed: 8346330]
54. Duits P, et al. Updated meta-analysis of classical fear conditioning in the anxiety disorders. *Depress Anxiety*. 32:239–253.2015; [PubMed: 25703487]
55. Mineka S, Oehlberg K. The relevance of recent developments in classical conditioning to understanding the etiology and maintenance of anxiety disorders. *Acta psychologica*. 127:567–580.2008; [PubMed: 18226795]
56. Kindt M. A behavioural neuroscience perspective on the aetiology and treatment of anxiety disorders. *Behaviour research and therapy*. 62:24–36.2014; [PubMed: 25261887]
57. Graham BM, Milad MR. The study of fear extinction: implications for anxiety disorders. *The American journal of psychiatry*. 168:1255–1265.2011; [PubMed: 21865528]
58. Britton JC, Lissek S, Grillon C, Norcross MA, Pine DS. Development of anxiety: the role of threat appraisal and fear learning. *Depress Anxiety*. 28:5–17.2011; [PubMed: 20734364]
59. Delgado MR, Nearing KI, Ledoux JE, Phelps EA. Neural circuitry underlying the regulation of conditioned fear and its relation to extinction. *Neuron*. 59:829–838.2008; [PubMed: 18786365]
60. McTeague LM, Lang PJ. The anxiety spectrum and the reflex physiology of defense: from circumscribed fear to broad distress. *Depress Anxiety*. 29:264–281.2012; [PubMed: 22511362]
61. Lang PJ, McTeague LM. Discrete and recurrent traumatization in PTSD: fear vs. anxious misery. *J Clin Psychol Med Settings*. 18:207–209.2011; [PubMed: 21626350]
62. Pine, DS, Klein, RG. Anxiety disorders. In: Thapar, A, , et al., editors. *Rutter's Child and Adolescent Psychiatry*. Jon Wiley & Sons; New York: 2015. 822–840.
63. Arch JJ, Craske MG. First-line treatment: a critical appraisal of cognitive behavioral therapy developments and alternatives. *Psychiatr Clin North Am*. 32:525–547.2009; [PubMed: 19716989]
64. Quirk GJ, et al. Erasing fear memories with extinction training. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 30:14993–14997.2010; [PubMed: 21068303]
65. Rodrigues H, et al. Does D-cycloserine enhance exposure therapy for anxiety disorders in humans? A meta-analysis. *PLoS One*. 9:e93519.2014; [PubMed: 24991926]

66. Bowers ME, Ressler KJ. An Overview of Translationally Informed Treatments for Posttraumatic Stress Disorder: Animal Models of Pavlovian Fear Conditioning to Human Clinical Trials. *Biological psychiatry*. 78:E15–27.2015; [PubMed: 26238379]
67. Monfils MH, Cowansage KK, Klann E, LeDoux JE. Extinction-reconsolidation boundaries: key to persistent attenuation of fear memories. *Science*. 324:951–955.2009; [PubMed: 19342552]
68. Schiller D, et al. Preventing the return of fear in humans using reconsolidation update mechanisms. *Nature*. 463:49–53.2010; [PubMed: 20010606]
69. Falls WA, Miserendino MJ, Davis M. Extinction of fear-potentiated startle: blockade by infusion of an NMDA antagonist into the amygdala. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 12:854–863.1992; [PubMed: 1347562]
70. Walker DL, Ressler KJ, Lu KT, Davis M. Facilitation of conditioned fear extinction by systemic administration or intra-amygdala infusions of D-cycloserine as assessed with fear-potentiated startle in rats. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 22:2343–2351.2002; [PubMed: 11896173]
71. Ressler KJ, et al. Cognitive enhancers as adjuncts to psychotherapy: use of D-cycloserine in phobic individuals to facilitate extinction of fear. *Archives of general psychiatry*. 61:1136–1144.2004; [PubMed: 15520361]
72. Ori R, et al. Augmentation of cognitive and behavioural therapies (CBT) with d-cycloserine for anxiety and related disorders. *Cochrane Database Syst Rev*. 5:CD007803.2015;
73. Sara SJ. Retrieval and reconsolidation: toward a neurobiology of remembering. *Learning & memory*. 7:73–84.2000; [PubMed: 10753974]
74. Auber A, Tedesco V, Jones CE, Monfils MH, Chiamulera C. Post-retrieval extinction as reconsolidation interference: methodological issues or boundary conditions? *Psychopharmacology*. 226:631–647.2013; [PubMed: 23404065]
75. Kredlow MA, Unger LD, Otto MW. Harnessing reconsolidation to weaken fear and appetitive memories: A meta-analysis of post-retrieval extinction effects. *Psychol Bull*. 142:314–336.2016; [PubMed: 26689086]
76. Peters J, Kalivas PW, Quirk GJ. Extinction circuits for fear and addiction overlap in prefrontal cortex. *Learning & memory*. 16:279–288.2009; [PubMed: 19380710]
77. Belin D, Jonkman S, Dickinson A, Robbins TW, Everitt BJ. Parallel and interactive learning processes within the basal ganglia: relevance for the understanding of addiction. *Behavioural brain research*. 199:89–102.2009; [PubMed: 18950658]
78. Di Chiara G. Nucleus accumbens shell and core dopamine: differential role in behavior and addiction. *Behavioural brain research*. 137:75–114.2002; [PubMed: 12445717]
79. Wise RA. Dopamine and reward: the anhedonia hypothesis 30 years on. *Neurotox Res*. 14:169183.2008;
80. Schultz W. Getting formal with dopamine and reward. *Neuron*. 36:241–263.2002; [PubMed: 12383780]
81. Schultz W, Dayan P, Montague PR. A neural substrate of prediction and reward. *Science*. 275:1593–1599.1997; [PubMed: 9054347]
82. Frank MJ. Dynamic dopamine modulation in the basal ganglia: a neurocomputational account of cognitive deficits in medicated and nonmedicated Parkinsonism. *Journal of Cognitive Neuroscience*. 17:51–72.2005; [PubMed: 15701239]
83. Hollerman JR, Schultz W. Dopamine neurons report an error in the temporal prediction of reward during learning. *Nat Neurosci*. 1:304–309.1998; [PubMed: 10195164]
84. Redish AD. Addiction as a computational process gone awry. *Science*. 306:1944–1947.2004; [PubMed: 15591205]
85. Marks KR, Kearns DN, Christensen CJ, Silberberg A, Weiss SJ. Learning that a cocaine reward is smaller than expected: A test of Redish's computational model of addiction. *Behavioural brain research*. 212:204–207.2010; [PubMed: 20381539]
86. Volkow ND, Fowler JS, Wang GJ, Baler R, Telang F. Imaging dopamine's role in drug abuse and addiction. *Neuropharmacology*. 56(Suppl 1):3–8.2009; [PubMed: 18617195]
87. Volkow ND, Wang GJ, Fowler JS, Tomasi D, Baler R. Food and drug reward: overlapping circuits in human obesity and addiction. *Curr Top Behav Neurosci*. 11:1–24.2012; [PubMed: 22016109]

88. Belin D, Belin-Rauscent A, Murray JE, Everitt BJ. Addiction: failure of control over maladaptive incentive habits. *Current opinion in neurobiology*. 23:564–572.2013; [PubMed: 23452942]
89. Janak PH, Tye KM. From circuits to behaviour in the amygdala. *Nature*. 517:284–292.2015; [PubMed: 25592533]
90. Namburi P, et al. A circuit mechanism for differentiating positive and negative associations. *Nature*. 520:675–678.2015; [PubMed: 25925480]
91. Stuber GD, et al. Excitatory transmission from the amygdala to nucleus accumbens facilitates reward seeking. *Nature*. 475:377–380.2011; [PubMed: 21716290]
92. Britt JP, et al. Synaptic and behavioral profile of multiple glutamatergic inputs to the nucleus accumbens. *Neuron*. 76:790–803.2012; [PubMed: 23177963]
93. Tye KM, Stuber GD, de Ridder B, Bonci A, Janak PH. Rapid strengthening of thalamo-amygdala synapses mediates cue-reward learning. *Nature*. 453:1253–1257.2008; [PubMed: 18469802]
94. White NM. Addictive drugs as reinforcers: multiple partial actions on memory systems. *Addiction*. 91:921–949.1996; [PubMed: 8688822]
95. Belin-Rauscent A, Fouyssac M, Bonci A, Belin D. How Preclinical Models Evolved to Resemble the Diagnostic Criteria of Drug Addiction. *Biological psychiatry*. 79:39–46.2016; [PubMed: 25747744]
96. Olmstead MC, Parkinson JA, Miles FJ, Everitt BJ, Dickinson A. Cocaine-seeking by rats: regulation, reinforcement and activation. *Psychopharmacology*. 152:123–131.2000; [PubMed: 11057515]
97. Belin D, Mar AC, Dalley JW, Robbins TW, Everitt BJ. High impulsivity predicts the switch to compulsive cocaine-taking. *Science*. 320:1352–1355.2008; [PubMed: 18535246]
98. Waselus M, et al. Long-term effects of cocaine experience on neuroplasticity in the nucleus accumbens core of addiction-prone rats. *Neuroscience*. 248:571–584.2013; [PubMed: 23811073]
99. Fligel SB, et al. Genetic background and epigenetic modifications in the core of the nucleus accumbens predict addiction-like behavior in a rat model. *Proceedings of the National Academy of Sciences of the United States of America*. 113:E2861–2870.2016; [PubMed: 27114539]
100. Xue YX, et al. A memory retrieval-extinction procedure to prevent drug craving and relapse. *Science*. 336:241–245.2012; [PubMed: 22499948]
101. Das RK, Freeman TP, Kamboj SK. The effects of N-methyl D-aspartate and B-adrenergic receptor antagonists on the reconsolidation of reward memory: a meta-analysis. *Neuroscience and biobehavioral reviews*. 37:240–255.2013; [PubMed: 23261501]
102. Yoon JH, et al. Association of dorsolateral prefrontal cortex dysfunction with disrupted coordinated brain activity in schizophrenia: relationship with impaired cognition, behavioral disorganization, and global function. *Am J Psychiatry*. 165:1006–1014.2008; [PubMed: 18519527]
103. Georgopoulos AP, et al. Synchronous neural interactions assessed by magnetoencephalography: a functional biomarker for brain disorders. *J Neural Eng*. 4:349–355.2007; [PubMed: 18057502]
104. Minzenberg MJ, et al. Gamma oscillatory power is impaired during cognitive control independent of medication status in first-episode schizophrenia. *Neuropsychopharmacology*. 35:2590–2599.2010; [PubMed: 20827271]
105. Spellman TJ, Gordon JA. Synchrony in schizophrenia: a window into circuit-level pathophysiology. *Curr Opin Neurobiol*. 30:17–23.2015; [PubMed: 25215626]
106. Uhlhaas PJ, Singer W. Oscillations and neuronal dynamics in schizophrenia: the search for basic symptoms and translational opportunities. *Biol Psychiatry*. 77:1001–1009.2015; [PubMed: 25676489]
107. Hall J, Trent S, Thomas KL, O'Donovan MC, Owen MJ. Genetic risk for schizophrenia: convergence on synaptic pathways involved in plasticity. *Biol Psychiatry*. 77:52–58.2015; [PubMed: 25152434]
108. Fromer M, et al. De novo mutations in schizophrenia implicate synaptic networks. *Nature*. 506:179–184.2014; [PubMed: 24463507]
109. Malenka RC, Nicoll RA. NMDA-receptor-dependent synaptic plasticity: multiple forms and mechanisms. *Trends Neurosci*. 16:521–527.1993; [PubMed: 7509523]

110. Schizophrenia_Working_Group_of_the_Psychiatric_Genomics_Consortium. Biological insights from 108 schizophrenia-associated genetic loci. *Nature*. 511:421–427.2014; [PubMed: 25056061]
111. Sekar A, et al. Schizophrenia risk from complex variation of complement component 4. *Nature*. 530:177–183.2016; [PubMed: 26814963]
112. Goldman-Rakic PS. Working memory dysfunction in schizophrenia. *J Neuropsychiatry Clin Neurosci*. 6:348–357.1994; [PubMed: 7841806]
113. Selemon LD, Kleinman JE, Herman MM, Goldman-Rakic PS. Smaller frontal gray matter volume in postmortem schizophrenic brains. *Am J Psychiatry*. 159:1983–1991.2002; [PubMed: 12450946]
114. Ahn S, Phillips AG. Daily Monitoring of Dopamine Efflux Reveals a Short-Lasting Occlusion of the Dopamine Agonist Properties of d-Amphetamine by Dopamine Transporter Blockers GBR 12909 and Methylphenidate. *ACS Chem Neurosci*. 2013
115. Preuss TM. Do rats have prefrontal cortex? The Rose-Woolsey-Akert program reconsidered. *J Cogn Neurosci*. 7:1–24.1995; [PubMed: 23961750]
116. Goghari VM, Sponheim SR, MacDonald AW 3rd. The functional neuroanatomy of symptom dimensions in schizophrenia: a qualitative and quantitative review of a persistent question. *Neuroscience and biobehavioral reviews*. 34:468–486.2010; [PubMed: 19772872]
117. Simen AA, DiLeone R, Arnsten AF. Primate models of schizophrenia: future possibilities. *Prog Brain Res*. 179:117–125.2009; [PubMed: 20302824]
118. Nuechterlein KH, et al. The MATRICS Consensus Cognitive Battery, part 1: test selection, reliability, and validity. *Am J Psychiatry*. 165:203–213.2008; [PubMed: 18172019]
119. Carter CS, Minzenberg M, West R, Macdonald A 3rd. CNTRICS imaging biomarker selections: Executive control paradigms. *Schizophrenia bulletin*. 38:34–42.2012; [PubMed: 22114099]
120. Barch DM, Moore H, Nee DE, Manoach DS, Luck SJ. CNTRICS imaging biomarkers selection: Working memory. *Schizophr Bull*. 38:43–52.2012; [PubMed: 22080498]
121. Butler PD, et al. Perceptual measurement in schizophrenia: promising electrophysiology and neuroimaging paradigms from CNTRICS. *Schizophr Bull*. 38:81–91.2012; [PubMed: 21890745]
122. Lee J, Park S. Working memory impairments in schizophrenia: a meta-analysis. *Journal of Abnormal Psychology*. 114:599–611.2005; [PubMed: 16351383]
123. Goldman-Rakic PS. Cellular basis of working memory. *Neuron*. 14:477–485.1995; [PubMed: 7695894]
124. Funahashi S, Bruce CJ, Goldman-Rakic PS. Mnemonic coding of visual space in the monkey's dorsolateral prefrontal cortex. *J Neurophysiol*. 61:331–349.1989; [PubMed: 2918358]
125. Chafee MV, Goldman-Rakic PS. Matching patterns of activity in primate prefrontal area 8a and parietal area 7ip neurons during a spatial working memory task. *J Neurophysiol*. 79:2919–2940.1998; [PubMed: 9636098]
126. Funahashi S, Bruce CJ, Goldman-Rakic PS. Dorsolateral prefrontal lesions and oculomotor delayed-response performance: evidence for mnemonic “scotomas”. *J Neurosci*. 13:1479–1497.1993; [PubMed: 8463830]
127. Chafee MV, Goldman-Rakic PS. Inactivation of parietal and prefrontal cortex reveals interdependence of neural activity during memory-guided saccades. *J Neurophysiol*. 83:1550–1566.2000; [PubMed: 10712479]
128. Sawaguchi T, Goldman-Rakic PS. The role of D1-dopamine receptor in working memory: local injections of dopamine antagonists into the prefrontal cortex of rhesus monkeys performing an oculomotor delayed-response task. *J Neurophysiol*. 71:515–528.1994; [PubMed: 7909839]
129. Park S, Holzman PS, Goldman-Rakic PS. Spatial working memory deficits in the relatives of schizophrenic patients. *Arch Gen Psychiatry*. 52:821–828.1995; [PubMed: 7575101]
130. Driesen NR, et al. Impairment of working memory maintenance and response in schizophrenia: functional magnetic resonance imaging evidence. *Biological psychiatry*. 64:1026–1034.2008; [PubMed: 18823880]
131. Eryilmaz H, et al. Disrupted Working Memory Circuitry in Schizophrenia: Disentangling fMRI Markers of Core Pathology vs Other Aspects of Impaired Performance.

Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology. 41:2411–2420.2016; [PubMed: 27103065]

132. Moghaddam B, Krystal JH. Capturing the angel in “angel dust”: twenty years of translational neuroscience studies of NMDA receptor antagonists in animals and humans. *Schizophrenia bulletin*. 38:942–949.2012; [PubMed: 22899397]
133. Javitt DC, Zukin SR, Heresco-Levy U, Umbricht D. Has an angel shown the way? Etiological and therapeutic implications of the PCP/NMDA model of schizophrenia. *Schizophrenia bulletin*. 38:958–966.2012; [PubMed: 22987851]
134. Wang M, et al. NMDA receptors subserve persistent neuronal firing during working memory in dorsolateral prefrontal cortex. *Neuron*. 77:736–749.2013; [PubMed: 23439125]
135. Skoblenick KJ, Womelsdorf T, Everling S. Ketamine Alters Outcome-Related Local Field Potentials in Monkey Prefrontal Cortex. *Cerebral cortex*. 26:2743–2752.2016; [PubMed: 26045564]
136. Ma L, Skoblenick K, Seamans JK, Everling S. Ketamine-Induced Changes in the Signal and Noise of Rule Representation in Working Memory by Lateral Prefrontal Neurons. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 35:11612–11622.2015; [PubMed: 26290238]
137. Skoblenick K, Everling S. NMDA antagonist ketamine reduces task selectivity in macaque dorsolateral prefrontal neurons and impairs performance of randomly interleaved prosaccades and antisaccades. *The Journal of neuroscience : the official journal of the Society for Neuroscience*. 32:12018–12027.2012; [PubMed: 22933786]
138. Evans S, et al. Performance on a probabilistic inference task in healthy subjects receiving ketamine compared with patients with schizophrenia. *Journal of psychopharmacology*. 2012
139. Jones JA, Sponheim SR, MacDonald AW 3rd. The dot pattern expectancy task: reliability and replication of deficits in schizophrenia. *Psychol Assess*. 22:131–141.2010; [PubMed: 20230159]
140. MacDonald AW 3rd. Building a clinically relevant cognitive task: case study of the AX paradigm. *Schizophr Bull*. 34:619–628.2008; [PubMed: 18487225]
141. Barch DM, Carter CS, MacDonald AW 3rd, Braver TS, Cohen JD. Context-processing deficits in schizophrenia: diagnostic specificity 4-week course and relationships to clinical symptoms. *Journal of Abnormal Psychology*. 112:132–143.2003; [PubMed: 12653421]
142. Blackman RK, Macdonald AW 3rd, Chafee MV. Effects of ketamine on context-processing performance in monkeys: a new animal model of cognitive deficits in schizophrenia. *Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology*. 38:2090–2100.2013; [PubMed: 23660706]
143. Dias EC, et al. Changing plans: neural correlates of executive control in monkey and human frontal cortex. *Experimental brain research*. 174:279–291.2006; [PubMed: 16636795]
144. Karayiorgou M, Simon TJ, Gogos JA. 22q11.2 microdeletions: linking DNA structural variation to brain dysfunction and schizophrenia. *Nature reviews Neuroscience*. 11:402–416.2010; [PubMed: 20485365]
145. Stark KL, et al. Altered brain microRNA biogenesis contributes to phenotypic deficits in a 22q11-deletion mouse model. *Nature genetics*. 40:751–760.2008; [PubMed: 18469815]
146. Sigurdsson T, Stark KL, Karayiorgou M, Gogos JA, Gordon JA. Impaired hippocampal-prefrontal synchrony in a genetic mouse model of schizophrenia. *Nature*. 464:763–767.2010; [PubMed: 20360742]
147. Fenelon K, et al. Deficiency of Dgcr8, a gene disrupted by the 22q11.2 microdeletion, results in altered short-term plasticity in the prefrontal cortex. *Proceedings of the National Academy of Sciences of the United States of America*. 108:4447–4452.2011; [PubMed: 21368174]

Summary of main findings

Animal models have failed to yield new treatments for psychiatric disorders. Some psychiatric disorders may result from pathology in plasticity mechanisms. Therefore, understanding plasticity mechanisms in model systems may provide insight into the disordered processes in patients.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

Reinstatement of fear following retrieval-extinction procedure. A. Conditioning paradigm used to induce and extinguish fear under either a standard extinction protocol or a retrieval-extinction protocol in which a fear memory is retrieved and rendered labile one hour before extinction. B. Behavioral performance following extinction and reinstatement.

Figure 2.

Reinstatement of nose-poke behavior following acquisition and extinction. A. Behavioral procedure used to extinguish heroin self-administration using either a standard extinction protocol or a retrieval-extinction procedure. B. Behavioral response during extinction. C. Active nose-poke behavior following reinstatement driven by heroin delivery.

Figure 3.

NMDAR antagonists in monkeys replicate both behavioral and neurophysiological features of cognitive deficits in patients with schizophrenia. **A.** BOLD signal reflecting maintenance of the B-cue in working memory is reduced in prefrontal cortex of schizophrenia patients relative to controls performing the AX-CPT. **B.** Focal iontophoretic application of an NMDAR antagonist in prefrontal cortex of monkeys performing the oculomotor delayed response (ODR) task reduces the spatially selective persistent firing of a prefrontal neuron associated with maintenance of the cue location in spatial working memory. **C.** Patients with schizophrenia (red) make selectively more errors on BX-trials relative to controls on the AX-CPT (black). **D.** In monkeys performing the AX-CPT, systemic injections of ketamine (an NMDAR antagonist) induce the same error pattern – a selective increase in errors on BX-trials (red line) relative to control (saline injections; black)¹⁴².