The author thanks Wg Cdr PVPD Singh for the keen interest shown and the critical review of the article. His letter has enabled us to elaborate various points. It has also assisted in improved presentation of the various critical points in the pathogenesis of cholelithiasis in this remote area of the country for which published data is scanty. The experience derived could be used to improve future studies.

REFERENCES

1. Menon PK, Mahapatra S. Study of serum and bile from patients of cholelithiasis in North Eastern India. *Medical Journal Armed Forces India* 1995; 51 : 9-12.
2. Prakash A. Chronic cholecystitis and cholelithiasis in India. *International Surgery* 1968; 49 : 79-85.
3. Malhotra SL. Epidemiological study of cholelithiasis among rail road workers in India with special reference to causation. *Gut* 1968; 9 : 290-5.
4. Krishnan AN, Narayanan K, Kapur BML. Regional variation in lithogenicity of bile in Indian patients. *Ind J Surg* 1984; 46 : 298-304.
5. Irwin JL, Johnston CG, Kapala J. A photometric method for determination of cholates in bile and blood. *J Biol Chem* 1953 : 439-57.
6. Carey JB. The serum trihydroxy-dihydroxy bile acid ratio in liver and biliary tract disease. *J Clin Invest* 1958; 37 : 1494-503.
7. Vlahcevic ZR, Bell CC, Buhac I, Farrar JT, Swell L. Diminished bile acid pool size in patients with gall stones. *Gastroenterology* 1970; 59 : 165-9.
8. Roslyn JJ, DenBesten L, Thompson JE, Cohen K. Chronic cholelithiasis and decreased bile salt pool size : cause or effect? *American J Surg* 1980; 139 : 119-23.
9. Connerty HV, Briggs AR, Eaton III Jr. Serum phospholipid estimation. *Clin Chem* 1961; 37 : 580-7.
10. Wani MA, Kariholu PL, Misgar MS, Wani MY, Khan M. Phospholipid and cholesterol estimation in hepatic duct and gall bladder bile in patients of cholelithiasis with functioning gall bladder in Kashmir. *Ind J Surg* 1986; 48 : 346-9.
11. Ramachandran K, Pushpalata M, Varkey L, Ravchandranath KK, Balasaraswathi KK. Biochemical assessment of cholelithiasis. *J Assoc Physicians India* 1992; 40 : 622-3.
12. Stott AW, Griffin TFR. Reversed serum lecithin : cholesterol ratios in relation to gall stone formation as a possible aid to Diagnosis. *Br J Clin Pract* 1973; 27 : 87-90.
13. Jones DP, Sosa FR, Shartsis J, Shah PT, Skromak E, Behr WT. Serum cholesterol esterifying and cholesterol ester hydrolyzing activities in Liver diseases. Relationships to cholesterol, Bilirubin and bile salt concentrations. *J Clin Invest* 1971; 50 : 259-65.
14. Taylor P. Cholinergic agonists. In : Goodman Gilman's *The pharmacological basis of Therapeutics*. 7th ed. New York, McMillan 100-9.

DIVING AT HIGH ALTITUDE. PRESSURE MEDICINE - A NEW DISCIPLINE ?

Dear Editor,

The lack of authoritative Indian research on high altitude illnesses has been lamented in relation to the dependence on Western literature in a recent editorial. The authors have acknowledged past contributions in this field by service medical officers and it is heartening to see renewed directions by Indian research workers after many decades [1].

We wish to place on record our experience with Indian Navy divers in conducting in 1988 the worlds highest altitude scuba dives at Pangong Tso in Ladakh at an altitude of 14,200 feet. The experiences were published in a leading international journal in the field of Diving Medicine [2]. This was acknowledged as being "Outstanding" by the Science Digest USA and is now used as a guideline for those venturing into the realm of high altitude, hyperbaric, diving and adventure medicine. The Limca Book of Records also documents this dive [3].

Divers of the Indian Navy performed "no decompression" dives to a maximum of 140 feet sea water (42.6 m of sea water) equivalent ocean depth at a maximum altitude of 14,200 feet (4328 m) and minimum water temperature of 2°C. A total of 100 dives were performed at

altitudes ranging from 7000 to 14,200 feet (2134 to 4328 m). Prior to this the highest documented dive was that at 12500 feet (3810 m) in 1968 in Lake Titicaca by Jacques Cousteau. Subsequently in 1989 a British group conducted diving in the Nepali Himalayas at a maximum altitude of 16000 feet (4877 m) in Donag Tsho and to a maximum depth of 98 feet of sea water (29.3 m of fresh water) [4].

Diving at extreme altitudes in remote areas has the fascination of adventure in addition to the thrill of diving in unknown waters. Sufficient documented dives have not been carried out to recommend standard procedures. Also equipment used is untested at these altitudes. Apart from the physiological adaptation to altitude and cold, the divers have to plan for survival in remote areas with sparse medical or rescue facilities.

Diving procedures need special consideration since the divers first have to adapt to the low pressures, hypoxemia and reduced physical capabilities at altitude. On diving they are exposed to increasing pressures and on return to surface to the low ambient pressure. For use in diving emergencies we were carrying a monoplace hyperbaric chamber but the divers could be exposed to low pressures while being evacuated by unpressurized helicopters to the hospital at Leh.

To appreciate the changes in diving principles while

diving at altitude we present a small aspect. At sea level (760 mmHg pressure = 1 Atmosphere Absolute (ATA)) a diver needs to descend to a depth of 10 m to be at a pressure of 2 ATA which is double that of sea level. At an altitude of 14000 feet (4267 m) the ambient pressure is 447 mmHg and a diver has to dive to only 6 m to feel the effects of double the pressure. Hence for every unit of depth under water the pressures increases at a much more rapid rate. Also on return he has to adapt to a surface pressure which is nearly half of sea level. His body has to adapt to these rapid changes of pressure and repeatedly depending on the number of dives carried out every day.

We reviewed all available literature in the fields of aviation and aerospace medicine which deals with rapid ascents to low pressures, in the field of high altitude medicine which relates to slower ascent and long term effects of residence at low pressures and in the field of diving medicine which relates to rapid increased in pressure under water. In the Armed Forces we have specialists experienced in each of these fields. It was when planning

these dives we felt that it may be time for a holistic view of these multidimensional problems to be addressed under the umbrella cover of "Pressure Medicine". This may be food for thought!

MAJ A DHALL^{*}, SURG CDR TK SAHNI⁺

***Graded Specialist (Medicine), 184 MH C/O 56 APO.**

⁺Classified Specialist (Medicine), Medical Directorate, Naval Headquarters, New Delhi - 110 011.

REFERENCES

1. Rao KS, Apte CV. Medical Research at High Altitude. *Medical Journal Armed Forces India* 1995; 51 : 81-2.
2. Sahni TK, John MJ, Dhall A, Chatterjee AK. High altitude dives from 7000 to 14200 feet in the Himalayas. *Undersea and Biomedical Research* 1991; 18 : 303-16.
3. Limca Book of Records 1995; 197.
4. Leach J, Mclean A, Mee FB. High Altitude dives in the Nepali Himalayas. *Undersea & Hyperbaric Medicine* 1994; 21 : 459-66.