ELECTROCARDIOGRAPHIC QUIZ

Maj RG PODUVAL*

Clinical information

A 60-year-old male patient, a known case of ischaemic heart disease was admitted with complaints of palpitations, orthopnea and anginal chest pain of one day duration. Examination revealed an obese patient with tachycardia, unrecordable blood pressure and peripheral pulses, raised JVP with no cannon waves, a loud first heart sound with unvarying intensity and bilateral pulmonary basal crepitations.

ECG before and after treatment with intravenous verapamil is shown in Fig 1 and Fig 2 respectively.

What is your diagnosis?

Fig. 1:

Fig. 2:

*Classified Specialist (Medicine), 153 General Hospital, C/O 56 APO.

ANSWER TO THE QUIZ (Page 180)**Interpretation**

The ECG (Fig 1) shows a wide-QRS tachycardia of 250/min, concordant positivity i.e. all complexes in V1-V6 are positive, monophasic R wave in V1, and axis in the right upper quadrant, all suggestive of ventricular tachycardia. However, close examination of V6 reveals distinct P waves related to the triphasic QRS complex, both of which are features of supraventricular tachycardia (SVT) with aberrant conduction. After administration of intravenous verapamil the tachycardia reverted to normal sinus rhythm (Fig 2).

Discussion

Wide-QRS tachycardia is a commonly encountered emergency in clinical medicine. Before the present day antiarrhythmic drugs were discovered, digitalis was the mainstay in the treatment of supraventricular tachycardia. However, the adverse consequences of giving it for ventricular tachycardia are well known. Hence stress was laid on differentiation between VT and SVT with aberrant conduction, by clinical and ECG evaluation. The clinical features favouring VT are cannon waves in neck veins, varying intensity of first heart sound (both indicating atrioventricular dissociation) and wide splitting of heart sounds (indicating atrioventricular asynchrony). A common misconception was that the blood pressure was low in VT and many VT were misdiagnosed as SVT when the blood pressure was normal [1].

To differentiate VT from SVT with aberrant conduction the following electrocardiographic features should be looked for [2,3] :

1. Evidence of preceding atrial activity for SVT. Esophageal leads are helpful if P waves are hidden in the QRS complex.
2. QRS duration more than 140 ms for VT.

3. QRS morphology : Features of QRS morphology that favour SVT are RBBB or triphasic patterns like rSR in V1 and qRS in V6. Monophasic pattern like R or qR in V1 and rS or QS in V6 or multiple morphology QRS complexes favour VT.

4. AV dissociation for VT.

In some cases SVT with aberration is initiated by an aberrant conducted beat which follows a long RR interval as the long cycle results in lengthening of the following refractory period and this favours aberrant conduction. SVT with aberration can sometimes be diagnosed by giving intravenous procainamide which blocks AV conduction and helps in identifying the P waves amongst the widely spaced ventricular complexes.

Ventricular aberration is of 3 types [4]. Type A - fascicular refractiveness (RBBB most common). Type B - anomalous activation and conduction via bypass tract. Type C - paradoxical critical rate (aberration following a sinus beat with a long refractory period).

Irrespective of the type, the primary importance of aberration is its misdiagnosis as ventricular tachycardia. Thus the emphasis on meticulous clinical and ECG evaluation for selecting the appropriate antiarrhythmic drug.

REFERENCES

1. Morady F, Dicarto AL, Krol BR. A prevalent misconception regarding wide complex tachycardia. *JAMA* 1985; 254: 2790-1.
2. Dancy MM, Canon AJ, Ward D. Misdiagnosis of chronic recurrent ventricular tachycardia. *Lancet* 1985; 2: 320-3.
3. Wellens HJJ, Frits WHM, Lie KI. The value of the electrocardiogram in the differential diagnosis of a tachycardia with a widened QRS complex. *Am J Med* 1978; 64: 27-32.
4. Marriot HJL. Aberrant ventricular conduction and the diagnosis of wide QRS tachycardia. In: *Practical electrocardiography*. 8th ed. Baltimore: Williams & Wilkins, 1988; 235-64.