Postgraduate Clinic CARCINOID SYNDROME

Sqn Ldr S SHANKAR ^{*}, Lt Col RAJAT KUMAR ⁺,
Brig SP KALRA [#], Col PS REDDY ^{**}, Lt Col SK SHARMA ⁺,
Lt Col KS RAO ⁺⁺, Maj A MENON ^{##}

MJAFI 1998; 54 : 340-342

KEY WORDS : Carcinoid crisis; Carcinoid syndrome.

Introduction

Carcinoid tumors are the commonest gastrointestinal endocrine tumours with an annual incidence of 1.5/lakh population [1]. systemic symptoms appear when the tumours metastasise to liver and manifest with features of carcinoid syndrome [2]. Though the symptoms and signs are characteristic of the illness, the diagnosis is often missed due to lack of awareness amongst the treating physicians.

Illustrative Case

A 37 years old labourer was admitted in Jan 97 with progressive breathlessness on exertion and palpitations of 6 months duration. Two months later he developed episodes of watery diarrhoea which lasted for 4-5 days. He had 3-4 such episodes every month which were accompanied by severe pain in upper abdomen. At his village he was treated with multiple brandings over abdomen.

Examination revealed: Pulse 100/mt, BP 100/60 mmHg, facial puffiness, bilateral pitting pedal edema and raised JVP to angle of jaw with prominent V waves. Cardiac examination showed findings of right ventricular hypertrophy, pulmonary stenosis and tricuspid regurgitation. On abdominal examination liver was palpable 8 cm below costal margins (Span 20 cms). It was hard, nodular, pulsatile and non tender. There was evidence of minimal ascites. A week later, a reddish hue was noticed over the chest which blanched on pressure and disappeared after 48 hrs. In view of his diarrhoea, pain abdomen, right sided cardiac lesions with cardiac failure, hard nodular liver and flushing-a clinical diagnosis of Carcinoid syndrome with multiple liver metastasis was made.

On investigations, his haemogram, urinalysis, metabolic parameters and chest Xray were normal. ECG showed right axis deviation (+135 degrees) with right ventricular hypertrophy. 2D echocardiography revealed a thickened pulmonary valve with pulmonary stenosis, tricuspid regurgitation and a dilated right atrium (Fig-1). 24 hour urinary 5-HIAA was markedly increased to 47.3 mg/day (normal 2-8 mg/day). Ultrasonography (USG) abdomen showed multiple hyperechoic lesions in liver consistent with metastasis. Fine needle aspiration cytology (FNAC) from a liver metastatic nodule under USG guidance showed cluster of cells consistent with neuroendocrine tumor metastasis to liver (Fig-2). An

attempt was made to locate the primary tumor by upper gastrointestinal endoscopy and colonoscopy which were normal. The Barium meal follow through showed rapid emptying of the contrast in the small intestine but the primary tumor was not visualised.

He was managed with bed rest, furesamide (40 mg/day), Cyproheptadine (4 mg twice daily) and ranitidine (150 mg twice daily). He showed improvement in cardiac failure with above therapy and is presently on regular follow up at this centre.

Discussion

Q.1) What was your clinical diagnosis at admission?

Ans. The classic triad [1] of carcinoid syndrome i.e. diarrhoea, flushing and right sided valvular heart lesions is present in this patient hence that was the first diagnosis considered.

Q 2) What is the genesis of flushing, diarrhoeas, valvular involvement and edema in a case of carcinoid syndrome?

Fig. 1 : Apical four chamber view in systole shows closed mitral valve with a thickened and opened tricuspid valve, consistent with tricuspid regurgitation.

^{*} Clinical Tutor, ⁺ Reader, [#] Consultant and Head, ^{##} Physician Trainee, Dept of Medicine, AFMC; ⁺⁺ Classified Specialist, Medicine and Cardiology, MH (CTC); Pune 411 040. ^{**} Senior Adviser in Pathology, Army Hospital (R & R), Delhi Cant 110 010.

Fig. 2 : Fine needle aspiration cytology (FNAC) liver showing uniform population of small tumour cells. Cytoplasm is scanty and nuclear chromatin shows 'salt and pepper' stippled appearance typical of neuroendocrine tumours

Ans. Flushing : the mediators implicated are histamine, tachykinins (neuropeptide K and substance P) and prostaglandins E and F. Serotonin does not cause flushing, but is known to provoke an attack [2].

Diarrhoeas: Serotonin is responsible for the diarrhoea and produces the same by its effects on gut motility, causing fat malabsorption and inducing a secretory state in the small intestine [1].

Valve involvement: Serotonin stimulates fibroblast growth and fibrogenesis which causes diffuse fibrosis of tricuspid and pulmonary valves. These fibrous deposits tend to cause constriction and deformity of pulmonary and tricuspid valves [3,4].

Edema: Edema is result of right heart failure and protein malnutrition due to excess utilisation of tryptophan in the diet (used for synthesis of serotonin [2].

Q 3) How is the development of symptoms explained in carcinoid syndrome?

Ans. The symptoms in carcinoid syndrome are due to different biological mediators produced by tumors that arise in enterochromaffin cells. Such tumors are typically located in the small intestine and the mediators which are released into the portal vein get metabolised by the liver normally. Hence the tumors usually become symptomatic only after hepatic metastasis occur, as the mediators bypass the hepatic degradation and enter the systemic circulation. In cases of bronchial and other extraintestinal carcinoids however, whose hormone products are not cleared by liver, symptoms are produced even in absence of metastasis [1,2].

Q.4) what is the incidence of flushing and what are

the types of flushing encountered in carcinoid syndrome?

Ans. Flushing occurs in 63-94% of patients with carcinoid syndrome [2,5]. Three types of flushing are seen in carcinoid syndrome. two of them occur in the usual metastatic carcinoid. the first is a sudden deep red or violaceous erythema of the face and upper part of the body of short duration which is often associated with an unpleasant feeling of warmth. It may be spontaneous or may be precipitated by stress, alcohol, cheese, exercise or drugs like adrenaline and calcium. the second type is usually more prolonged, produces venous dilatation and purplish hue and may give rise to telangiectasia. the third variety of flushing is seen in bronchial and gastric carcinoid which produce an intense and erythematous flush, sometimes associated with lacrimation, conjunctival suffusion, itching, palpitation and facial edema[6].

Q.5) What are the causes of isolated right sided valvular lesions?

Ans. The causes are (i) Congenital-e.g. pulmonary stenosis, tricuspid stenosis, Ebstein's anomaly. (ii) Infective endocarditis in intravenous drug abusers (Tricuspid regurgitation). (iii) Endomyocardial fibroelastosis (Tricuspid regurgitation). (iv) Carcinoid syndrome (Pulmonary stenosis, tricuspid regurgitation).

Q6) what are the common sites of carcinoid tumor?

Ans. 90% of all carcinoid tumors are found in the gastrointestinal tract (predominantly-appendix, ileum and rectum) with appendix alone accounting for 50% of all tumors [1,7]. Pancreas and bronchi are the other common sites and rarer sites are thymus, esophagus, Meckel's diverticulum and ovary. While upto 60% of small bowel carcinoids are associated with systemic manifestation, only 3.5% of bronchial, 1% of appendix and virtually no rectal carcinoid produce the symptoms [2,8].

Q 7) How do you diagnose carcinoid syndrome?

Ans. When the clinical suspicion of carcinoid syndrome is present, the diagnosis is based on the measurement of urinary levels of serotonin and its metabolites. Once dietary 5-hydroxyindole containing substances like bananas, plantain, pineapple, plums, walnuts and acetaminophen are excluded, urinary 5-HIAA levels of 25mg/day are diagnostic. Elevation in the range of 9-25 mg may be seen in non tropical sprues, acute intestinal obstruction and carcinoid syndrome. Raised urinary 5 HIAA level has a 73% sensitivity and 100% specificity for the diagnosis of carcinoid syndrome [9]. additional information may be

gained by urinary and platelet concentrations of serotonin which are elevated.

Q 8) How do you localise the primary in a case of carcinoid syndrome?

Ans. The small intestinal carcinoids, the commonest to cause the syndrome, are also the most difficult to localise. A barium meal follow through or small bowel enteroclysis may reveal the carcinoid, though it is often missed by them. A CT scan or angiography are marginally better. The best results are however obtained with radio-nucleide scanning with ^{123}I MIBG, [10] and ^{123}I octreotide [11] which has 87% sensitivity. An UGI endoscopy/colonoscopy can help localise a gastric, rectal or colonic carcinoid. Bronchial carcinoids are relatively easier to localise and can be done so by chest Xray, CT scan, MRI scan and bronchoscopy.

Q 10) How do you treat a case of carcinoid syndrome?

Ans. Resection of the primary tumor causes relief of symptoms in an extra intestinal or bronchial carcinoid without metastasis. However, in a patient with hepatic metastasis the first line therapy consists of symptomatic treatment [12]. Flushing can be prevented by avoiding precipitating foods. Diarrhoea responds to antidiarrhoeal agents like loperamide or 5 Hydroxy Tryptamine antagonists like cyproheptadine, ketanserin or ondansetron. Wheezing can be treated with aminophylline or selective β_2 agonists like salbutamol. Heart failure responds to diuretics and occasionally may require a tricuspid valve replacement. Octreotide, a somatostatin analogue, has been shown to provide a good control of diarrhoea, flushing and wheezing in 75% of cases. It is given in the dose of 75-150 μg subcutaneously 8 hourly which can be increased upto 750 μg daily to achieve control of symptoms [13]. Interferon alpha in dose of 24 millions units/ M^2 of body surface area thrice a week for one year has been found to be of limited role [4].

Q 11) What is carcinoid crisis?

Ans. Carcinoid crisis is the acute manifestation of carcinoid syndrome characterized by hypotension, tachycardia, diarrhoea, flushing and occasionally angina. It is usually seen with foregut carcinoids and in patients excreting 200 mg/day urinary 5 HIAA. It may be spontaneous or brought on by stress, anaesthesia, chemotherapy or liver biopsy. It is managed by administration of octreotide [15,16].

Q 12) What is the prognosis of carcinoid tumor?

Ans. The prognosis is dependent on the stage of

tumor [7]. For local tumors the 5 years survival is 94%, with lymph node involvement it is 65% and only 18% with distant metastasis. Urinary 5 HIAA levels correlate inversely with the prognosis [2]. The median survival for patients excreting 10-49 mg/day is 29 months, 50-149 mg/day is 24 months and 150 mg/day only 13 months.

REFERENCES

1. Kaplan LM. Endocrine tumors of the gastrointestinal tract and pancreas. In: Isselbacher, Braunwald, Wilson, Martin, Fauci, Kasper, editors. Principles of Internal Medicine. New York: McGraw Hill Inc, 1994;1537-42.
2. Norton JA, Levin B, Jensen RT. Cancer of the endocrine system. In: DeVita VT, Hellman S, Rosenberg SA, editors. Cancer, principles and practice of oncology. 4th ed. Philadelphia: Lippincott, 1983; 1333-435.
3. Lundil L. Carcinoid heart disease. Acta Oncol 1991;30: 499-502.
4. Ross EM, Roberts WC. The carcinoid syndrome: comparison of 21 necropsy subjects with carcinoid heart disease to 15 subjects without carcinoid heart disease. Am J Med 1985;79:339-54.
5. Norheim I, Oberg K, Theodorsson-Norheim E. Malignant carcinoid tumors. Ann Surg 1987;206:115-25.
6. MacDonald J, Haller D, Weigel RJ. Endocrine system. In: Abeloff MD, Armitage JO, Lichter AS, Neiderhuer JE, editors. Clinical Oncology. 2nd ed. New York: Churchill Livingstone, 1995;1047-81.
7. Godwin JD. Carcinoid tumors: an analysis of 2837 cases. Cancer 1975;36:560-69.
8. Rothmund M, Stinner D, Arnold R. Endocrine pancreatic carcinoma. Eur J Surg Oncol 1991;17:191-8.
9. Feldman JM. Carcinoid tumors and syndrome. Semin Oncol 1987;15:237-46.
10. Hanson MW, Feldman JM, Blinder RH, Moore JO, Coleman RE. Carcinoid tumors- ^{123}I MIBG scintigraphy. Radiology 1989;172:699-703.
11. Lamberta SW, Bakker WH, Reubi JC, Krenny EP. Somatostatin receptor imaging in the localisation of endocrine tumors. N Engl J Med 1990;323:1246-9.
12. Saini A, Waxman J. Management of carcinoid syndrome. Postgrad Med 1991;67:505-12.
13. Oates JA. Carcinoid syndrome. In: Bennet JC, Plum F, editors. Cecil Textbook of medicine. 20th ed. Philadelphia, WB Saunders: 1996;1348-50.
14. Oberg K. The action of interferon alpha on human carcinoid tumors. Semin Cancer Biol 1992;3:35-44.
15. Kvols LK, Moertel CG, O'Connell MJ. The treatment of the malignant carcinoid syndrome with a long acting somatostatin analogue. N Engl J Med 1986;315:663-6.
16. Dissonette RT, Gibne RG, Berry RR et al. Fatal carcinoid crisis after percutaneous fine needle biopsy of hepatic metastasis: case report and literature review. Radiology 1990;174:751-5.