High risk mothers are those receiving anticonvulsants, antitubercular drugs, coumarin derivatives and salicylates.

High risk neonates are those with prematurity, birth asphyxia and IUGR

REFERENCES

1. MK Behra, SD Kulkarni. Vitamin K deficiency hemorrhagic disease of newborn and present controversies. *MJAFI* 1998;54:143-5.
2. Malia RG, Preston FE, Mitchell VE. Evidence against vitamin K

deficiency in normal neonates. *Thromb Hemost* 1980; 44 :159-60.

3. Singh M. Do breast fed healthy neonates need vitamin K supplementation? *Academy Today* April 1991,p-12
4. Singh M. Vitamin K during infancy: Current status and recommendations. *Indian Pediatr* 1997; 34 : 708-12

Gp Capt TSRAGHU RAMAN

Senior Advisor (Pediatrics) Command Hospital, Bangalore-560 007.

LET'S TAKE A BIG LEAP

Dear Editor,

The states and Union Territories have been asked to gear up measures to meet the possible shortage in blood supply following the ban of professional blood donation system immediately. This was the news in Navhind Times on 25 Jan 98. The news continued "In public interest litigation between common cause vs Union Of India and others, the Supreme Court had directed that steps should be taken to discourage the prevalent system of professional blood donors and this system should be completely eliminated by the end of December last. Following the Supreme Court direction, the Government has initiated measures to promote voluntary blood donation to meet the possible short fall of blood supply". Will it be successful?

Blood donation is said to be a matter of routine and a well accepted norm in the Armed Forces and therefore no significant shortage is envisaged during peace time. But it is also alleged that volunteers are stage-managed through orders and concealed force. If the fact is that people have to be coerced and forced for blood donation, then it is certain that they need education. One of the effective methods prescribed for education is to start from 'known' and proceed to 'unknown'. An effort was made to find out the level of awareness, knowledge and misgivings about blood donations through a survey conducted amongst 1000 service personnel and their families in a Training Centre.

68% of them including 61% ladies volunteered to bleed. More than half of them considered that blood donation was a noble deed. 15% of them would donate blood as it was life and limb saving. About one sixth of them thought that it was their duty to donate blood to a colleague, friend or relative.

But the most significant finding is that 15% of them knew, an occasional bleeding is harmless and it could even stimulate blood

production. Curiously maximum number of such believers hailed from the fairer sex. Forced volunteers were limited to a mere 3%.

One third of the surveyed population remained non-believers in the sanctity of voluntary blood donation. Their reasons ranged from fear of needles (30%), weakness generation (27%), fear of infection(9%), inadequacy of blood (8%) to a ridiculous thought of causing infertility (1%). 25% of them feared that their diseases will be known to others during blood donation tests.

These inferences are significant and valuable. Majority of the non-volunteers can be easily converted into proud volunteers through small lessons of education and demonstration. Many of them are unaware of the volume of blood that will be drawn from each individual at a time and the total volume of blood that an individual has. Some of them think that the whole process is a painful episode of the invasion of needles. Many are amazed to know that the healthy donors can go to their normal duties immediately after blood donation. The blood donated never comes back. It is true. But what they should know is that a donation triggers off the whole assembly line of blood production soon replacing the losses with fresh and young blood. They are ignorant that there are many reasons beyond their control for their rejection as donors. They should be informed in no uncertain terms that any detection of diseases during the blood testing process will be kept confidential. The medical fraternity has the moral responsibility to alleviate their apprehensions about causing infertility and infection.

Let's take a big leap to create a net work of 100% voluntary blood donors in the Armed Forces and the nation will follow and reward us in our crisis.

Lt Col TOMMY VARGHESE

Commanding Officer, MH Panaji.

EPIDEMIOLOGY OF UNDER FIVE MALNUTRITION: SEX DIFFERENTIAL IN HEALTH CARE AND NUTRITIONAL STATUS (UNDER FIVE MALNUTRITION)

Dear Editor,

This is in reference to the article titled 'Epidemiology of under five malnutrition: Sex differential in health care and nutritional status (under five malnutrition) (1). The following comments are offered.

1. Definition of the population under study in terms of ethnic background, socioeconomic status including dependency ratio and other environmental influences is a prerequisite criterion for any attempt on epidemiological study.

2. Classification of malnutrition as per Indian Academy of Paediatrics is graded into four groups based on the national centre health statistics (NCHS) charts. The cut off point taken is 80% of

the 50th percentile for the age and sex.

3. Anaemia is a nutrient deficiency and usually part of overall malnutrition syndrome. Unless correlation is made between the grade of malnutrition and haemoglobin levels, it will be inappropriate to draw conclusion regarding gender bias leading to lower haemoglobin levels in girls.

4. In a greater context, maternal malnutrition reflects the outcome of pregnancies. There is greater incidence of low birth weight babies born to malnourished mothers. A female low birth weight baby subjected to nutritional inadequacy remains nutritionally compromised during the first five years. Hence the important factors of maternal malnutrition and birth weight should be included in the

study parameters.

REFERENCE

1. A Banerjee. Epidemiology of under five malnutrition: sex differential in health care and nutritional status (under five malnutrition). MJAFI, 1998, 54; 119-20

1998, 54; 119-20

Gp Capt TS RAGHU RAMAN

Senior Advisor (Pediatrics). Command Hospital (Air Force), Bangalore-560 007

SEZARY SYNDROME-CLINICO HISTOLOGICAL CORRELATION

Dear Editor,

This refers to the report on two cases of Sezary Syndrome (MJAFI, 1998;54: 76-78).

Sezary Syndrome (SS) comprises a triad of fiery red erythroderma, generalized lymphadenopathy and circulating atypical cells with cerebriform nuclei (Sezary cells) exceeding an absolute value of 1000/cumm or exceeding 10 percent of circulating cells [1].

Erythroderma, defined as an inflammatory skin disease which affects more than 90 percent of the body surface [2], is a characteristic feature of this syndrome. though 'multiple erythrodermic patches' with 'butterfly rash on face' in the first case and 'multiple erythematous scaly lesions' in the second case of the report do not indicate the extent of cutaneous involvement, these descriptions could well suggest the clinical impression of disseminated lupus erythematosus and psoriasis respectively; more so in the absence of generalised lymphadenopathy as was in both these cases. Cytomorphologically, Sezary cell is not pathognomonic of SS, as it can be seen in many benign dermatoses like chronic eczema, psoriasis, lupus erythematosus, parapsoriasis, atopic dermatitis, vasculitis, etc. [3].

SS, sometimes described as a leukaemic variant of Mycosis Fungoides (MF), is not only clinically and hematologically distinct from it, but also distinct from a histological view, thus discouraging the interchangeable use of both these terms. The infiltrate in SS is less pleomorphic than in MF; and eosinophils, plasma cells, macrophages and accessory cells are less frequent or lacking. [4]. Proliferation Kinetics of the dermal infiltrate using 3H-thymidine labelling (3H index) have revealed widely distinct patterns in both [5]. Lymph nodes in MF have a higher number of interdigitating reticulum cells and the ratio of mononuclear cells having cerebriform

nuclei to the interdigitating reticulum cells varies from 1:1 to 2:1, in contrast to a ratio of 5:1 to 10:1 in SS[6].

The case report serves to highlight the diagnostic dilemma a clinician might have to face when confronted with cutaneous markers of internal disease and also underscores the importance of multi disciplinary approach which is inevitable to overcome such situations.

REFERENCES

1. Arnold HL, Odom RB, James WD. Diseases of the skin (Sezary Syndrome). 8th ed. Philadelphia : WB Saunders Company, 1990; 863-5.
2. Burton JL. Eczema, lichenification, prurigo and erythroderma. In : Champion RH, Burton JL, Ebling FJG. Text Book of Dermatology. 5th ed., Oxford : Blackwell Scientific Publications, 1992; 584-8.
3. Wieselthier JS, Koh HK. Sezary syndrome : Diagnosis, Prognosis and critical review of treatment options. J Am Acad Dermatol 1990; 22 : 381-401.
4. Imai S, Burg G, Braun-Falco O. Mycosis Fungoides and Sezary Syndrome show distinct histomorphological features. Dermatologica 1986; 173 : 131-5.
5. Sterry W, Pullmann H, Steigleder GK. Proliferation kinetics of the dermal infiltrate in cutaneous malignant lymphoma. Arch Dermatol Res 1981; 270 : 285-90.
6. Scheffer E, Meijer CJLM, Van Vloten WA, Willemzen R. A histological study of lymph nodes from patients with the Sezary syndrome. Cancer 1986; 57 : 2375-80.

Lt Col KEDARNATH DASH, Wg Cdr GURCHARAN SINGH, Sqn Ldr SANJIV GROVER

Command Hospital Air Force, Bangalore-560007

ECG DIAGNOSIS: ISORRHYTHMIC ARRHYTHMIA

Dear Editor,

A 52-year-old male patient was admitted with left sided hemiplegia caused by ischemic cerebral infarct along with hypertension. He was then placed on tab Atenolol and tab Disprin. On third day his electrocardiogram (ECG) revealed ventricular tachycardia. His subsequent ECG (Fig. 1) revealed AV junctional rhythm at the rate of 56 beats/min, inverted P-wave indicating retrograde atrial depolarisation, P-R interval 0.18 sec, and QRS duration 0.08 sec (Upper strip). This rhythm was interrupted by a normally conducted sinus beat indicating capture beat (first beat in second strip), a ventricular premature beat (third beat) followed by incomplete compensatory pause. It revealed AV dissociation with both atria and ventricle under control of separate pacemaker (Fig. 2). Rate of both pacemaker is 50 beats/min, however AV junctional rate is slightly faster than sinus rate hence P-wave is seen as marching through the QRS complexes. There is gradual shortening of R-P interval and there is no correlation between P-wave and QRS complexes. This is consistent with diagnosis of incomplete AV

Fig. 1: Electrocardiogram showing AV junctional rhythm