after 3 days of quinine; and pyrimethamine-sulfadoxine resistance too is not infrequent.

1.3% of patients in this series had symptoms persisting for 1 to 2 months and such patients are usually repeatedly treated with empirical chloroquine as reported and observed by us [3]. For such patients the ideal treatment would be quinine with doxycycline to avoid delayed sudden deterioration and high mortality.

The message is that quinine for 5 days with tetracycline or pyrimethamine-sulfadoxine is mandatory for all uncomplicated MT Malaria cases and there should be no hesitation in using artemisine derivatives for proved and ill looking MT Malaria cases.

Col SR MEHTA *, **Lt RAVISHANKAR MK ***.

* Senior Advisor (Medicine), + Medical officer, Medical Division, 92 Base Hospital, C/o 56 APO.

REFERENCES

1. Nandi B. Clinical study of Falciparum Malaria-302 cases Medical Journal Armed Forces India. 1998;54:9-12.
2. Looareesuwan S, Charoan P, Ho M et al. fatal PF Malaria after an inadequate response to quinine treatment. J Infect Dis 1990; 161:577.
3. Mehta SR, Kohle VS, Chauhan SS. The changed clinical spectrum of Malaria due to drug resistance. J Assoc Physician India 1998;46:360-2.

REPLY

I thank Colonel SR Mehta for his valuable comments and advice on my study.

Quinidine was used successfully in seven complicated cases of falciparum malaria (not included in the study). Monitoring QT interval is essential when using quinidine, and doing so in a large number of patients undergoing treatment simultaneously would have been cumbersome. Hence quinidine was not used freely. The initial encouraging results from the alternate regimes used in the study further obviated the need for use of quinidine.

The study was done in 1993. At the time Artemether and its derivatives were not available in the local market.

There is no denying that five day course of quinidine with or without doxycycline/sulpha-pyrimethamine is ideal for uncomplicated falciparum malaria. The alternate regimes used in the study were not meant to be a substitute for the above regime under ideal circumstances. The challenge was to find an alternate regime which can be used in a peripheral hospital with its resource constraints, to treat a large number of patients in the face of acute quinine shortage without undue increase in morbidity/mortality. Within this perspective the alternate regimes used were a reasonable success.

Sqn Ldr B NANDI,

Graded Specialist Medicine, No.12 Air Force Hospital, New Project Area, Gorakhpur -273002.