MULTIPLE MENINGIOMAS

Lt Col HARIQBAL SINGH^{*}, Col SK KHANNA⁺,
Maj VIKAS RASTOGI[#], Lt Col L SATIJA^{**},
Maj SC CHAMOLI⁺⁺

MJAFI 1999; 55 : 263-264

KEY WORDS: Meningioma; Multiple meningiomas.

Introduction

Meningioma are the most common of all extra axial neoplasm occurring within the cranial cavity. They are also the most common non glial primary tumours of the central nervous system. However the occurrence of multiple meningiomas (MM) is uncommon and is usually seen in association with neurofibromatosis. A case of MM not associated with neurofibromatosis is being presented.

Case Report

A 28-year-old woman presented with dull fronto parietal headache of 3 years duration which had increased in intensity and frequency for last 6 months. No history of hypertension or seizures. All routine investigations were within normal limits. Clinical examination revealed cerebellar signs in the form of ataxia and nystagmus. CT brain revealed 3 well defined oval to round high attenuating brightly homogeneously enhancing intracranial mass lesions without calcifications, or perifocal oedema. One seen in the right cerebellar lobe posterior to CP angle with destruction and erosion of petrous bone (Fig 1A). Second seen in the left cerebellar lobe posteriorly with mass effect on fourth ventricle (Fig 1B). It was the largest measuring 33 x30 mm. Third one was seen arising from falx cerebri (Fig 1C and D). Histopathology confirmed it as fibroblastic meningioma.

Discussion

MM not associated with neurofibromatosis are uncommon tumours [1]. Most cases present with chronic headache or complex partial seizures [2]. Among meningiomas the incidence of MM has been reported as low as 3% in pre CT scan era but with advent of CT and MRI its incidence has risen to as high as 16% [3,4].

Detection of meningiomas is more obvious on CT because of superior demonstration as compared to MR because of their low MR tissue contrast characteristics.

MM was first defined as a distinct clinical entity in 1938 by Cushing and Eisenharodt [5]. A series of 8

cases of MM without neurofibromatosis have been described by Turgul et al [1]. Fortunately all his patients

Figs. 1A & B : Cranial CT showing multiple meningiomas (A and B) delineated as brightly enhancing mass lesions in the cerebellum

^{*}, ^{**} Classified Specialist, ⁺ Senior Advisor, [#] Department of Radiology, Command Hospital (Southern Command), Pune 411 040. ⁺⁺ Graded Specialist (Radiology), MH Jalandar Cantt.

Figs. 1C & 1D : Pre and post contrast scans showing bright enhancement in the meningioma arising from the falx cerebri

tolerated multiple surgical interventions well, although the removal of these tumours in critical areas may be difficult [1]. Thus it is important to examine and supervise all patients who have had a meningioma for a possible occurrence of a second meningioma.

The cause of multiplicity of meningiomas is uncertain. Andrioli et al [6] were of the view that there are multiple foci of simultaneous development as a result of mesodermal germ layer susceptibility to unknown tumour producing factors.

REFERENCES

1. Turgul M, Palaogio S, Ozcan OE, Gurcay O, Eryilmaz M. Multiple meningiomas of central nervous system without the stigmata of neurofibromatosis. Clinical and therapeutic study. *Neuro Surg Rev* 1997;20:117-23.
2. Nickell PV. Panic attacks, Complex partial seizures and multiple meningiomas. *Anxiety* 1994;1:40-2.
3. Sahu B, Ranganadham P, Kishore LT, Dinkar J. Multiple meningiomas. *J Assoc Physicians of India* 1992;40:616-7.
4. O'Neill, Booth AE. Multiple meningiomas: Case report and review of literature. *Surg Neurol* 1984;21:80-2.
5. Cushing H, Eisenhardt L. Meningiomas: Their classification regional behaviour, life history and surgical end results. Illinois. Springfield 1938;115-32.
6. Andrioli GC, Rigobello L, Job I, Casentini L. Multiple meningiomas. *Neurochirurgia (Stuttg)* 1981;24:67-9.