Letters to the Editor

OCULAR DEVIATIONS-IS ANAESTHETIC TO BLAME?

Dear Editor,

The article on ocular deviations following cataract surgery was interesting and thought provoking. I have the following queries and observations:

If anaesthetic myotoxicity is primarily responsible for ptosis and ocular deviations then there should have been deviations in some of the 12 patients who underwent a repeat block.

One of the most important causes of post-operative ptosis is the induced uveitis. That's why one invariably finds cases of Acute Iridocyclitis manifesting ptosis. Some degree of ptosis is evident in cases who undergo phacoemulsification through a clear corneal incision and under topical anaesthesia. In this situation conjunctival

flaps and anaesthetic myotoxicity are eliminated.

Though ocular deviations persisted in 6 cases only 3 of them had diplopia which again is inconsistent with their assumed myological origin.

The authors have concluded that incidence of myotoxicity decreases by the end of 12 weeks whereas in their study 75% of the vertical deviations, 66.6% combined deviations and 60% of "ptosis and deviations" persisted. Only horizontal deviations recovered significantly.

Maj RAKESH MAGGON

Gd Spl (Ophthalmology), 153 General Hospital, C/o 56 APO.

REPLY

In 12 patients who underwent repeat blocks, 2 patients developed post-operative ptosis and one developed an ocular deviation, which resolved by 12 weeks. In none of the earlier studies, there is any mention of an association of repeat blocks with ptosis. Only 4 case reports have been published of deviations after cataract surgery with repeat blocks. Moreover in these blocks, adrenaline was used along with bupivacaine (0.75%) which potentiates the effect of the anaesthetic. Recent studies indicate that there may be some degree of direct trauma to the muscle with the needle and some amount of intramuscular injection of LA would lead to more muscle damage and scarring with subsequent ocular deviation.

Out of the 6 patients with ocular deviation, only 3 had diplopia. Three patients had significant lenticular opacities in the non-operated eye with vision <6/60 which could explain the absence of diplopia in these cases.

As seen in the table, if we take deviations alone, there were 10 at the end of first week which had decreased to 3 at the end of XII week. But as pointed out, deviations associated with ptosis as well as vertical deviations have a slower recovery and need careful

follow-up for a longer duration. Moreover in vertical deviations the effect of LA could be compounded by direct surgical trauma to the muscles itself and these cases may need surgical intervention later.

If there are any further queries, I would be glad to answer them.

REFERENCES

1. Esswein MB, Noorden GKV. Paresis of a vertical rectus muscle after cataract extraction. *Am J Ophthalmol* 1993; 116 : 424-30.
2. Capo H, Roth E, Johnson T, et al. Vertical Strabismus after cataract surgery. *Ophthalmology* 1996; 103, 918-21.
3. Capo H, Guyton DL. Ipsilateral hypertropia following cataract surgery. *Ophthalmology* 1996; 103 : 721-30.
4. Burns CL, Seigel LA. Inferior rectus recession for vertical tropia after cataract surgery. *Ophthalmology* 1988; 95 : 1120-4.
5. Hunter DG, Lam GC, Guyton DL. Inferior oblique muscle injury from local anaesthesia for cataract surgery. *Ophthalmology* 1995; 102 : 501-9.

Maj K SHYAM SUNDER

Gd Spl (Ophthalmology), Military Hospital, Jalandhar Cantt.

HIV: CHANGING SCENARIO AND NEW CHALLENGES

Dear Editor,

This is with reference to Editorial "HIV: changing scenario and new challenges" [1] and your reply to the letter "HIV: Scenario in Andaman and Nicobar Islands" [2].

From our unit two HIV positive cases have been reported in last two years. One of them gave history of receiving blood transfusion in Tripura and other gave a history of sexual contact with a prostitute while on temporary duty in Meghalaya. Both are serving soldiers, one from Kerala and other from Uttaranchal.

REFERENCES

1. Sancheti PC, Rai R. Editorial HIV: Changing scenario and new challenges. *MJAFI* 1998;54:91-2.
2. Rai R. HIV: Scenario in Andaman and Nicobar Islands (Reply to letter to editor). *MJAFI* 1999;55:84.

Capt MADHUSUDAN CHEREKAR

RMO, 9 Assam Rif, C/o 99 APO.

ECG DIAGNOSIS : SUPRAVENTRICULAR TACHYCARDIA

Dear Editor,

This 30-year old lady presented with palpitation. Her ECG (Fig-1) revealed regular heart rate of 214/min with cycle length of 280 msec. P-wave morphology was negative in inferior leads and upright in lead I, aVR and aVL with caudocranial activation pattern.

P-R interval (80 msec) was less than fixed R-P interval (200 msec). QRS complexes were narrow with duration of 40 msec. QRS alternans was present in lead V₅ and V₆. She was treated initially with injection deltiagem (Repeated three times) without any response. Three DC shocks of 50,50 and 100 joules had no effect on her

Fig. 1: Electrocardiogram

tachycardia. She was placed on amiodarone drip with which she responded after 6 hours. There was no evidence of preexcitation on her normal ECG.

Regular narrow complex tachycardia in the absence of preexcitation can be due to atrioventricular nodal reentry (AVNRT), atrioventricular reentry with concealed bypass tract (AVRT), sinus node reentrant tachycardia (SNRT), and other atrial tachycardia (AT). Altered P-wave morphology and axis excluded possibility of SNRT and AT. In AVNRT the atria and ventricle are activated simultaneously, thus P-waves are usually buried within QRS complex. Hence P-waves are either not seen or can be noted as continuation of terminal QRS wave as pseudo 's' wave in inferior leads. In AVRT with concealed bypass tract, P-wave must follow the QRS. In 90% of this tachycardia the bypass tract is fast conducting, hence R-P interval is less than P-R interval. But in 10% slow conducting bypass tract produces long R-P and short P-R interval, which is seen in our patient. Rarely uncommon variety of AVNRT, in which retrograde conduction proceeds up the slow pathway and antegrade conduction down the fast pathway, may produce long R-P tachy-

cardia [1]. However tachycardia associated with slow conducting bypass tract are incessant in contrast to long R-P AVNRT, which is virtually always paroxysmal. Furthermore the presence of QRS alternans during narrow complex tachycardia had a specificity of 96% and a predictive accuracy of 92% for the concealed bypass tract [2]. Hence clinical profile and ECG feature is consistent with diagnosis of AVRT with slow conducting bypass tract in our case.

REFERENCES

1. Josephson ME, Wellens HJJ. Differential diagnosis of supraventricular tachycardia. *Cardiology Clinics* 1990;8:411-42.
2. Green M, Heddle B, Dassen W. Value of QRS alteration in determining the site of origin of narrow QRS supraventricular tachycardia. *Circulation* 1983;68:368-74.

Lt Col JS DUGAL *, Lt Col MK GARG *, Col AP SINGH *

* Classified Specialist (Medicine) & Cardiologist, * Classified Specialist (Medicine) & Endocrinologist, * Senior Adviser (Medicine & Neurology), Department of Medicine, Command Hospital (SC), Pune 411 040.