1996;49-68.

4. Matovinac J. Complications of goiter prophylaxis. In: Stansbury JB, Hetzel BS, eds. *Endemic Goitre and Endemic Cretinism*, New York, Wiley Med J Publications 1980;533-49.
5. Indian Council of Medical Research. Report of task force on salt consumption pattern in India. ICMR, New Delhi 1996.
6. Joseph P, et al. Iodine intake and thyroid function. *J Molecular Med* 1980;4:21-37.
7. Thilly CH, Hetzel BS. An assessment of prophylactic programmes: Social, political, cultural and economic issues. In: Stansbury JB, Hetzel

BS, eds. *Endemic Goitre and Endemic Cretinism*, New York, Wiley Med J Publications 1980;445-60.

8. Anonymous. Prevention and Control of Iodine Deficiency Disorders. *Lancet* 1986;23:433-4.
9. Pandav CS, Malik A, Anand K, Pandav S, Karmarkar MG. Prevalence of IDD in NCT of Delhi. IDD Delhi Report No 1, 1996.

Maj MP CARIAPPA

Advance Course Trainee (Command Phase), Station Health Organisation (L), Pune 411 040.

STUDY OF GLOMERULAR FUNCTION IN NEONATES

Dear Editor,

This is with reference to the original article on the subject of glomerular function in neonates [1]. The ultimate objective of any research should be to translate the benefits to the betterment of patient care. That neonates have a high plasma creatinine levels at birth and normalizing over period of time is a well-established text book fact. However, the riddle of the high plasma creatinine levels in term and particularly in preterms seems to be solved. Once the umbilical cord is severed, the perfect intrauterine maternal-fetal biochemical balance is disturbed. Thereafter, the already transferred exogenous adult level creatinine will rapidly disappear in the first urine specimens passed by the now autonomous newborn infant. A new steady state is achieved in due time, based on independent neonatal factors. One of the factors is the unusual occurrence of tubular creatinine reabsorption. The authors hypothesize that this latter temporary phenomenon is attributable to back flow of creatinine across leaky immature tubular and vascular structures. With time, maturational renal changes will impose a barrier to creatinine [2]. Secondly by suggesting that there is scanty Indian literature on the subject of renal functions in neonates, are the authors contemplating difference in different racial groups? Narang et al [3] have studied the maturation of neonatal glomerular function as evidenced by serum creatinine and creatinine clearance in 15 preterm small for dates infants (group 1) and compared with values obtained in 15 preterm appropriate for date babies (group 2) on 3rd, 7th and 14th postnatal days. Creatinine clearance was shown to be statistically less ($p < 0.005$ on day 3, $p < 0.001$ on day 7 and $p < 0.01$ on day 14 respectively) in group 1 as compared to group 2, thereby implying slower renal maturation in small for date preterm babies. This highlights the methodology error in the present report [1]. Thirdly

adaptation in neonatology of the once daily concept of aminoglycoside administration has been well studied. Langhendries JP et al [4] have been able to demonstrate after a trial period of single dose amikacin in 37 babies that minimum inhibitory concentration profiles tested in 43 successive bacterial strains remained adequate. A prospective randomized study compared the pharmacokinetics of once daily gentamicin regime in preterms to the conventional twice-daily regime. The observations suggested that once daily dosage regime guarantees initial peak serum concentration above the minimum inhibitory concentration of gram negative bacteria and trough concentration below potentially toxic levels of 2 µg/ml [5].

REFERENCES

1. Wilson CG, Sarkar PK, Mazumdar JPS, Bharadwaj B. Study of glomerular functions in neonates. *Medical Journal Armed Forces India* 1999;55:183-6.
2. Guignard JP, Drukker A. Why do newborn infants have a high plasma creatinine? *Paediatrics* 1999;103(4): e49.
3. Narang A, Bhakoo ON, Majumdar S, Kumar CH. Renal function in SFD and FD preterm babies. *Indian Paediatr* 1993;30:201-5.
4. Langhendries JP, Battisti O, Bertrand JM, Francois A, Kalenga M, Scalais DJ. Adaptation in neonatology of the once daily concept of aminoglycoside administration: Evaluation of a dosing chart for amikacin in an intensive care unit. *Biol Neonate* 1998;74:351-62.
5. Krishnan L, George SA. Gentamicin therapy in preterms: A comparison of two dosage regimens. *Indian Pediatr* 1997;34:1075-80.

Gp Capt TS RAGHU RAMAN

Senior Adviser (Paediatrics), Command Hospital (AF), Bangalore 560 007.

REPLY

We thank the reader for expressing his views, which however, turn out to be his own assumptions but not based on the facts presented in the article. We, the authors, did not postulate any hypothesis that decrease in a serum creatinine in neonatal period is due to backflow of creatinine across leaky immature tubular and vascular structures as quoted by the reader but we stated that it is due to increase in Renal blood flow and redistribution of intrarenal pattern of glomerular blood flow as viewed by Arant BS Jr also [1]. Similarly mentioning the fact by the authors that "there is scanty Indian literature on this subject" is not expected to lead one to think (by any stretch of imagination) that racial differences are being postulated by the authors. The study is descriptive presentation of facts as observed in subgroups of neonates classified as per gestational age. Pointing out this to be a methodological error in the study is not justifiable and unwanted because various other authors [2,3] showed renal functional maturation is gestational age dependent

rather than birth weight dependent though the study by Narang et al as quoted by the reader narrates Renal function in SFD and AFD preterm babies.

Further, once daily dosage regimen of aminoglycoside in term guaranteeing initial peak serum concentration is a well known fact which was not at all countered by the authors. In the study [4] quoted by the reader the workers used Gentamicin in preterm neonates at a dosage of 4 mg/kg/24 hours when serum creatinine was less than 1 mg/dl. We recommended that dosage of aminoglycoside is to be adjusted as per glomerular functions (Serum creatinine and GFR) more so in preterms.

To reassure the reader, the authors feel that the purpose of research is very well served in the study, as it clearly reemphasises and draws the attention of one and all to be extremely cautious in the dosage of drugs to be administered to preterm neonates.

Thus, all the assumptions made by the reader are just unfortunate

as they serve no purpose.

Col CG WILSON

Professor and Head, Department of Paediatrics, Armed Forces Medical College, Pune 411 040.

REFERENCES

1. Arnat BS Jr. Development patterns of renal functional maturation

compared in human neonate. *J Pediatr* 1978; 92:705-12.

2. Feldman 4, Guignard JP. Plasma creatinine in the first month of life. *Arch Dis Child* 1982;57:123-6.
3. Walia AK, Hall JC, Saini AS, et al. Renal functions in small for gestational age newborn infants. *Indian Pediatr* 1983;20:83-5.
4. Krishnan L, George SA. Gentamicin therapy in preterms: A comparison of two dosage regimens. *Indian Pediatr* 1997;34:1075-80.

TRIAGE IN THE MODERN BATTLE FIELD

Dear Editor,

Proper triage and allotment of priority is an important aspect of smooth and efficient medical cover in the battlefield. The situation in a modern battlefield has drastically changed in view of the high intensity conflicts with all modern weapons and the consequent number of casualties. With increasing risks of nuclear, biological and chemical warfare, the problems can be well appreciated. About 40% of the wounded casualties who otherwise survive, die before reaching a medical aid post [1]. It should also be understood that most trauma patients are reasonably stable for the first one hour after injury (Physiological compensation) and their vital organs will regain function if resuscitated within this period (Golden Hour). After this period decompensation and organ failure is progressively more common [2].

The existing system of prioritising the battlefield casualties into 3 groups (P-I: those needing immediate resuscitation and urgent surgery, P-II: those needing early surgery and possible resuscitation, P-III: remaining casualties) needs to be reconsidered in view of the changing battlefield scenario. In order to do maximum good to maximum number of casualties, the following 5 (Five) category triage is suggested, as modified from the text book [3].

- P-I: Life threatening but savable
- P-II: Serious but stable injury
- P-III Non-walking wounded
- P-IV: Walking wounded
- P-V: Life threatening but unsavable

The priority discs for identification can be issued to only the first 2 or 3 categories for sake of simplicity. The remaining two categories (P-IV and P-V) are self identifiable. By using this system of triage, precious time can be saved in managing P-I casualties than on

moribund P-V casualties, keeping in view the golden hour concept.

It may be argued that the existing categorisation into P-I to P-III is simpler and helps in quicker calculations of surgical load, fluids etc. In that case to achieve more realistic figures in calculation, a P-IV categorisation should be included in the existing system for moribund or unsavable casualties (Equivalent to P-V of the above suggested triage). Thus some of the P-I casualties will be diverted to P-IV and help in achieving more realistic figures in calculations and subsequent medical planning.

Although there is ethical issue related to such categorisation, but the tremendous number of casualties generated by modern warfare cannot be ignored. Hence, to achieve more realistic planning figures for prompt management of salvageable casualties, it is suggested that one of the above classification should be adopted.

Views of the editorial board and the readers are welcome to reach a consensus.

REFERENCES

1. Bacelis-Brite JL. Casualty evacuation in an Armor Task Force: A Riddle in Today's Battlefield. *Medical Corps International* 1990;6:8-13.
2. Stene JK, Grande CM and Giesecke A. Shock resuscitation. In: Stene JK and Grande CM, eds. *Trauma Anaesthesia*. Williams and Wilkins, Baltimore, USA 1994:100-32.
3. Atkinson RS, Rushman GB, Davies NJH. Trauma and multiple injuries. In: Lee JA, ed. *Lee's synopsis of Anaesthesia*, 11th edition. Oxford, Butterworth-Heinemann, 1997:835-54.

Lt Col KC KHANDURI

Classified Specialist (Anaesthesiology), Military Hospital, Ranikhet-263645.

HIGH INCIDENCE OF SEVERE PARASITEMIA IN FALCIPARUM MALARIA

Dear Editor,

We are writing to you in order to share our recent experience of a high incidence of severe parasitemia in falciparum malaria that we have been noting, this malaria season.

During the past 105 days, with effect from 1st August, we admitted 111 cases of falciparum malaria, of these 10 patients were found to have more than 10% parasitemia and 3 of these patients had more than 50% parasitemia. Only 2 of these patients expired, one from ARDS and the other from renal failure, although all were severely ill with renal, hepatic and cerebral dysfunction. Exchange transfusion was not performed and quinine resistance has not been

noted so far. All patients with severe parasitemia were local residents and no immunocompromise state was noted. These cases were seen in all age groups and both sexes. It is possible that a new strain has been recently introduced in the region. We would be extremely interested in the views of our colleagues on this phenomena being observed for the first time by us, with such high incidence.

**Lt Col D ROSHA^{*}, Lt Col VK KATARIA⁺,
Brig KK MAUDAR[#]**

^{*} Pulmonologist and critical care specialist, ⁺ Pathologist, [#] Commandant, Military Hospital Namkum, Ranchi, Bihar 834010.