incidence of pneumothorax. The cough reflex is absent and postoperatively there is less active cough. Bronchial anastomoses heal poorly and are often fragile. Hence airway pressures should be kept low during mechanical ventilation.

As a consequence of refined technique and better care, these patients have a long survival rate and they often return to the operating theatre for cardiac catheterisation, angiography and other incidental surgeries such as appendicectomy, herniotomy or dental procedures. A careful assessment and attention to the aspects highlighted above, may aid in successful anaesthetic management of

these cases.

References

1. Dash A. Anaesthesia of a patient with previous heart transplant. *Int Anesthesiol Clin* 1995;33(2):1-9.
2. Black AE. Anaesthesia for a paediatric patient who has had a transplant. *Int Anesthesiol Clin* 1995;33(2):107-23.

Maj DK SREEVASTAVA

Graded Specialist (Anaesthesia), On Study Leave, The Royal Children Hospital, Melbourne, Victoria 3052. Australia.

UTILISING RESIDUAL 2% METHYL CELLULOSE

Dear Editor,

I would like to disseminate through this letter a simple way to utilize leftover 2% Methylcellulose.

Most anterior chamber (AC) intraocular surgeries, mainly cataract removal with intraocular lens implantation, involve use of viscoelastics. Viscoelastics are critically essential to any procedure involving AC manipulations as they help maintain AC, protect corneal endothelium and assist in breaking synechiae, if any. 2% Methylcellulose, though not strictly a viscoelastic, is used most frequently for this purpose and comes pre-filled in a syringe. Almost invariably some of it remains unspent at the end of the surgery and has to be discarded.

I have been autoclaving this residual 2% Methylcellulose and

accumulating it over 3-4 surgical sessions. This viscous transparent 2% solution can be diluted with equal or three-fourth volume of 4% lignocaine, commonly used as topical anaesthetic. The resultant solution is an excellent coupling fluid for application of contact lenses e.g. Goldmann-3-mirror and Abraham's lens.

Not only can one tailor the viscosity of the solution according to one's preference, there is also a distinct advantage of constant topical anaesthesia. This becomes important in cases of laser photocoagulation where prolonged topical anaesthesia is required. Of course, preventing a wastage has its own charm.

Maj RAKESH MAGGON

Graded Specialist (Ophthalmology), 153 General Hospital, C/o 56 APO.

MULTIPLE MENINGIOMA

Dear Editor,

This refers to article 'Multiple Meningiomas' published in *Medical Journal Armed Forces India* 1999;55:263-4.

Firstly the lady was 50 years old as known to us and as seen on CT scan (Fig 1A) of the case report. Secondly the patient had presented with severe headache and vomiting due to raised intracranial pressure even CT Scan (Fig 1B) shows severe hydrocephalus. Lastly I had the opportunity to operate this lady, two of her three

lesions were operated, the one arising from tentorium (not cerebellum) on left, was confirmed histopathologically to be meningioma. The right CP angle tumour was found to be schwannoma on histopathology. The third seen in the falx was too small to be operated, so histopathologically not confirmed. I fail to understand, how this case can be labelled a case of multiple meningioma.

Lt Col PRAKASH SINGH

Command Hospital (NC), C/o 56 APO.

REPLY

Dear Editor,

This refers to article 'Multiple Meningiomas' published in *Medical Journal Armed Forces India* 1999;55:263-4.

The lady was 50-years old mother of a serving soldier, age typed as 28 years is wrong and deeply regretted.

Hydrocephalous was associated due to primary pathology of intracranial SOL's and hence not emphasized.

All the 3 lesions showed similar CT findings and Imaging diag-

nosis of multiple meningioma was offered on the basis of CT imaging as the case report was essentially depicting the imaging aspect. It is after surgery that the lowermost lesion came out to be a schwannoma which shows similar finding on CT as meningioma.

Col HARIQBAL SINGH

Senior Adviser (Radiodiagnosis), Command Hospital, Pune 411 040.

SAY NO TO MMR; EPIDEMIOLOGICAL REASONS

Dear Editor,

There are some practitioners who advocate Measles, Mumps and Rubella (MMR) vaccinations among infants. The epidemiological impact of this in our country can have adverse consequences. The following clinico-epidemiological factors have to be considered.

Measles is a severe disease in infancy and milder in older children and adults, therefore measles vaccine has a well earned place in the Universal Immunisation Programme (UIP).

Mumps on the other hand, is a mild disease in infancy and more

severe during adulthood. Orchitis occurs in about one in four males who develop mumps after puberty and may lead to sterility [1]. In developed countries, 85% of infections occurred in children younger than 15 years prior to widespread childhood immunisation-now disease occurs in young adults, producing epidemics in colleges/workplace [2]. Why should we repeat the mistakes of developed countries? More so as we cannot afford immunisation at puberty/adulthood as advocated by Western medical literature [3]. Having the disease in childhood is preferable to escaping it then and possibly acquiring it in adult life when protection afforded by

vaccine may have been lost [4].

Rubella. Prior to rubella vaccine programmes in developed countries, the peak incidence of the disease was in children 5-14 years of age—now most cases occur in susceptible teenagers and young adults [2]. It is especially important that girls have immunity to rubella during child bearing age, either by contracting the natural disease or by active immunisation. In USA, Australia and Nordic countries, a second dose of MMR is recommended for teenagers for both genders [3]. In our country the priority should be first to protect women of childbearing age and only then to interrupt transmission of rubella by universal immunisation of children (and not the other way round as is being advocated by premature introduction of MMR in UIP). As no programme exists for teenage/adult women immunisation in our country (and not likely in the near future), MMR during infancy should be withheld lest it cause epidemiological shift towards older age group of teenage/adult women leading to increased incidence of congenital rubella syndrome.

Because of above considerations, only measles vaccination should continue during infancy. MMR is like a chess gambit, con-

cealing a lot of dangerous traps and snares [5], which we can ill afford.

References

1. Geddes AM, Bryceson ADM, Thin CN, Mitchell DM. Diseases due to infection. In: Davidson's principles and practice of medicine. Edwards CRW, Bouchier IAD, Haslett C, eds. 17th ed. ELBS 1995:66-189.
2. Maldonado Y. Rubella/Mumps. In: Nelson's text book of paediatrics. Nelson WE, Behrman RE, Kliegman RM, Arvink AM, eds. WB Saunders Co. 1996:871-5.
3. Benenson AS. Control of communicable diseases manual. 16th ed. American Public Health Association. 1015, 15th street NW, Washington DC, 1995.
4. Christie AB. Infectious disease; epidemiology and clinical practice, 3rd ed, Churchill Livingstone 1980.
5. Yudovich Y. The Gambit. Planeta Publishers, Moscow 1989.

Lt Col A Banerjee,

Classified Specialist (Preventive and Social Medicine) and Epidemiologist, Air Force Station, Amla, Dist Betul-460 553.

METASTATIC MALIGNANT MELANOMA BREAST-FNAC DIAGNOSIS

Dear Editor,

A 31-year-old wife of a serving soldier presented with a lump in Lt breast outer upper quadrant of 2 months duration. It measured 10 x 10 cms and was hard non-tender and freely mobile. It was rapidly increasing in size for the past two months. Past history of a surgery for a nodular lesion on planter aspect of right foot was obtained. However, no further clinical/histological details were available. USG revealed a lobulated mixed echoic large mass with irregular necrotic areas within. Multiple small rounded lesions were seen surrounding the large lesion. FNAC of lump revealed a brownish aspirate. Giemsa and H & E stained smears revealed a mixed population of epithelioid cells and spindle cells. Epithelioid cells displayed eccentric large nuclei, prominent macronucleoli and nuclear vacuoles. There was also presence of binucleated and multinucleated tumor giant cells. Intra cellular melanin pigment was noticed in most spindle cells better appreciated in Giemsa stain as a brownish-black pigment. An FNAB diagnosis of metastatic malignant melanoma breast was concluded (Fig-1).

Metastasis to breast from an extra mammary site is uncommon. In a large study they have been reported to constitute 0.4 to 2% of all breast malignancies [1]. Agarwal et al [2] reported an incidence of 7.6% for all malignancies of the breast. Malignant melanoma is the most common tumor metastasizing to the breast [3] followed by lymphoma, lung cancer, ovarian carcinoma, soft tissue sarcoma, gastrointestinal tumours and genitourinary tumours. The most common breast metastasis in men is from prostatic carcinoma [4] and in children Rhabdomyosarcoma [5]. Clinically, a breast metastasis usually presents as rapidly growing single firm mass that is usually moveable, non-tender and located more in the left breast in upper outer quadrant [6]. Skin involvement or nipple discharge is usually absent. Mammographically, metastasis to breast usually presents as a rounded well circumscribed discrete dense mass that mimics a fibro-adenoma. However, they usually show indistinct margins that help distinguish them from benign lesions. Microcalcifications and spiculations are absent.

FNAB has proven to be an excellent method for detecting metastatic tumours in the breast and of confirming clinically and radiologically suspected breast metastasis. We think that FNAB must be

Fig. 1: Mixed population of epithelioid and spindle cells with macronucleoli. Intracellular melanin pigment seen in spindle cells.

the first approach to obtain a correct diagnosis because not only it is faster, safer, cheaper and more reliable, but also at times it will be the only proof available and thus avoid unnecessary mastectomy and ensure appropriate chemotherapy and radiotherapy.

References

1. Eisenberg AJ, Hajdu SI, Wilhelmus JJ. Preoperative aspiration cytology of breast tumours. Acta cytol 1986;30:135-46.
2. Agarwal J, Gupta JK. Cytodiagnosis of metastasis in the breast. A report of eight cases. J Cytol 1996;13:51-4.
3. Sandison AT. Metastatic tumours in the breast. Br J Surg 1959;47:54-8.
4. Nielsen M, Andersen JA, Henriksen FW. Metastasis to the breast from extramammary carcinomas. APMIS 1981;89:251-6.
5. Rogers DA, Lobo TE, Raw BN. Breast malignancy in children. J Paediatr Surg 1994;29:48-51.
6. American Society of Clinical Pathologists. Check sample. Chicago. ASCP 1994:55-61.

Maj T Chatterjee

Classified Specialist (Pathology), Base Hospital, Delhi Cantt 110 010