ADENOCARCINOMA OF SMALL BOWEL

An Unusual Presentation

Lt Col ATUL GUPTA^{*}, Lt Col VIJAY SINGH⁺,
Brig VC OHRI[#]

MJAFI 2000; 56 : 361-362

KEY WORDS : Adenocarcinoma; Small bowel.

Introduction

Adenocarcinoma is a malignant neoplasm arising from the epithelial cell (of endodermal origin) of the lining of the gastrointestinal tract which microscopically shows glandular pattern. In spite of the rapid turnover rate frequency of adenocarcinoma is very low in small bowel though lymphomas are not so uncommon. Usually the patients present late with features of intestinal colic, nausea, vomiting and at times with intestinal obstruction [7]. Sometimes the patients report with advanced disease with distant metastasis carrying poor prognosis. We present a case of adenocarcinoma of ileum which presented as acute on chronic intestinal obstruction, required emergency palliative resection and anastomosis since there were distant metastasis in lymph nodes and liver.

Case Report

A 63 year old man presented with history of recurrent episodes of colicky abdominal pain of one month duration and had nausea, vomiting and constipation for 3 days. General examination and examination of abdomen was non-contributory. Radiographical examination of abdomen revealed few gas filled bowel loops but no free fluid levels. While under conservative treatment for sub acute intestinal obstruction in hospital he developed acute intestinal obstruction on fifth day of his admission and was taken up for emergency exploratory laparotomy. Per-operative findings revealed blood stained peritoneal fluid (approximately 1.5 litres), a stricture approx. 5 cm long in the wall of ileum at the junction of upper 2/3rd and lower 1/3rd, multiple firm to hard lymph nodes adjacent to the bowel segment and in the mesentery and multiple nodular secondaries were found in the right lobe of the liver. Resection and end-to-end anastomosis was done. Postoperative period was complicated with the development of left pleural effusion, ascitis and slow wound healing. He was discharged from the hospital after three weeks of surgery. Histopathological examination of the resected specimen revealed poorly differentiated, infiltrative adenocarcinoma of small intestine; infiltration was seen till subserosa.

Discussion

Small bowel malignancies constitute only 1-2% of

all the malignancies although some rise in incidence has been reported in recent past [1]. The relative immunity of small bowel is probably due to absence of bacteria, an alkaline environment and low levels of activating enzymes of precarcinogens [2]. In industrialized countries, small bowel cancers are predominantly adenocarcinomas, in developing countries, lymphomas are much more common. The risk factors are similar to the risk factor of carcinoma colon and include cigarette smoking, alcohol intake, familial adenomatous polyposis, post cholecystectomy, peptic ulcer disease, cystic fibrosis, Crohn's disease [2] and at times in the ileal conduit [3]. The small bowel may be affected by adenocarcinoma, lymphomas, carcinoid tumour, leiomyosarcoma and rarely by sarcomatoid carcinoma. The frequency of adenocarcinoma is reported to decrease in aboral direction while it increases in carcinoids [4]. Males have higher incidence than females. Ninety percent patients are older than 40 years. Early diagnosis is difficult since the initial symptoms are non-specific and instrumental tests do not allow a differential diagnosis to be made between adenocarcinoma and benign inflammatory bowel disease, benign tumours and other neoplasias. The commonest presenting features include pain abdomen (83%), nausea and vomiting (54%), weight loss (29%) abdominal distension (24%) and guarding and rigidity (10%). Abdominal mass was the most common finding on physical examination [5]. Laboratory examinations are normal in 40% but the radiological findings viz. barium contrast studies are abnormal in majority (87%) while ultrasonography and CT scan are not reliable for diagnosing ill tumours [5,6]. The preoperative diagnosis is possible in only 30-51% cases [1,2]. Few patients present as surgical emergencies-obstruction and perforation [1]. Tumour resection for palliation or cure is possible in 68% cases with low operative mortality [7]. 10 years actuarial survival rate for

^{*}Classified Specialist (Surgery), ⁺Graded Specialist (Pathology), [#]Commandant, 167, Military Hospital, C/O 56 APO.

Fig. 1 : Photomicrograph of small bowel showing normal villous pattern of mucosa with malignant cells. Infiltration with glandular pattern seen in submucosa. (x10, H & E stain)

Fig. 2 : Photomicrograph of small bowel showing infiltration of malignant cell in outer part of muscularis propria and serosa (x 10, H & E stain)

adenocarcinoma is reported to be 24%(all stages) 75% for stage I, 25% for stage II, 0% for stage III [7]. The most common recurrence for adenocarcinoma is peritoneal carcinomatosis [7].

References

1. Severson RK, Schenk M et al. Epidemiol-Increasing Incidence of adenocarcinomas and carcinoid tumours of the small intestine in adults. *Cancer - Epidemiol - Biomarkers* 1996 Feb;5 (2) :81-4.
2. Neught AI, Jacobson JS, Suc J, Mukherjee R, Arber J. The epidemiology of cancer of the small bowel. *Cancer - Epidemiol - Biomarkers* 1998 Mar; 7(3):243-51.
3. Tsuboniwa N, Miki T, Kuroda M et al. Primary adenocarcinoma in ileal conduit. *Int.Journal Urology*, 1996 Jan;3(1):64-6.
4. Gabos S, Berkel J, Band P et al - Small bowel cancer in western Canada. *Int-Journal-Epidemiology*. 1993 Apr;22(2): 198-206.
5. Garcia Marcilla JA, Sanchez Bueno F, Aguilar J et al - Primary small bowel malignant tumours. *Eur J Surg Oncol* 1994 Dec; 20(6):630-4.
6. Kusumoto H, Takahashi I, Yoshida M et al. Primary malig-

Fig. 3 : Photomicrograph of small bowel showing tumour cells arranged in irregular acinar pattern in submucosa

nant tumours of the small intestine:analysis of 40 Japanese patients. *J Surg Oncol* 1992 Jul;50(3):139-43.

7. Frost DB, Mercado PD, Tyrell JS. Small bowel cancer : A 30 years review. *Ann.Surg Oncol*. 1994 Jul;1(4):290-5.