Paediatric traumatic pneumomediastinum: the spinnaker sail sign

Maria João Ferreira, Mafalda Couto, Rita Loureiro, António Horta Oliveira

Serviço de Cirurgia 2, Centro Hospitalar Tondela-Viseu EPE, Viseu, Portugal

Correspondence to
Dr Maria João Ferreira,
mjp_ferreira@hotmail.com

Accepted 29 March 2017

DESCRIPTION

A 20-month-old female infant was admitted to the emergency room after being run over by a car. She presented with pallor and signs of impaired peripheral perfusion, despite being conscious and alert. Signs of respiratory distress were recognised. There was a bilateral decrease of respiratory murmurs at pulmonary auscultation and her heart sounds were also diminished. Her peripheral oxygen saturation was 94% and her respiratory rate was 54 breaths per minute. A cervical and thoracic crepitus were present. There were no alterations at abdominal examination. Her blood pressure was 142/58 mmHg.

She presented with a Glasgow Coma Scale score of 15 and had normal reactive pupils. Facial oedema was observed and an epicranial haematoma was also detected. Haematological workup revealed a haemoglobin value of 12.9 g/dL.

A supine chest radiograph showed left hydropneumothorax, right pneumothorax and pulmonary contusion. This exam also showed significant pneumomediastinum revealing the *spinnaker sail sign*, which consists on the accumulation of air in the mediastinum in a wedge shape, displacing the thymus superiorly and laterally (figure 1). Two chest drains were immediately placed and the infant remained stable. A thoracic CT scan revealed

Figure 2 Thoracic CT scan revealing extensive cervical subcutaneous emphysema, left hydropneumothorax, and right pneumothorax and pneumomediastinum.

extensive cervical subcutaneous emphysema and hydropneumothorax, and showed no evidence of tracheobronchial, oesophageal or vascular injuries (figure 2). There was no need for mechanical ventilatory support and the child showed both clinical and radiological daily improvement. The chest drains were removed at day 3 and she was discharged 7 days after the injury without complaints or any sequelae.

Figure 1 Supine chest radiograph showing left hydropneumothorax, right pneumothorax, pulmonary contusion and significant pneumomediastinum revealing the spinnaker sail sign.

To cite: Ferreira MJ, Couto M, Loureiro R, et al. *BMJ Case Rep* Published Online First: [please include Day Month Year]. doi:10.1136/bcr-2017-219494

Learning points

- ▶ Traumatic pneumomediastinum, although relatively rare, can be associated with potentially lethal tracheobronchial, oesophageal or vascular thoracic injuries.¹
- ▶ When a thoracic CT shows no signs of tracheobronchial, oesophageal or vascular thoracic injuries and the child remains stable, there is no evidence supporting the need for bronchoscopy or oesophageal imaging.² In most of the cases the treatment is conservative.²
- ▶ The spinnaker sail sign happens when the air trapped in the mediastinum displaces the thymic tissue superiorly and laterally, which is similar to the headsail of a boat in appearance.³

Contributors MJF was part of the surgical team that assisted the child during her admission to the emergency room. She was responsible for clinical data acquisition and writing the manuscript. MC was part of the surgical team that assisted the children during her stay at the nursery. She also had an important role when revising the contents of the manuscript. RL played a major role in acquisition of all the clinical data and collecting the figures. A HO designed and developed this clinical case report. He was responsible for revising the content and ensured veracity of the manuscript.

Competing interests None declared.

Patient consent Obtained from guardian.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Bejvan SM, Godwin JD. Pneumomediastinum: old signs and new signs. *AJR Am J Roentgenol* 1996;166:1041–8.
- 2 Pryor SD, Lee LK. Clinical outcomes and diagnostic imaging of pediatric patients with pneumomediastinum secondary to blunt trauma to the chest. *J Trauma* 2011;71:904–8.
- 3 Monteiro R, Paulos L, Agro J, *et al.* Neonatal spontaneous pneumomediastinum and the spinnaker-sail sign. *Einstein* 2015;13:642–3.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- Submit as many cases as you like
- Enjoy fast sympathetic peer review and rapid publication of accepted articles
- Access all the published articles
- Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow