Lingual hyperpigmentation after 5-fluorouracil chemotherapy

Michael Lawrenz Co, Marcus Juan Esteban

Department of Medicine, Rush University Medical Center, Chicago, Illinois, USA

Correspondence to
Dr Michael Lawrenz Co,
michael_co@rush.edu

Accepted 3 April 2017

DESCRIPTION

A 76-year-old woman presented to our hospital with abdominal pain, diarrhoea and new-onset hyperpigmented spots on her tongue. Her medical history includes metastatic colon cancer, acute myeloid leukemia (AML) in remission after bone marrow transplant, hypertension, chronic kidney disease (CKD), osteoarthritis and prior internal jugular deep venous thrombosis (DVT). She had just received second cycle of 5-fluorouracil, bevacizumab and leucovorin 3 days prior to admission. Examination revealed hyperactive bowel sounds, diffuse abdominal tenderness and multiple hyperpigmented papules on her tongue ([figure 1](#)). Upright chest film did not show any gastrointestinal perforation. Abdominal CT scan was consistent with pancolitis. PCR for *Clostridium difficile* toxin was positive. She was started on a 14-day course of vancomycin and metronidazole and was discharged a few days later after her diarrhoea and abdominal pain improved. On follow-up 4 weeks later, the hyperpigmented papules on her tongue were noted to have improved significantly ([figure 2](#)).

The temporal sequence between the development of hyperpigmentation in this patient and her recent chemotherapy suggests a possible association. Indeed, previous studies have reported a similar association between chemotherapeutic drugs and hyperpigmentation.¹⁻⁴ The mechanism behind this association is not clear; however, some have hypothesised that certain chemotherapeutic

Figure 2 Almost complete spontaneous resolution of hyperpigmentation 4 weeks after chemotherapy.

drugs upregulate the activity of melanocyte-stimulating hormone leading to hyperpigmentation.² The previous reports on this association have mentioned spontaneous resolution of the hyperpigmentation after a few weeks to months from chemotherapy infusion. This knowledge on clinical course is important in both the clinic and hospital settings in order to guide further workup including a possible skin biopsy to rule out more serious conditions.

Figure 1 Hyperpigmented papules and plaques on the tongue 3 days after chemotherapy.

Learning points

- ▶ Certain chemotherapeutic agents such as capecitabine, 5-fluorouracil, doxorubicin, and cyclophosphamide are associated with hyperpigmentation most commonly in the tongue, palms and soles that usually spontaneously resolve after a few weeks to months.
- ▶ Biopsy is not usually required for these lesions, although if they fail to improve, a dermatology consult for biopsy may be warranted.
- ▶ 5-Fluorouracil and bevacizumab may be associated with gastrointestinal toxicities including diarrhoea and gastrointestinal perforation.⁵

CrossMark

To cite: Co ML, Esteban MJ.
BMJ Case Rep Published
Online First: [please
include Day Month Year].
doi:10.1136/bcr-2017-
219806

Contributors Both authors were involved in writing up the case and preparing for publication.

Competing interests None declared.

Patient consent Obtained.

Provenance and peer review Not commissioned; externally peer reviewed.

© BMJ Publishing Group Ltd (unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- 1 Vickers MM, Easaw JC. Palmar-plantar hyperpigmentation with capecitabine in adjuvant colon cancer. *J Gastrointest Cancer* 2008;39:141–3.
- 2 Alfreijat M. Tongue hyperpigmentation associated with chemotherapy. *J Community Hosp Intern Med Perspect* 2013;3:21047.
- 3 Sreeja C, Ramakrishnan K, Vijayalakshmi D, *et al*. Oral pigmentation: a review. *J Pharm Bioallied Sci* 2015;7:403–8.
- 4 Marcelo B, Nakhle S. Chemotherapy-induced hyperpigmentation of the tongue. *N Engl J Med* 2011;365:e20.
- 5 Petrelli F, Cabiddu M, Barni S. 5-Fluorouracil or capecitabine in the treatment of advanced colorectal cancer: a pooled-analysis of randomized trials. *Med Oncol* 2012;29:1020–9.

Copyright 2017 BMJ Publishing Group. All rights reserved. For permission to reuse any of this content visit <http://group.bmj.com/group/rights-licensing/permissions>.
BMJ Case Report Fellows may re-use this article for personal use and teaching without any further permission.

Become a Fellow of BMJ Case Reports today and you can:

- ▶ Submit as many cases as you like
- ▶ Enjoy fast sympathetic peer review and rapid publication of accepted articles
- ▶ Access all the published articles
- ▶ Re-use any of the published material for personal use and teaching without further permission

For information on Institutional Fellowships contact consortiasales@bmjgroup.com

Visit casereports.bmj.com for more articles like this and to become a Fellow