Published in final edited form as:

Angew Chem Int Ed Engl. 2017 July 17; 56(30): 8736–8741. doi:10.1002/anie.201703844.

Electronically Activated Organoboron Catalysts for Enantioselective Propargyl Addition to Trifluoromethyl Ketones

Nicholas W. Mszar, Malte S. Mikus, Dr. Sebastian Torker, Dr. Fredrik Haeffner, and Prof. Amir H. Hoveyda*

Department of Chemistry, Merkert Chemistry Center, Boston College, Chestnut Hill, MA 02467 (USA)

Abstract

A broadly applicable, practical, scalable, efficient and highly α - and enantioselective method for addition of a silyl-protected propargyl moiety to trifluoromethyl ketones has been developed. Reactions, promoted by 2.0 mol % of a catalyst that is derived in situ from a readily accessible aminophenol compound at ambient temperature, were complete after only 15 minutes at room temperature. The desired tertiary alcohols were isolated in up to 97% yield and 98.5:1.5 enantiomeric ratio. Alkyl-, alkenyl-, alkynyl-, aryl- or heteroaryl-substituted trifluoromethyl ketones can be used. Utility is highlighted by application to a transformation that is relevant to enantioselective synthesis of BI 653048, a compound active against rheumatoid arthritis.

Fast and selective

When armed with a trifluoromethyl group, readily accessible small-molecule chiral organoboron catalysts can promote highly enantioselective addition of a silyl-protected propargylboron compound to a wide range of trifluoromethyl ketones with high efficiency. Utility is highlighted by demonstrated through a process that may be used for enantioselective synthesis of BI 653048 (active against rheumatoid arthritis).

Keywords

catalysis; enantioselective synthesis; homopropargylic alcohols; propargyl groups; trifluoromethyl group

Organic molecules that contain a trifluoromethyl unit often possess desirable properties,^[1] and efficient and reliable methods for their enantioselective synthesis are needed.^[2] Catalytic enantioselective additions to trifluoromethyl ketones is an attractive way to synthesize trifluoromethyl-substituted tertiary alcohols,^[3, 4] but developing such transformations is hardly straightforward.^[5] One issue is the high electrophilicity of trifluoromethyl ketones and rapid competitive non-catalyzed/non-enantioselective background reactions. Another problem is the smaller size difference between a trifluoromethyl group and the other carbonyl substituent (more so than methyl ketones), rendering enantiotopic face differentiation based on steric factors less tenable. As part of a program designed to address these general issues, we have developed an ammonium containing boron-based chiral catalyst capable of promoting enantioselective additions of allyl- and allenyl-boron compounds to trifluoromethyl ketones.^[6] High enantioselectivity in these reactions arises from transition state organization induced through electrostatic attraction (**I**, Scheme 1).

A range of aminophenol-based catalysts can be synthesized easily. For instance, as supported by mechanistic investigations and computational studies,^[7] H-bonding interactions lead to enantioselective organoboron additions to phosphinoylimines, isatins (e.g., **II**, Scheme 1) and Boc-protected aldimines.^[8] In the additions to ketones, where there is just one point of catalyst-substrate contact (i.e., no electrostatic attraction with the ammonium group), high e.r. arises from a more sizeable triphenylsilyl moiety in the catalyst (**III**), which helps prevent the larger ketone substituent to orient pseudo-axially.^[9] With Zn(OMe)_2 as the co-catalyst, reactivity can be enhanced and/or alternative pathways made available.^[10] We have proposed that, among other factors, chelation of the Lewis acid to phenoxy oxygen (**V**) destabilizes an inactive form of the chiral catalyst, increasing boron center Lewis acidity and accelerating 1,3-borotropic shift so that it occurs prior to the addition step (i.e., net γ vs. α selectivity).^[10]

The present studies were inspired by the complications associated with enantioselective synthesis of glucocorticoid agonist BI 653048, developed for the treatment of rheumatoid arthritis (Scheme 2a).^[11e,12] Towards this end, Fandrick *et al.* have reported what is, as far as we know, the only available method for catalytic enantioselective propargyl group addition to trifluoromethyl ketones.^[12] Homopropargyl alcohol **4** was synthesized by reaction of ketone **1** with silyl-protected propargyl-B(pin) compound **2**^[13,14] promoted by a catalyst generated *in situ* from 27 mol % *N*-iso-propyl-L-proline (**3**) and 25 mol % Et_2Zn (Scheme 2a). After two days at -40°C , tertiary alcohol **4** was isolated in 78% yield, 90:10 diastereomeric ratio and >98:2 propargyl:allenyl selectivity. We wondered if with an appropriate aminophenol-based organoboron catalyst, products such as **6** (Scheme 2b; precursor to BI 653048),^[12] might be synthesized more efficiently (e.g., <4 h), with higher enantioselectivity (e.g., 95:5 e.r.), at room temperature (vs. -40°C), and with lower

catalyst loading (e.g., <5.0 vs. 25 mol %). It has been demonstrated that the silyl-alkyne moiety, in its protected or unmasked form, can be converted to several desirable derivatives, including those not easily accessible by modification of the corresponding homoallylic alcohols.^[13c,15] Another aim was therefore to develop a method that would be applicable to an assortment of trifluoromethyl ketones.

Preliminary studies were performed with ketone **5**, which may be used to access BI 653048. Control experiments showed ~15% conversion in the absence of a chiral organoboron catalyst (conv. to **6**, Scheme 3). We were also met with a more significant challenge: silyl-protected propargyl-B(pin) **2** undergoes complete proto-deboration within 15 minutes at 22 °C (Scheme 3; >98% by ¹H NMR analysis).^[16] A catalyst would have to be identified for bringing the transformation to completion within the same brief length of time.

There was appreciable enantioselectivity (88:12 e.r.) with 2.0 mol % **7a**, but the yield was low (38% yield; Scheme 3). Selectivity improved to 92:8 e.r. with triphenylsilyl-substituted **7b**, but there was little change in efficiency. Based on the reasoning that a smaller aryloxy group might better accommodate the large alkyl moiety of **5** (see complex **VI**, Scheme 2), reactions with **7c** and **7d** were probed; efficiency did improve (87% and 84% conv., 61% and 60% yield, respectively) and there was a boost in e.r. (97:3). We attributed the enantioselectivity increase to the ability of the smaller catalyst to compete more effectively with the non-catalytic pathway. There was complete α selectivity in all cases (i.e., >98:2 propargyl:allenyl addition).

The key question then was whether further catalyst activation could be achieved electronically, as represented in **VII**. We wondered if, while bulkier than **7c** and **7d**, an aminophenol ligand containing an electron-withdrawing substituent on its aryloxy unit (e.g., vs. Me in **7c**) could provide appropriate rate enhancement without adversely impacting e.r. We prepared trifluoromethyl-substituted aminophenol ligand **7e** (from commercially available 3-trifluoromethyl salicylaldehyde) according to the logic that a trifluoromethyl differs slightly from a methyl group (similar in size to an Et unit^[17]). Indeed, after 15 minutes at room temperature with 2.0 mol % **7e**, **6** was isolated in 94% yield, >98% α selectivity and 98.5:1.5 e.r. Electronic activation of the catalyst not only led to considerable increase in efficiency, it resulted in some enhancement in e.r. as well. To obtain the enantiomer for synthesis of the biologically active enantiomer of BI 653048,^[11e,12] we used the aminophenol ligand derived from D-valine and obtained *ent*-**6** in nearly the same yield and e.r. (Scheme 3).^[18] The aryl bromide moiety in **6** was readily converted to the corresponding carboxylic acid, ready for use for synthesis of the target molecule, by a one-pot procedure (*i*PrMgClO₂LiCl, -78-22 °C, 2 h; CO₂, 22 °C, 6 h; 89% yield).^[16]

Yields and enantioselectivities were high for the additions to ketones with smaller alkyl substituents (vs. the moiety in **5**), for which enantiotopic differentiation is more challenging; only one moderately selective example was previously disclosed (80:20 e.r.).^[12] Tertiary alcohols **8a-c**, derived from additions to a ketone with an α -branched, β -branched or an *n*-alkyl substituent, respectively, were isolated in 62–89% yield and 95.5–97.5:2.5 e.r. Formation of enyne **9** was similarly efficient and enantioselective (88% yield, 98.5:1.5 e.r.). Diyne **10** was obtained in lower enantiomeric purity (76% yield, 86:14 e.r.) perhaps because

of the increased steric repulsion between the catalyst's CF₃ group and the conformationally less mobile phenylacetylene moiety (see **VI**, Scheme 2). There was no improvement in enantioselectivity at higher catalyst loading.

Aryl- and heteroaryl-substituted trifluoromethyl ketones were similarly suitable (Scheme 5), despite background processes being more competitive (~30–60% conv. without **7e**). Reactions of trifluoromethyl-substituted ketones with an aromatic moiety that contains either an electron-withdrawing or -donating group proceeded to >90% conversion within 15 minutes, affording **11a–m** in 85–97% yield and 92:8–98.5:1.5 e.r.^[16] As indicated by efficient preparation of pyrrole-substituted **11l**, protection of basic amines was not required.

Nonetheless, *o*-fluorophenyl substituted alcohol **11n** (86:14 e.r.) and most notably *o*-chloro- and *o*-bromophenyl derivatives **11o** and **11p** (56:44 e.r.) were formed in markedly lower e.r. This was surprising for two reasons: 1) Compounds **11b** and **11c** (Scheme 5), derived from additions to the related *o*-tolyl- and *o*-methoxy-substituted ketones, were generated with considerably higher e.r. (92:8 and 97:3, respectively). 2) Allyl-B(pin) additions to methyl ketones bearing the same *o*-halo-substituted aryl groups were much more enantioselective with the same class of catalysts (93:7–99:1 e.r.).^[9]

The above data suggest that steric repulsion between an *ortho* substituent and the catalyst structure can cause diminution in enantioselectivity (e.g., **11a–c**, Scheme 5); this explains why e.r. is more favorable (85:15 vs. 59:41) with a smaller fluorine atom and a shorter C–F bond in **11n** compared to a bromine atom and a longer C–Br bond in **11p**. Electronic factors probably play a role as well. In what we suggest as the preferred mode of addition (**VIII**, Scheme 6) there would be some degree of electron-electron repulsion between the non-bonding electrons of the halogen atom and the catalyst's phenolic oxygen. Unlike a methyl ketone, the alternative conformer **IX** cannot offer relief as a result repulsive interactions with the substrate's trifluoromethyl group. Consequently, competitive pathways become more variable and enantioselectivity suffers. *o*-Methoxyphenyl- and *o*-fluorophenyl-substituted **11c** and **11n** can be generated in higher e.r. because of the better overlap involving period two elements and the resulting resonance stabilization that can reduce electron-electron repulsion (see **X**, Scheme 6).^[19] The above trends underscore key electrostatic interactions that impact enantioselectivity and are particular to trifluoromethyl ketones.

In summary, we have developed the first broadly applicable method for efficient and enantioselective addition of a propargyl group to trifluoromethyl ketones. The chiral ligand and the organoboron reagent can be prepared in gram quantities from commercially available materials. Reactions may be conveniently performed at 22 °C, are complete in just 15 minutes with 2.0 mol % catalyst and can be performed on gram scale, as the examples in Scheme 7 show.

Development of additional enantioselective processes catalyzed by small-molecule organoboron chiral catalysts is in progress.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was provided by the NIH (GM-57212). We are grateful to Kyung A Lee and Farid W. van der Mei for helpful discussions.

References

1. For example, see: Berger R, Resnati G, Metrangolo P, Weber E, Hulliger J. *Chem Soc Rev.* 2011; 40:3496–3508. [PubMed: 21448484] Gillis EP, Eastman KJ, Hill MD, Donnelly DJ, Meanwell NA. *J Med Chem.* 2015; 58:8315–8359. [PubMed: 26200936] Fujiwara T, O'Hagan D. *J Fluorine Chem.* 2014; 167:16–29.
2. For selected reviews, see: Shibata N, Mizuta S, Kawai H. *Tetrahedron: Asymmetry.* 2008; 19:2633–2644. Nie J, Guo HC, Cahard D, Ma JA. *Chem Rev.* 2011; 111:455–529. [PubMed: 21117644] Yang X, Wu T, Phipps RJ, Toste FD. *Chem Rev.* 2015; 115:826–870. [PubMed: 25337896]
3. For enantioselective synthesis of tertiary alcohols through additions to ketones, see: Yus M, González-Gómez JC, Foubelo F. *Chem Rev.* 2011; 111:7774–7854. [PubMed: 21923136]
4. For examples of catalytic enantioselective additions of C-based nucleophiles to trifluoromethyl ketones (with no secondary chelating group), see: Zhang GW, Meng W, Ma H, Nie J, Zhang WQ, Ma JA. *Angew Chem Int Ed.* 2011; 50:3538–3542. Motoki R, Kanai M, Shibasaki M. *Org Lett.* 2007; 9:2997–3000. [PubMed: 17629284] Cook AM, Wolf C. *Angew Chem Int Ed.* 2016; 55:2929–2933. Noda H, Amemiya F, Weidner K, Kumagai N, Shibasaki M. *Chem Sci.* 2017; 8:3260–3269. [PubMed: 28553529] Zheng Y, Tan Y, Harms K, Marsch M, Riedel R, Zhang L, Meggers E. *J Am Chem Soc.* 2017; 139:4322–4325. [PubMed: 28290685]
5. For example, see: Zhang X, Chen D, Liu X, Feng X. *J Org Chem.* 2007; 72:5227–5233. [PubMed: 17559275] Loh TP, Zhou JR, Li XR. *Tetrahedron Lett.* 1999; 40:9333–9336. Haddad TD, Hirayama LC, Taynton P, Singaram B. *Tetrahedron Lett.* 2008; 49:508–511.
6. Lee K, Silverio D, Torker S, Robbins DW, Haefner F, van der Mei FW, Hoveyda AH. *Nat Chem.* 2016; 8:768–777. [PubMed: 27442282]
7. Silverio DL, Torker S, Pilyugina T, Vieira EM, Snapper ML, Haefner F, Hoveyda AH. *Nature.* 2013; 494:216–221. [PubMed: 23407537]
8. Wu H, Haefner F, Hoveyda AH. *J Am Chem Soc.* 2014; 136:3780–3783. [PubMed: 24588835]
9. Robbins DW, Lee K, Silverio DL, Volkov A, Torker S, Hoveyda AH. *Angew Chem Int Ed.* 2016; 55:9610–9614.
10. van der Mei FW, Miyamoto H, Silverio DL, Hoveyda AH. *Angew Chem Int Ed.* 2016; 55:4701–4706.
11. a) Betageri R, Zhang Y, Zindell RM, Kuzmich D, Kirrane TM, Bentzien J, Cardozo M, Capolino AJ, Fadra TN, Nelson RM, Paw Z, Shih DT, Shih CK, Zuvela-Jelaska L, Nabozny G, Thomson DS. *Bioorg Med Chem Lett.* 2005; 15:4761–4769. [PubMed: 16112571] b) Bekkali, Y., Betageri, R., Emmanuel, MJ., Hammach, A., Harcken, HJJ., Kirrane, TM., Kuzmich, D., Lee, TW-h, Liu, P., Patel, UR., Razavi, H., Reither, D., Takahashi, H., Thomson, DS., Wang, J., Zindell, R., Proudfoot, JR. *PCT Int Appl. WO 2005030213.* 2005. c) Betageri, R., Bosanac, T., Burke, MJ., Harcken, C., Kim, S., Kuzmich, D., Lee, TW-H., Li, Z., Liu, Z., Pingrong, L., Lord, J., Razavi, H., Reeves, JT., Thomson, D. *PCT Int Appl. WO 2009149139.* 2009. d) Betageri R, Gilmore T, Kuzmich D, Kirrane TM, Bentzien J, Wiedenmayer D, Bekkali Y, Regan J, Berry A, Latli B, Kukulka AJ, Fadra TN, Nelson RM, Goldrick S, Zuvela-Jelaska L, Souza D, Pelletier J, Dinallo R, Panzenbeck M, Torcellini C, Lee H, Pack E, Harcken C, Nabozny G, Thomson DS. *Bioorg Med Chem Lett.* 2011; 21:6842–6851. [PubMed: 21963986] e) Reeves JT, Fandrick DR, Tan Z, Song JJ, Rodriguez S, Qu B, Kim S, Niemeier O, Li Z, Byrne D, Campbell S, Chitroda A, DeCroos P, Fachinger T, Fuchs V, Gonnella NC, Grinberg N, Haddad N, Jäger B, Lee H, Lorenz JC, Ma S, Narayanan BA, Nummy LJ, Premasiri A, Roschangar F, Sarvestani M, Shen S, Spinelli E, Sun X, Varsolona RJ, Yee N, Senanayake CH. *J Org Chem.* 2013; 78:3616–3635. [PubMed: 23544738]
12. Fandrick DR, Reeves JT, Bakonyi JM, Nyalapatla PR, Tan Z, Niemeier O, Akalay D, Fandrick KR, Wohlleben W, Ollenberger S, Song JJ, Sun X, Qu B, Haddad N, Sanyal S, Shen S, Ma S, Byrne D, Chitroda A, Fuchs V, Narayanan BA, Grinberg N, Lee H, Yee N, Brenner M, Senanayake CH. *J Org Chem.* 2013; 78:3592–3615. [PubMed: 23544787]

13. Fandrick DR, Roschangar F, Kim C, Hahm BJ, Cha MH, Kim HY, Yoo G, Kim T, Reeves JT, Song JJ, Tan Z, Qu B, Haddad N, Shen S, Grinberg N, Lee H, Yee N, Senanayake CH. *Org Process Res Dev.* 2012; 16:1131–1140. For previous reports from these laboratories involving catalytic enantioselective reactions with the reagent 2, see: Mszar NW, Haeffner F, Hoveyda AH. *J Am Chem Soc.* 2014; 136:3362–3365. [PubMed: 24555471] Shi Y, Jung B, Torker S, Hoveyda AH. *J Am Chem Soc.* 2015; 137:8948–8964. [PubMed: 26172476]
14. a) Hoffman RW, Brinkmann H, Frenking G. *Chem Ber.* 1990; 123:2387–2394. b) Fandrick, DR., Reeves, JT., Song, JJ. International Patent. WO 2010/141328 A2. 2010. c) Fandrick DR, Roschangar F, Kim C, Hahm BJ, Cha MH, Kim HY, Yoo G, Kim T, Reeves JT, Song JJ, Tan Z, Qu B, Haddad N, Shen S, Grinberg N, Lee H, Yee N, Senanayake CH. *Org Process Res Dev.* 2012; 16:1131–1140.
15. Vieira EM, Haeffner F, Snapper ML, Hoveyda AH. *Angew Chem Int Ed.* 2012; 51:6618–6621.
16. See the Supporting Information for details. Protodeboration probably occurs through formation of the derived methoxy borate followed by the rupture of C-B(pin) bond by the proton counter-ion.
17. Jagodzinska M, Huguenot F, Candiani G, Zanda M. *Chem Med Chem.* 2009; 4:49–51. [PubMed: 19065645]
18. Consistent with a previous report (see ref. [6]) additions to mono- and difluoromethylphenyl ketone were considerably less enantioselective (with 7e under conditions shown in Scheme 3: 95% conv., 76% yield, >98:2 propargyl:allenyl, 50:50 e.r. and >98% conv., 94% yield, >98:2 propargyl:allenyl, 86:14 e.r., respectively). See the Supporting Information for details.
19. It is unlikely that the enantioselectivity profile for *o*-halophenyl-substituted trifluoromethyl ketones (11n-p) is due to differences in rates of competitive un-catalyzed background additions (i.e., less in the case of the more enantioselective *o*-fluorophenyl substrate). In fact, control experiments point to the opposite trend: 61%, 40% and 29% conversion was observed for 11n-p, respectively, under the same conditions but in the absence of 2.0 mol % 7e.

High enantioselectivity and reactivity with chiral boron-based catalysts

Scheme 1.

Readily modifiable small-molecule chiral aminophenol-derived catalysts allow for mechanism-based optimization of reaction efficiency and/or enantioselectivity.

a. Previous work (Fandrick, *et al.*):

b. Objectives of this study:

Can the catalytic method be extended to a broader range of trifluoromethyl ketones?

Scheme 2.

A problem in catalytic enantioselective synthesis and the objectives of this study.

Key initial findings:
 1. There is ~15% conv. to *rac*-**6** in the absence of a chiral ligand.
 2. Propargyl-B(pin) **2** decomposes within 15 min under the reaction conditions.

Scheme 3.

Identification of an optimal catalyst for enantioselective synthesis of homopropargyl alcohol **6**. Conversion ($\pm 2\%$) determined by analysis of the ¹⁹F NMR spectra of unpurified product mixtures (trifluoro-toluene as reference); $>98\%$ α selectivity in all cases. E.r. ($\pm 1\%$) was determined by HPLC analysis. See the Supporting Information for details. Abbreviation: EWG = electron-withdrawing group; L = methoxy or silyl-allenyl group.

Scheme 4.

Additions to alkyl-, alkenyl- and alkynyl-substituted trifluoromethyl ketones. Conversion ($\pm 2\%$) determined by analysis of the ^{19}F NMR spectra of unpurified product mixtures (trifluoro-toluene as reference); $>98\%$ α selectivity in all cases. E.r. ($\pm 1\%$) determined by HPLC analysis. See the Supporting Information for details.

Scheme 5.

Additions to aryl- and heteroaryl-substituted ketones. Conversion ($\pm 2\%$) determined by analysis of the ^{19}F NMR spectra of unpurified product mixtures (trifluoro-toluene as reference); >98% α selectivity in all cases. E.r. ($\pm 1\%$) determined by HPLC analysis. See the Supporting Information for details.

Scheme 6.

The unusual trends in enantioselectivity in propargyl group additions to *o*-halophenyl-substituted trifluoromethyl ketones. Conversion ($\pm 2\%$) determined by analysis of the ^{19}F NMR spectra of the unpurified product mixtures; $>98\%$ α selectivity in all cases. E.r. ($\pm 1\%$) was determined by HPLC analysis. See the Supporting Information for details.

Scheme 7.

Gram scale. Conversion ($\pm 2\%$) determined by analysis of the ^{19}F NMR spectra of the unpurified product mixtures; >98% α selectivity in all cases. E.r. ($\pm 1\%$) was determined by HPLC analysis. See the Supporting Information for details.