Published in final edited form as:

Brain Behav Immun. 2017 October ; 65: 262–273. doi:10.1016/j.bbi.2017.05.012.

Intracranial IL-17A overexpression decreases cerebral amyloid angiopathy by upregulation of ABCA1 in an animal model of Alzheimer's disease

Junling Yang¹, Jinghong Kou¹, Robert Lalonde², and Ken-ichiro Fukuchi¹

¹Department of Cancer Biology and Pharmacology, University of Illinois College of Medicine at Peoria, P.O. Box 1649, Peoria, Illinois, 61656, USA

²Department of Psychology, University of Rouen, 76821 Mont-Saint-Aignan, Rouen, France

Abstract

Neuroinflammation is a pervasive feature of Alzheimer's disease (AD) and characterized by activated microglia, increased proinflammatory cytokines and/or infiltrating immune cells. T helper 17 (Th17) cells are found in AD brain parenchyma and interleukin-17A (IL-17A) is identified around deposits of aggregated amyloid β protein ($A\beta$). However, the role of IL-17A in AD pathogenesis remains elusive. We overexpressed IL-17A in an AD mouse model via recombinant adeno-associated virus serotype 5 (rAAV5)-mediated intracranial gene delivery. AD model mice subjected to injection of a vehicle (PBS) or rAAV5 carrying the lacZ gene served as controls. IL-17A did not exacerbate neuroinflammation in IL-17A-overexpressing mice. We found that IL-17A overexpression markedly improved glucose metabolism, decreased soluble $A\beta$ levels in the hippocampus and cerebrospinal fluid, drastically reduced cerebral amyloid angiopathy, and modestly but significantly improved anxiety and learning deficits. Moreover, the ATP-binding cassette subfamily A member 1 (ABCA1), which can transport $A\beta$ from the brain into the blood circulation, significantly increased in IL-17A-overexpressing mice. In vitro treatment of brain endothelial bEnd.3 cells with IL-17A induced a dose-dependent increase in protein expression of ABCA1 through ERK activation. Our study suggests that IL-17A may decrease $A\beta$ levels in the brain by upregulating ABCA1 in blood-brain barrier endothelial cells.

Keywords

Alzheimer's disease; ATP-binding cassette transporters; blood-brain barrier; cerebral amyloid angiopathy; $A\beta$; interleukin 17; mouse behavior; ERK signaling; endothelial cells

Address correspondence to: Ken-ichiro Fukuchi, MD, PhD, Department of Cancer Biology and Pharmacology, University of Illinois College of Medicine at Peoria, P.O. Box 1649, Peoria, Illinois 61656 USA; kfukuchi@uic.edu.

Conflict of interest

The authors declare no conflict of interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

1. Introduction

Alzheimer's disease (AD) is the most prevalent form of dementia (Fratiglioni *et al.*, 2000; Kawas *et al.*, 2000; Kukull *et al.*, 2002), characterized by extracellular amyloid β ($A\beta$) deposition in the brain parenchyma and cerebral blood vessels. Additionally, neuroinflammation is a consistent feature of AD and thought to influence the disease pathogenesis (McManus *et al.*, 2014). Immune responses occurring in brains of AD patients can be driven by resident immune cells and/or brain infiltrating peripheral immunocompetent cells (Ghosh and Geahlen, 2015; Togo *et al.*, 2002). Microglia, the innate immune cells in the central nervous system (CNS), are activated by $A\beta$ deposits and secrete inflammatory cytokines and chemokines, contributing to AD progression (Hickman *et al.*, 2008; Ridolfi *et al.*, 2013; Rodriguez *et al.*, 2016).

Central-memory $CD4^+$ T lymphocytes account for more than 80% of the cells in the cerebrospinal fluid (CSF) and routinely penetrate the CNS. Lymphocytes in the CNS are replenished by newly immigrating cells approximately twice per day (Engelhardt and Ransohoff, 2005; Hickey, 2001; Smolders *et al.*, 2013). Numerous lines of evidence indicate the presence of T cells in the brains of AD patients (Ferretti *et al.*, 2016; Lombardi *et al.*, 1999; Monsonego *et al.*, 2003; Shalit *et al.*, 1995; Togo *et al.*, 2002) although their possible roles in AD pathogenesis have been studied less than those of microglia. T cells in the brains of AD patients increased as compared with subjects with non-AD degenerative dementias and age-matched controls (Lueg *et al.*, 2015; Togo *et al.*, 2002). T cells breaching the brain vasculature and parenchyma can be either detrimental or beneficial for brain homeostasis (Kipnis *et al.*, 2004; Monsonego *et al.*, 2006; Ziv *et al.*, 2006). Depletion of regulatory T (Treg) cells in an AD mouse model accelerated the onset of cognitive deficits without altering brain $A\beta$ load (Dansokho *et al.*, 2016). Administration of $A\beta$ -specific type 1 T helper (Th1) cells into the brain lateral ventricle of an AD mouse model enhanced removal of cerebral $A\beta$ deposits and promoted neurogenesis without harmful effects (Fisher *et al.*, 2014). Thus, these T cells play beneficial roles in the AD pathophysiology. In contrast, intravenous administration of $A\beta$ -specific Th1 cells caused brain infiltration, increased microglial activation and $A\beta$ deposition, and impaired cognitive function in an AD mouse model but similarly administered $A\beta$ -specific IL-17-producing helper T (Th17) cells infiltrated into the brain without exacerbating microglia activation, $A\beta$ deposition and cognitive deficits in the same mice (Browne *et al.*, 2013).

IL-17A or IL-17 is a signature cytokine of Th17 cells and is involved in antimicrobial host defense and inflammation. A plethora of evidence supports the notion that Th17 cells and IL-17A play a pathogenetic role in certain autoimmune diseases such as multiple sclerosis, rheumatoid arthritis, psoriasis, and inflammatory bowel disease. Respiratory infection with *Bordetella pertussis* in an AD mouse model (McManus *et al.*, 2014) and intrahippocampal administration of $A\beta_{42}$ into rats (Zhang *et al.*, 2013) induced infiltration of Th17 cells and expression of IL-17A in the brain, which were associated with exacerbation of an AD-like phenotype in these experimental animals. It is interesting to note that plasma IL-17A levels have been identified as one of the plasma biomarkers for AD diagnosis and neocortical $A\beta$ load (Burnham *et al.*, 2014; Doecke *et al.*, 2012). However, the role of IL-17A in AD pathophysiology is unclear. In this study, we investigated the effects of IL-17A

overexpression via intracranial delivery of recombinant adeno-associated virus serotype 5 (rAAV5)-mediated IL-17A on A β load, microglia activation, glucose metabolism and behavioral function using a transgenic animal model of AD.

2. Materials and Methods

2.1. Experimental animals and intracranial injection of rAAV-IL-17A

A rAAV vector, rAAV5-IL-17A, was prepared as described previously (Yang *et al.*, 2016). A control rAAV vector encoding LacZ, rAAV5-LacZ, was similarly prepared. B6. Cg-Tg (APPswe, PSEN1dE9) 85Dbo/J mice (TgAPPswe/PS1dE9 mice) purchased from Jackson Laboratory, were used as an AD mouse model to study a potential role of IL-17A in the AD pathogenesis. For intracranial injection, 2-month-old TgAPPswe/PS1dE9 mice were anesthetized by Nembutal® (pentobarbital) and placed on a stereotaxic instrument with a motorized stereotaxic injector (Stoelting, Wood Dale, IL). A midline incision was made to expose the bregma. A hole in the skull was made by a drill 0.5 mm posterior to the bregma and 1.0 mm right to the midline. rAAV5-IL-17A (1.5×10^{10} vector genomes (vg) in 10 μ l PBS/mouse) was injected unilaterally into the right lateral brain ventricle at the depth of 2 mm at a rate of 1 μ l/min. Mice subjected to rAAV5-IL-17A injection are referred to as IL-17A-overexpressing mice (n=8). Mice similarly treated with 10 μ l PBS or rAAV5-LacZ (1.5×10^{10} vg/mouse) served as controls and are referred to as control mice (n=9): Four PBS-injected and five rAAV5-LacZ-injected mice were combined together because no phenotypic differences between two controls were found (more information in Supplementary Figure 1). Due to the difference in brain amyloid content between male and female TgAPP/PS1 mice (Akhter *et al.*, 2015; Gallagher *et al.*, 2013; Sierksma *et al.*, 2013; Taniuchi *et al.*, 2007; Wang *et al.*, 2003), we used only male mice for this study. All animal protocols were prospectively approved by the Institutional Animal Care and Use Committee of the University of Illinois College of Medicine at Peoria.

2.2. Behavioral tests

2.2.1. Behavioral schedule and statistics—A battery of behavioral tests was performed on experimental mice at 12 months of age as previously described (Lim *et al.*, 2011). After measuring body weight and adapting mice to handling, the tests were conducted over a 16-day period as follows: spontaneous alternation (days 1–10), open-field (days 6–8), elevated plus-maze (days 9 and 10), and Morris water maze (days 11–16). In each test, whenever possible, the apparatus was wiped clean with a wet cloth and dried before the next mouse was introduced to minimize odor cues. For T-maze, Mann-Whitney U test was used to assess the chance of alternation compared with theory (50%) and unpaired t-test was used for the intergroup differences. Intergroup differences were assessed by 2×3 analysis of variance (ANOVA) for 2 independent groups and 3 days of testing with repeated measures on the second factor for the open-field and 2×2 ANOVA for two groups and 2 days of testing for the elevated plus-maze. The unpaired t-test was used for the probe test of the Morris water maze and a 2×5 ANOVA for 2 independent groups and 5 days of testing with repeated measures on the second factor was used for 5-day acquisition of the Morris water maze. Results are expressed as mean $\pm$ standard error of the mean (SEM). In all cases, $P < 0.05$ was considered to be significant.

2.2.2. Exploration and anxiety—Spontaneous alternation was measured in a T-maze, made of white acrylic and consisting of a central stem flanked on each side by 2 arms. The maze width was 9 cm, the wall height 20 cm, and each arm 30 cm in length. On the initial trial, the mice were placed in the stem with the right arm blocked by a plastic barrier (forced choice). After entering the available arm, the mice were kept in it for 1 min by closing the barrier behind them. The mice were then retrieved and after removing the barrier, placed back in the stem for a free-choice trial, either into the same arm or the opposite arm (4-paw criterion). On the following 10 days, the same 2-trial procedure was repeated, except that the blocked arm was switched from right on odd days to left on even days.

The number of alternations and the latencies before responding during the choice trial were measured. In the absence of any decision within 1 min, the mice were briefly prodded from behind, usually not more than once, far from the choice point so that a response could be recorded on every trial.

Motor activity was measured in the open-field chamber that was made of white acrylic with a 50 cm × 50 cm surface area. Each mouse was placed in a corner of the open-field. The activity in central (25 cm × 25 cm surface area) and peripheral zones was recorded in a 5-min session for 3 consecutive days and analyzed by video tracking software (SD Instruments, San Diego, CA). The distance traveled and the time spent resting (<2 cm/s), moving slow (2–5 cm/s), or moving fast (>5 cm/s) in each zone were measured, as well as the time spent in the periphery and center of the apparatus.

The elevated plus-maze apparatus consisted of 4 arms in a cross-shaped form 70 cm in length with a 10 cm × 10 cm central region. Two of the arms were enclosed on 3 sides by walls (10 cm in height) facing each other, while the other two were open, except for a minimal border (0.5 cm in height) used to minimize falls. A mouse was placed in the central region and then the number of entries and the time spent inside enclosed and open arms were measured in a 5-min session on 2 consecutive days with the same video tracking system. The open/total arm entries and duration ratios were also calculated.

2.2.3. Spatial learning and memory—The Morris water maze consisted of a pool of blue opaque plastic, 116 cm in diameter with 75-cm high walls, filled with water (20°C) at a height of 31 cm. Powdered milk was evenly spread over the water surface to camouflage the escape platform (8 × 8 cm) made of white plastic and covered with a wire mesh grid to ensure a firm grip. The watered milk was removed every day after a few hours of training and the pool rinsed with clean water. The pool was contained in a room with external visual cues such as light fixtures and a ladder. The mice were placed next to and facing the wall successively in north (N), east (E), south (S), and west (W) positions, with the escape platform hidden 1 cm below water level in the middle of the NW quadrant. The same video tracking equipment was used to estimate path length and escape latencies in 4-trial sessions for 5 days with a 15-min inter-trial interval. The mice remained on the platform for at least 5 s. When the mice failed to reach the escape platform within the 1 min cut-off period, they were retrieved from the pool and placed on the platform for 5 s. After swimming, the mice were kept dry in a plastic holding cage filled with paper towels. In the morning after the acquisition phase, a probe trial was conducted by removing the platform and placing the

mouse next to and facing the N side. The time spent in the previously correct quadrant was measured for a single 1-min trial. In the afternoon, the visible platform subtask was conducted, with the escape platform lifted 1 cm above water level and shifted to the SE quadrant. A 17-cm high pole was inserted on top of the escape platform as a viewing aid. The same procedure was adopted as with place learning except that the subtest was conducted on a single day.

2.3. Intraperitoneal glucose tolerance test (IPGTT) and insulin tolerance test (ITT)

Forty-two and forty-four weeks after rAAV injection, IPGTT and ITT were conducted, respectively, as previously described (Ize-Ludlow *et al.*, 2011). Animals were fasted for 16 h, weighed and intraperitoneally injected with D-glucose (1.5 g/kg) for IPGTT. Blood was obtained using the tail nick method before and at 30, 60, 90 and 120 min after the D-glucose injection for monitoring glucose levels using Glucometer (ReliOn® Prime, Arkray USA, Minneapolis MN). Plasma insulin levels were determined by an insulin ELISA kit (Crystal Chem Inc., Downers Grove, IL). For ITT, mice were fasted for 4 h, weighed, and intraperitoneally injected with human insulin (0.75 units/kg; Novolin® R, Novo Nordisk, Princeton, NJ). Blood sugar was monitored before and at 15, 30, 60, and 90 min.

2.4. Murine CSF isolation

CSF was isolated from the cisterna magna compartment using the method previously described (DeMattos *et al.*, 2002). Forty-four weeks after rAAV5-IL-17A injection, mice were deeply anesthetized by Nembutal® (pentobarbital, 60mg/kg) and fixed face down on a narrow platform. An incision was made from the top of the skull to the dorsal thorax. The musculature from the base of the skull to the first vertebrae was carefully removed to expose the meninges overlying the cisterna magna. The surrounding area was gently cleaned with PBS using cotton swabs to remove any residual blood or other interstitial fluid. The arachnoid membrane covering the cistern was punctured with a 29-gauge insulin syringe. A polypropylene narrow bore pipette was immediately placed in the hole to collect CSF. As the primary CSF exiting the compartment was collected, a second collection was performed after the cistern was refilled within 2 min. About 10 to 15 µl CSF was collected from each mouse. Immediately after CSF collection, mice were killed and other tissues were also collected for analysis.

2.5. Immunohistochemical analyses

Forty-four weeks after rAAV5-IL-17A intracranial injection, mice were deeply anesthetized with Nembutal® (pentobarbital) and cardinally perfused with cold PBS and their brains were quickly removed. The neocortices and hippocampi of the left hemispheres were separately dissected and stored at -80°C for further studies. The right hemispheres were fixed in 4% paraformaldehyde for 48 h, stored overnight in 30% sucrose in 0.1 M PBS and frozen in Tissue-Tek optimal cutting temperature compound. Coronal sections (35 µm) of the brains were immuno-stained with anti-IBA1 antibody (Wako, Osaka, Japan) for detection of activated microglia. Endogenous peroxidase was eliminated by 1% H₂O₂ in 10% methanol Tris-buffered saline (TBS) treatment. After washing with 0.1 M Tris buffer (pH 7.5) and 0.1M TBS (pH 7.4), sections were blocked with 5% normal serum in 0.1 M TBS with 0.5% triton-X-100 (TBS-T) and then incubated with primary antibodies in 2%

serum in TBS-T for 18–48 h at 4°C. For the negative controls, slides were processed without primary antibody. After rinsing, the sections were incubated with biotinylated secondary antibodies in 2% serum TBS-T for 2 h at room temperature. The avidin-biotin peroxidase method using 3, 3'-diaminobenzidine as a substrate (Vector, Burlingame, CA) was performed. Sections were counterstained with hematoxylin. For the detection of fibrillar A β plaque, tissue sections were stained in 1% thioflavin S (Sigma) and rinsed with 70% ethanol. After washing with H₂O, the sections were mounted in 75% glycerol in H₂O. Histomorphometry was performed for quantification of amyloid deposits and reactive/activated glial cells using an Olympus BX61 automated microscope, Olympus Fluoview system, and Image Pro Plus v4 image analysis software (Media Cybernetics, Silver Spring, MD, USA) capable of color segmentation and automation via programmable macros. Five coronal brain sections from each mouse were analyzed, each separated by an approximately 400- μ m interval, starting at 1.3 mm posterior to the bregma to caudal. Both neocortex and hippocampus were found in all the brain sections. For fibrillar A β plaques, stained areas were expressed as a percentage of total neocortex or hippocampus, respectively. Activation of microglial was evaluated as previously reported (Hovens *et al.*, 2014). Because the IBA1 staining in cell body is darker than that in total cell, all the pixels that are darker than the background are traced to determine the total cell size and pixels that are above an applied staining threshold and size filter are traced to determine the total cell body size, as well as the number of microglia. For cerebral amyloid angiopathy (CAA), thioflavin S-positive blood vessels in 3 to 4 right hemisphere sections for each animal were enumerated. Each animal section starting at 1.3 mm posterior to the bregma to caudal was separated by 400- μ m interval. The quantification data were expressed as a number of thioflavin S-positive blood vessels including cross/transverse, oblique and longitudinal planes per square millimeter of neocortex area. Data were expressed as mean $\pm$ SEM (Kou *et al.*, 2015).

2.6. Quantification of plasma and CSF IL-17A by ELISA

Approximately 100 μ l of blood/mouse was drawn from the tail vein at 3, 7, and 11 months after injection by using microhematocrit heparin tubes (Fisher Scientific, Pittsburgh, PA) and centrifuged for 20 min at 2,000 $\times$ g for plasma isolation. Levels of IL-17A in plasma and CSF were determined by ELISA (eBioscience, Inc. San Diego, CA).

2.7. Quantification of brain A β and CSF by ELISA, and ABCA1 by western blotting

The neocortex and hippocampus were removed from -80°C , lysed using the Bio-Plex cell lysis kit (Bio-Rad Laboratories, Hercules, CA) and homogenized according to the manufacturer's protocol, and centrifuged at 16,000 $\times$ g for 30 min at 4°C. The supernatants containing buffer soluble A β were collected and the protein concentrations in the supernatants were determined by Bio-Rad Protein Assay (Bio-Rad Laboratories, Hercules, CA). The pellets containing insoluble A β were further dounce homogenized in guanidine hydrochloride (final concentration, 5 M) and then rock-shaken for 3 to 4 h at room temperature. Levels of buffer soluble and insoluble A β in the neocortex and hippocampus, and A β in the CSF were quantified by A β 42 and A β 40 ELISA kits (Invitrogen, Carlsbad, CA) according to the manufacturer's protocol.

The protein lysates were used to determine ABCA1 levels by western blotting, also. After determining protein concentrations, the proteins were electrophoresed under reducing conditions in 10% SDS–PAGE gels and transferred to PVDF membranes. The membranes were incubated overnight at 4°C anti-ABCA1 antibody (1:1000 dilution) (Abcam, Cambridge, MA), or anti- β -actin antibody (Sigma, St. Louis, MO) and specific bands were visualized by an enhanced chemiluminescence system (Amersham, Arlington Heights, IL). The optical densities of the protein bands were determined by densitometric scanning using an HP Scanjet G3010 Photo Scanner and Image J V1.40 (NIH, MD).

2.8. Cell culture and western blot

The bEnd.3 cells (ATCC® CRL2299™) were grown in ATCC-formulated Dulbecco's modified eagle's medium containing 10% fetal bovine serum (Atlanta Biologicals, Flowery Branch, GA) in a humidified 37°C incubator with a 5% CO₂ atmosphere. Cells were plated on culture dishes at 4×10^4 /ml cells. Forty-eight hours after seeding, bEnd.3 cells were treated with 0, 1, 10, 100 ng/ml of IL-17A (Cell Signaling Technology, Danvers, MA) for 24 h to determine ABCA1 protein expression. Other sets of bEnd.3 cells were treated with 100 ng/ml of IL-17A for 0 (no IL-17A), 5, 15 and 30 min to detect the activation of p-ERK1/2. Different sets of bEnd.3 cells were pre-treated with U0126 (25 μ M) or DMSO (vehicle) for 1 h followed by treatment with or without IL-17A (100 ng/ml) for 30 min or 24 h to determine expression levels of ERK or ABCA1, respectively. The cell lysates were prepared by using radioimmune precipitation assay (RIPA) lysis buffer containing complete mini-protease-inhibitor and phosphatase inhibitor cocktail tablets (Roche Diagnostics Corporation, Indianapolis, IN) and centrifuged at 12,000 xg for 10 min at 4°C for collecting supernatants. After determining protein concentrations, the proteins were electrophoresed under reducing conditions in 10% SDS–PAGE gels and transferred to PVDF membranes. The membranes were incubated overnight at 4°C with anti-phospho-ERK1/2, anti-ERK1/2 antibody (1:1000 dilution) (Cell Signaling Technology, Danvers, MA), anti-ABCA1, or anti- β -actin antibody and specific bands were visualized by an enhanced chemiluminescence system (Amersham, Arlington Heights, IL). The optical densities of the protein bands were determined by densitometric scanning using an HP Scanjet G3010 Photo Scanner and Image J V1.40 (NIH, MD). The optical density of p-ERK1/2 band was divided by that of total ERK1/2 band on the same lane from the same membrane for normalization. β -actin was used to normalize expression levels of ABCA1.

2.9. Statistical analysis

Data were expressed as mean $\pm$ SEM. The data were analyzed using SPSS version 22 software (IBM, Armonk, NY). One-way ANOVA followed by a least-significant-difference test was used for statistical comparisons among multiple groups and t-test for statistical comparisons between two groups. P values less than 0.05 were considered statistically significant.

3. Results

3.1. Overexpression of IL-17A via intracranial delivery of rAAV5 improves glucose metabolism

The bioactivity of IL-17A produced by the AAV-IL-17A vector was confirmed using 3T3 cells described previously (Yang *et al.*, 2016). To explore the potential role of IL-17A in AD, we intracranially delivered rAAV5-IL-17A into 2-month-old TgAPP^{swe}/PS1dE9 mice. Three, seven, and eleven months after injection, the average plasma IL-17A levels in IL-17A-overexpressing mice were 14.88 ± 1.95 , 14.69 ± 1.28 , and 9.31 ± 0.95 ng/ml respectively and the average CSF IL-17A level in IL-17A-overexpressing mice was 270.73 ± 48.83 ng/ml 11 months after injection, while IL-17A was undetectable in control mice by ELISA (Figure 1A). The regional distribution of rAAV5 in brain was mostly restricted to hippocampal neuronal cells, determined by beta-gal immunostaining (supplementary Figure 2A and B). Similarly, the copy number of rAAV5 in the hippocampus is 2.5-fold and 12-fold higher than that in the neocortex and cerebellum by qPCR method, respectively (supplementary Figure 2C and methods, $P < 0.05$ for both). The distribution of rAAV5 in peripheral tissues, including heart, lung, liver, spleen, kidney and intestine were undetectable by the same methods (data not shown).

Blood glucose and insulin levels in IL-17A-overexpressing mice were significantly lower than those in control mice throughout IPGTT (Figure 1B and C). There was no significant difference in ITT between groups (data not shown).

3.2. Overexpression of IL-17A improves anxiety and spatial learning but not long-term memory

3.2.1. Exploration and anxiety—In the T-maze spontaneous alternation test, mice in each group alternated above chance ($P < 0.05$) (Figure 2A) and choice latencies showed no difference between groups (Figure 2B). In the open-field evaluated by 2×3 ANOVA with repeated measures on the day factor (Figure 3), IL-17A-overexpressing mice had a higher center/total time ratio than control mice ($F(1,15)=10.77$, $P < 0.01$). Center/total ratio declined over days in both groups ($F(2,30)=5.08$, $P < 0.001$). Resting time and moving slow time increased over days, whereas moving fast time decreased over days, a sign of habituation to experimental conditions (day effects in all 3 measures, $P < 0.05$). There was also a significant group $\times$ day interaction for moving slow time ($F(2,30)=3.8$, $P < 0.05$) which increased more slowly and moving fast time ($F(2,30)=3.69$, $P < 0.05$) which decreased more slowly in IL-17A-overexpressing mice than controls. In contrast, distance travelled declined equally in both groups ($F(2,30)=23.67$, $P < 0.001$). In the elevated plus-maze test, there were group $\times$ day interaction ($F(1,15)=14.04$, $P < 0.01$) and day ($F(1,15)=35.11$, $P < 0.001$) effects, as values of open arm duration were higher in IL-17A-overexpressing mice only on day 1 ($t(15)=2.57$, $P < 0.05$), not day 2 when values were lower (Table 1).

3.2.2. Spatial learning and memory—In the acquisition phase of the Morris water maze test, there were significant day effects and no day vs group interactions for distance swum ($F(4,60)=5.06$, $P < 0.01$) and escape latencies ($F(4,60)=2.58$, $P < 0.05$) caused by a reduction in both values over days in both groups (Figure 4A and B). Furthermore, there was

a significant group effect ($F(1,15)=8.27$, $P<0.05$), because IL-17A-overexpressing mice were quicker than control mice. In the probe trial, the percentages of time spent in the correct quadrant were 27.40 ± 3.09 s and 26.73 ± 3.58 s for IL-17A-overexpressing and control mice, respectively ($P>0.05$) (Figure 4C). There were no intergroup differences in the visible platform subtask, also (data not shown).

3.3. Overexpression of IL-17A in mouse brain does not increase microglia activation but reduces A β in CSF and CAA

Fibrillar A β deposits are closely associated with activated microglia. Microglia can be activated by various endogenous and exogenous factors through their receptors including toll-like receptors, scavenger receptors, and numerous cytokine and chemokine receptors (Kierdorf and Prinz, 2013). IL-17A overexpression may aggravate microglia activation in an AD mouse model as observed in mouse models of intracerebral hemorrhage (Yu *et al.*, 2016). Activation of microglial is characterized by retraction and thickening of processes and increased cell body size. Microglial cell body to cell size ratio by IBA1 immunostaining has been shown to be positively correlated with microglial activation (Hovens *et al.*, 2014). Therefore, brain sections were subjected to immunohistochemical analysis with anti-IBA1 antibody for detection of activated microglia (Figure 5A). There was no difference in microglial cell body to cell size ratio (Figure 5B) and the total number of microglial cells (Figure 5C) between groups.

To determine the effect of IL-17A overexpression on A β load in the brain, A β deposits were visualized by thioflavin S fluorescence and quantified by morphometry (Figure 6A and B). The areas positive for thioflavin S fluorescence in the neocortex and hippocampus showed no statistic differences between groups: $0.36 \pm 0.03\%$ and $0.27 \pm 0.03\%$ for IL-17A-overexpressing mice, $0.37 \pm 0.03\%$ and $0.21 \pm 0.22\%$ in control mice, respectively. However, both CSF A β 40 and A β 42 were dramatically decreased in IL-17A-overexpressing mice (1.88 ± 0.37 and 0.50 ± 0.09 ng/ml, respectively) compared with that in control mice (7.35 ± 1.58 and 2.05 ± 0.35 ng/ml, respectively) ($P<0.01$ and $P<0.001$, respectively) (Figure 6C). Hippocampal soluble A β 40 levels in IL-17A-overexpressing mice (0.20 ± 0.05 ng/mg protein) were also significantly decreased compared with that in control mice (0.81 ± 0.39 ng/mg protein) ($P<0.05$) while no differences were found in hippocampal soluble A β 42 levels and in neocortical soluble A β 40 and A β 42 levels between groups (Figure 6D and E). No differences were found in insoluble A β 40 and A β 42 levels in the neocortex and hippocampus between groups (Figure 6F and G) ($P>0.05$).

Vascular A β deposition (CAA) is associated with vascular dysfunction and thought to play an important role in the AD pathogenesis (de la Torre, 2004; Smith and Greenberg, 2009). We enumerated the number of blood vessels positive for thioflavin S fluorescence in the right neocortices. The number of thioflavin S stained blood vessels in IL-17A-overexpressing mice ($0.05 \pm 0.02/\text{mm}^2$) considerably decreased compared with that in control mice (0.24 ± 0.06) (Figure 6H and I) ($P<0.01$).

3.4. IL-17A increases ABCA1 expression by ERK1/2 activation

In order to investigate possible mechanisms by which IL-17A decreases A β in CSF and CAA, protein levels of ABCA1 in IL-17A-overexpressing mice were determined by immunoblot analysis. Hippocampal ABCA1 protein increased in IL-17A-overexpressing mice compared with control mice ($P<0.05$) (Figure 7A and B), while there were no changes in cortex (data not shown). To further explore signal pathways of the increase in ABCA1 by IL-17A a mouse brain endothelial cell line, bEnd.3 cell, was used to determine the level of ERK1/2. Immunoblot analysis consistently revealed a dose-dependent increase in ABCA1 expression by IL-17A treatment (control vs 100 ng/ml IL-17A, $P<0.05$) (Figure 8A). Moreover, phospho-ERK1/2 (p-ERK1/2) levels were significantly increased in bEnd.3 cells ($P<0.05$) by 100 ng/ml IL-17A (Figure 8B). U0126 (25 μ M), an ERK1/2 kinase inhibitor, completely inhibited both the activation of ERK1/2 and the upregulation of ABCA1 expression by IL-17A (Figure 8C and D) ($P<0.05$).

4. Discussion

IL-17A is produced by Th17 cells and other immune cells, such as neutrophils, macrophages, dendritic cells, and natural killer, lymphoid tissue inducer and $\gamma\delta$ -T cells (Korn *et al.*, 2009; Zenaro *et al.*, 2015). Th17 cells infiltrate into the brain parenchyma in AD patients and mouse models, suggesting that Th17 cells may contribute to neuroinflammation by release of proinflammatory cytokines IL-17A and IL-22 (McManus *et al.*, 2014; Zhang *et al.*, 2013). IL-17A is released by neutrophils surrounding A β deposits in two transgenic mouse models of AD (5XFAD and 3xTg-AD mice), which suggests that IL-17A may be involved in AD pathogenesis and cognitive impairment (Zenaro *et al.*, 2015). However, the role of the IL-17A in the AD brain has not been well studied. To the best of our knowledge, this is the first study to demonstrate the effects of IL-17A overexpression on AD-like pathophysiology in an AD mouse model. The distribution of rAAV5 was discernible in the brain but not in the peripheral tissues (heart, lung, liver, spleen, kidney and intestine). However, IL-17A levels in blood was readily detectable in IL-17A overexpressing mice although the plasma IL-17A levels in IL-17A overexpressing mice were one twenty-ninth of the CSF IL-17A levels. There exist few possibilities. Despite of undetectable levels of rAAV5 in the peripheral tissues, we still cannot exclude the possibility that rAAV5 was released into brain vessels during intraventricular injection. Alternatively, because the blood-brain barrier plays an important role in regulating the rate of transport in (K_{in}) and out of (K_{out}) the brain (Wong *et al.*, 2013), the difference in K_{in} and/or K_{out} may be the main cause to distribution of IL-17A in CSF and plasma in IL-17A overexpressing mice.

Insulin signaling impairment/insulin resistance (IR) is a common and important pathophysiologic alteration between type 2 diabetes mellitus (T2DM) and AD (Craft and Watson, 2004; Sebastiao *et al.*, 2014; Verdile *et al.*, 2015), although the cellular and molecular mechanism that associates T2DM with AD is unknown (Rios *et al.*, 2014). IR may be one of the causes of AD. Hyperinsulinemia/insulin resistance may lead to alterations of protein expression or posttranslational modification central to AD pathogenesis, including tau, A β and the A β -degrading protease neprilysin (Morales-Corraliza *et al.*, 2016). A longitudinal study found that peripheral IR is associated with a higher risk of AD (Schrijvers

et al., 2010). IR in the CNS is involved in AD progression (Biessels and Reagan, 2015) and intranasal insulin delivery improved cognitive function in a subset of patients with AD or mild cognitive impairment (Benedict *et al.*, 2011; Claxton *et al.*, 2015; Craft *et al.*, 2012). Inversely, AD may be one of the causes of IR. Fasting glucose levels and/or insulin sensitivity are altered in 81% of AD patients suggesting that AD may lead to a diabetic phenotype (Janson *et al.*, 2004). A recent study demonstrated that intracerebroventricular infusion of A β oligomers induces peripheral insulin resistance possibly by triggering neuroinflammation (Clarke *et al.*, 2015). Therefore, AD appears to be reciprocally associated with IR, suggesting that improving IR may alleviate AD and vice versa. In this study, IL-17A significantly improved glucose metabolism in IL-17A-overexpressing mice compared with that in control mice. Although the IL-17A level in CSF was about 29 times higher than that in plasma, it is unclear whether the effect of IL-17A on glucose metabolism is due to activation of IL-17A signal transduction pathways in CNS, peripheral tissues or both.

A β accumulation occurs in the brain parenchyma and CAA and plays a causal role in AD pathogenesis. CAA is the deposition of A β in the walls of cerebral blood vessels and identified at autopsy in 75–98% of AD patients (Jellinger and Attems, 2003). A β deposition in CAA affects cerebral blood flow, alters BBB permeability, interferes with A β clearance or efflux from the brain and triggers deleterious inflammatory responses, contributing to development of vascular fragility, hemorrhages and, ultimately, dementia (Cupino and Zabel, 2014; Ghiso *et al.*, 2014; Kalaria, 1992). According to the two-hit vascular hypothesis of AD, brain vascular damages (hit 1) initiate A β accumulation (hit 2) in the brain, leading to neurodegeneration and memory decline in AD (Sagare *et al.*, 2012). Clinical imaging, epidemiological and pharmacotherapy studies support the notion that vascular changes play a central role in early AD pathogenesis (de la Torre, 2004; Iturria-Medina *et al.*, 2016). If not all, a subset of AD patients have disruptions of the BBB (Erickson and Banks, 2013), resulting in the reduction of cerebral blood flow (Spetsieris *et al.*, 2015) and A β clearance in late-onset AD (Mawuenyega *et al.*, 2010). We found that IL-17A-overexpressing mice show an approximately 4-fold decrease in A β deposition in CAA compared with control mice. Soluble A β levels in the hippocampus and CSF are similarly decreased in IL-17A-overexpressing mice. Contrary to this observation, a decrease in CSF A β , especially A β 42, is accompanied by an increase in brain A β deposition in the early stages of AD patients (Kuhlmann *et al.*, 2016; Maia *et al.*, 2013). Therefore, CSF A β levels do not always reflect A β load in the brain. Altogether, IL-17A may protect the brain vasculature from A β accumulation and prevent CAA. It is worth noting that IL-17A does not alter microglia activation in our experiments. Microglial cells are known to be important for amyloid plaque formation and clearance. However, eliminating microglia in AD mice failed to change amyloid plaque formation and modulate A β pathology (Grathwohl *et al.*, 2009; Spangenberg *et al.*, 2016). Thus, microglia may not be involved in reduction of A β deposits in IL-17A-overexpressing mice.

Cognitive impairment along with anxiety in AD animal models, which resembles that in AD patients, has been reported (Foidl *et al.*, 2016; Pietrzak *et al.*, 2015; Sil and Ghosh, 2016; Starkstein *et al.*, 2007). Protofibrillar A β 1–42 injection induced anxiety and AD-like pathology in rats (Sharma *et al.*, 2016), suggesting the deleterious effect of elevated A β

levels on behavior in AD. In our current study, IL-17A-overexpressing mice manifested behavioral alterations relative to controls. IL-17A-overexpressing mice were characterized by slower habituation in the open-field test, as indicated by less pronounced increases in moving slowly and less pronounced decreases in moving quickly across days of testing. Moreover, IL-17A-overexpressing mice had a higher center/total time ratio, indicating less anxiety in exploring the open-field. The lesser anxiety displayed by this group was also observed in the elevated plus-maze test, where open arm duration was increased relative to controls on day 1 though not on day 2 of testing when exploration is usually minimal. Therefore, IL-17A overexpression may improve anxiety by reducing soluble A β levels in the hippocampus. During acquisition of the Morris water maze task, IL-17A-overexpressing mice reached the escape platform faster than control mice. In contrast, there was no change in probe or visible platform subtasks, indicating cognitive improvement in the acquisition process but not long-term memory, visuomotor control, or swimming speed. Our results are consistent with the previous studies reporting the lack of association between A β deposits in brain and behavioral changes (Heikkinen et al. 2004; Holcomb et al. 1999; Howlett et al. 2004; Montarolo et al. 2013; Morgan 2003). It is possible that spatial memory deficits in AD model mice correlate with the selective modulation of the levels of hippocampal A β 42 (Faucher et al. 2015; Puolivali et al. 2002), which are unchanged in IL-17A-overexpressing mice.

ABCA1 is one of the routes of A β clearance from the brain across the BBB. The depletion and overexpression of ABCA1 increases and decreases brain A β in AD mouse models, respectively (Wahrle *et al.*, 2005; Wahrle *et al.*, 2008). In human, a loss-of-function mutation in ABCA1 with occurrence of 0.2% is associated with a high risk of AD and cerebrovascular disease (Nordestgaard *et al.*, 2015). Rarer non-synonymous variants of ABCA1 increase the risk for late onset AD, suggesting its protective effect against AD (Lupton *et al.*, 2014). Interestingly, the level of hippocampal ABCA1 in IL-17A-overexpressing mice is increased compared with that in control mice. Given that soluble A β levels in the hippocampus and CSF were similarly decreased in the IL-17A-overexpressing mice and neurons in the hippocampus highly expressed IL-17A by the rAAV5 gene delivery, we infer that IL-17A secreted from neurons induced ABCA1 expression in brain endothelial. Because IL-17A can induce activation of ERK1/2 signal (Guo *et al.*, 2014; Rodgers *et al.*, 2015; Song *et al.*, 2015) and because ERK1/2 signaling regulates ABCA1 expression in different types of cells (Chang *et al.*, 2013; Mulay *et al.*, 2013), we determined if IL-17A would increase ABCA1 expression via ERK1/2 activation in endothelial bEnd.3 cells. As expected, IL-17A increased phospho-ERK1/2 but not total ERK1/2. The upregulation of ABCA1 by IL-17A was blocked by U0126, an ERK inhibitor. The ERK1/2 signaling pathway also participates in the spatial memory (Gao *et al.*, 2016; Ghasemi *et al.*, 2014; Liu *et al.*, 2014). Thus, IL-17A overexpression may reduce CAA and soluble A β in the CSF and hippocampus by upregulating ABCA1 through the ERK1/2 signaling pathway.

5. Conclusions

Overexpression of IL-17A via intracranial delivery of rAAV5-IL-17A improved glucose metabolism, reduced A β levels in the CSF, hippocampus and CAA, and alleviated the anxiety in AD mice. Furthermore, our study suggests that IL-17A may decrease A β in the

brain by upregulating expression of ABCA1 through the ERK1/2 signaling pathway. However, IL-17A neither improved parenchymal A β deposits nor exacerbated neuroinflammation in an AD mouse model. Our findings suggest that IL-17A may play a protective role in the pathogenesis of AD.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Ms. Linda Walter (University of Illinois College of Medicine at Peoria) for assistance in preparation of this manuscript. This work was supported by grants from the National Institutes of Health (Grant Numbers AG030399, AG042082 and AG050854).

References

- Akhter H, Ballinger C, Liu N, van GT, Postlethwait EM, Liu RM. Cyclic ozone exposure induces gender-dependent neuropathology and memory decline in an animal model of Alzheimer's disease. *Toxicol Sci.* 2015; 147:222–234. [PubMed: 26116027]
- Benedict C, Frey WH, Schioth HB, Schultes B, Born J, Hallschmid M. Intranasal insulin as a therapeutic option in the treatment of cognitive impairments. *Exp Gerontol.* 2011; 46:112–115. [PubMed: 20849944]
- Biessels GJ, Reagan LP. Hippocampal insulin resistance and cognitive dysfunction. *Nat Rev Neurosci.* 2015; 16:660–671. [PubMed: 26462756]
- Browne TC, McQuillan K, McManus RM, O'Reilly JA, Mills KH, Lynch MA. IFN-gamma production by amyloid beta-specific Th1 cells promotes microglial activation and increases plaque burden in a mouse model of Alzheimer's disease. *J Immunol.* 2013; 190:2241–2251. [PubMed: 23365075]
- Burnham SC, Faux NG, Wilson W, Laws SM, Ames D, Bedo J, Bush AI, Doecke JD, Ellis KA, Head R, Jones G, Kiiveri H, Martins RN, Rembach A, Rowe CC, Salvado O, Macaulay SL, Masters CL, Villemagne VL. A blood-based predictor for neocortical A β burden in Alzheimer's disease: results from the AIBL study. *Mol Psychiatry.* 2014; 19:519–526. [PubMed: 23628985]
- Chang YC, Sheu WH, Chien YS, Tseng PC, Lee WJ, Chiang AN. Hyperglycemia accelerates ATP-binding cassette transporter A1 degradation via an ERK-dependent pathway in macrophages. *J Cell Biochem.* 2013; 114:1364–1373. [PubMed: 23239052]
- Clarke JR, Lyra E, Silva NM, Figueiredo CP, Frozza RL, Ledo JH, Beckman D, Katashima CK, Razolli D, Carvalho BM, Frazao R, Silveira MA, Ribeiro FC, Bomfim TR, Neves FS, Klein WL, Medeiros R, LaFerla FM, Carvalheira JB, Saad MJ, Munoz DP, Velloso LA, Ferreira ST, De Felice FG. Alzheimer-associated A β oligomers impact the central nervous system to induce peripheral metabolic deregulation. *EMBO Mol Med.* 2015; 7:190–210. [PubMed: 25617315]
- Claxton A, Baker LD, Hanson A, Trittschuh EH, Cholerton B, Morgan A, Callaghan M, Arbuckle M, Behl C, Craft S. Long-acting intranasal insulin detemir improves cognition for adults with mild cognitive impairment or early-stage Alzheimer's disease dementia. *J Alzheimers Dis.* 2015; 44:897–906. [PubMed: 25374101]
- Craft S, Baker LD, Montine TJ, Minoshima S, Watson GS, Claxton A, Arbuckle M, Callaghan M, Tsai E, Plymate SR, Green PS, Leverenz J, Cross D, Gerton B. Intranasal insulin therapy for Alzheimer disease and amnesic mild cognitive impairment: a pilot clinical trial. *Arch Neurol.* 2012; 69:29–38. [PubMed: 21911655]
- Craft S, Watson GS. Insulin and neurodegenerative disease: shared and specific mechanisms. *Lancet Neurol.* 2004; 3:169–178. [PubMed: 14980532]
- Cupino TL, Zabel MK. Alzheimer's silent partner: cerebral amyloid angiopathy. *Transl Stroke Res.* 2014; 5:330–337. [PubMed: 24323728]

- Dansokho C, Ait AD, Aid S, Toly-Ndour C, Chaigneau T, Calle V, Cagnard N, Holzenberger M, Piaggio E, Aucouturier P, Dorothee G. Regulatory T cells delay disease progression in Alzheimer-like pathology. *Brain*. 2016; 139:1237–1251. [PubMed: 26912648]
- de la Torre JC. Is Alzheimer's disease a neurodegenerative or a vascular disorder? Data, dogma, and dialectics. *Lancet Neurol*. 2004; 3:184–190. [PubMed: 14980533]
- DeMattos RB, Bales KR, Parsadanian M, O'Dell MA, Foss EM, Paul SM, Holtzman DM. Plaque-associated disruption of CSF and plasma amyloid-beta (A β) equilibrium in a mouse model of Alzheimer's disease. *J Neurochem*. 2002; 81:229–236. [PubMed: 12064470]
- Doecke JD, Laws SM, Faux NG, Wilson W, Burnham SC, Lam CP, Mondal A, Bedo J, Bush AI, Brown B, De RK, Ellis KA, Fowler C, Gupta VB, Head R, Macaulay SL, Pertile K, Rowe CC, Rembach A, Rodrigues M, Rumble R, Szoek C, Taddei K, Taddei T, Trounson B, Ames D, Masters CL, Martins RN. Blood-based protein biomarkers for diagnosis of Alzheimer disease. *Arch Neurol*. 2012; 69:1318–1325. [PubMed: 22801742]
- Engelhardt B, Ransohoff RM. The ins and outs of T-lymphocyte trafficking to the CNS: anatomical sites and molecular mechanisms. *Trends Immunol*. 2005; 26:485–495. [PubMed: 16039904]
- Erickson MA, Banks WA. Blood-brain barrier dysfunction as a cause and consequence of Alzheimer's disease. *J Cereb Blood Flow Metab*. 2013; 33:1500–1513. [PubMed: 23921899]
- Ferretti MT, Merlini M, Spani C, Gericke C, Schweizer N, Enzmann G, Engelhardt B, Kulic L, Suter T, Nitsch RM. T-cell brain infiltration and immature antigen-presenting cells in transgenic models of Alzheimer's disease-like cerebral amyloidosis. *Brain Behav Immun*. 2016; 54:211–225. [PubMed: 26872418]
- Fisher Y, Strominger I, Biton S, Nemirovsky A, Baron R, Monsonego A. Th1 polarization of T cells injected into the cerebrospinal fluid induces brain immunosurveillance. *J Immunol*. 2014; 192:92–102. [PubMed: 24307730]
- Foidl BM, Do-Dinh P, Hutter-Schmid B, Bliem HR, Humpel C. Cholinergic neurodegeneration in an Alzheimer mouse model overexpressing amyloid-precursor protein with the Swedish-Dutch-Iowa mutations. *Neurobiol Learn Mem*. 2016; 136:86–96. [PubMed: 27670619]
- Fratiglioni L, Launer LJ, Andersen K, Breteler MM, Copeland JR, Dartigues JF, Lobo A, Martinez-Lage J, Soininen H, Hofman A. Incidence of dementia and major subtypes in Europe: A collaborative study of population-based cohorts. Neurologic Diseases in the Elderly Research Group. *Neurology*. 2000; 54:S10–S15. [PubMed: 10854355]
- Gallagher JJ, Minogue AM, Lynch MA. Impaired performance of female APP/PS1 mice in the Morris water maze is coupled with increased A β accumulation and microglial activation. *Neurodegener Dis*. 2013; 11:33–41. [PubMed: 22627185]
- Gao L, Tian M, Zhao HY, Xu QQ, Huang YM, Si QC, Tian Q, Wu QM, Hu XM, Sun LB, McClintock SM, Zeng Y. TrkB activation by 7, 8-dihydroxyflavone increases synapse AMPA subunits and ameliorates spatial memory deficits in a mouse model of Alzheimer's disease. *J Neurochem*. 2016; 136:620–636. [PubMed: 26577931]
- Ghasemi R, Zarifkar A, Rastegar K, maghsoudi N, Moosavi M. Insulin protects against A β -induced spatial memory impairment, hippocampal apoptosis and MAPKs signaling disruption. *Neuropharmacology*. 2014; 85:113–120. [PubMed: 24881967]
- Ghisso J, Fossati S, Rostagno A. Amyloidosis associated with cerebral amyloid angiopathy: cell signaling pathways elicited in cerebral endothelial cells. *J Alzheimers Dis*. 2014; 42(Suppl 3):S167–S176. [PubMed: 24670400]
- Ghosh S, Geahlen RL. Stress granules modulate SYK to cause microglial cell dysfunction in Alzheimer's disease. *EBioMedicine*. 2015; 2:1785–1798. [PubMed: 26870803]
- Grathwohl SA, Kalin RE, Bolmont T, Prokop S, Winkelmann G, Kaeser SA, Odenthal J, Radde R, Eldh T, Gandy S, Aguzzi A, Staufenbiel M, Mathews PM, Wolburg H, Heppner FL, Jucker M. Formation and maintenance of Alzheimer's disease beta-amyloid plaques in the absence of microglia. *Nat Neurosci*. 2009; 12:1361–1363. [PubMed: 19838177]
- Guo X, Jiang X, Xiao Y, Zhou T, Guo Y, Wang R, Zhao Z, Xiao H, Hou C, Ma L, Lin Y, Lang X, Feng J, Chen G, Shen B, Han G, Li Y. IL-17A signaling in colonic epithelial cells inhibits pro-inflammatory cytokine production by enhancing the activity of ERK and PI3K. *PLoS ONE*. 2014; 9:e89714. [PubMed: 24586980]

- Hickey WF. Basic principles of immunological surveillance of the normal central nervous system. *Glia*. 2001; 36:118–124. [PubMed: 11596120]
- Hickman SE, Allison EK, El Khoury J. Microglial dysfunction and defective beta-amyloid clearance pathways in aging Alzheimer's disease mice. *J Neurosci*. 2008; 28:8354–8360. [PubMed: 18701698]
- Hovens IB, Nyakas C, Schoemaker RG. A novel method for evaluating microglial activation using ionized calcium-binding adaptor protein-1 staining: cell body to cell size ratio. *Neuroimmunol Neuroinflammation*. 2014; 1:82–88.
- Iturria-Medina Y, Sotero RC, Toussaint PJ, Mateos-Perez JM, Evans AC. Early role of vascular dysregulation on late-onset Alzheimer's disease based on multifactorial data-driven analysis. *Nat Commun*. 2016; 7:11934. [PubMed: 27327500]
- Ize-Ludlow D, Lightfoot YL, Parker M, Xue S, Wasserfall C, Haller MJ, Schatz D, Becker DJ, Atkinson MA, Mathews CE. Progressive erosion of beta-cell function precedes the onset of hyperglycemia in the NOD mouse model of type 1 diabetes. *Diabetes*. 2011; 60:2086–2091. [PubMed: 21659497]
- Janson J, Laedtke T, Parisi JE, O'Brien P, Petersen RC, Butler PC. Increased risk of type 2 diabetes in Alzheimer disease. *Diabetes*. 2004; 53:474–481. [PubMed: 14747300]
- Jellinger KA, Attems J. Incidence of cerebrovascular lesions in Alzheimer's disease: a postmortem study. *Acta Neuropathol*. 2003; 105:14–17. [PubMed: 12471455]
- Kalaria RN. The blood-brain barrier and cerebral microcirculation in Alzheimer disease. *Cerebrovasc Brain Metab Rev*. 1992; 4:226–260. [PubMed: 1389957]
- Kawas C, Gray S, Brookmeyer R, Fozard J, Zonderman A. Age-specific incidence rates of Alzheimer's disease: the Baltimore Longitudinal Study of Aging. *Neurology*. 2000; 54:2072–2077. [PubMed: 10851365]
- Kierdorf K, Prinz M. Factors regulating microglia activation. *Front Cell Neurosci*. 2013; 7:44. [PubMed: 23630462]
- Kipnis J, Cohen H, Cardon M, Ziv Y, Schwartz M. T cell deficiency leads to cognitive dysfunction: implications for therapeutic vaccination for schizophrenia and other psychiatric conditions. *Proc Natl Acad Sci USA*. 2004; 101:8180–8185. [PubMed: 15141078]
- Korn T, Bettelli E, Oukka M, Kuchroo VK. IL-17 and Th17 Cells. *Annu Rev Immunol*. 2009; 27:485–517. [PubMed: 19132915]
- Kou J, Yang J, Lim JE, Pattanayak A, Song M, Planque S, Paul S, Fukuchi K. Catalytic immunoglobulin gene delivery in a mouse model of Alzheimer's disease: prophylactic and therapeutic applications. *Mol Neurobiol*. 2015; 51:43–56. [PubMed: 24733587]
- Kuhlmann J, Andreasson U, Pannee J, Bjerke M, Portelius E, Leinenbach A, Bittner T, Korecka M, Jenkins RG, Vanderstichele H, Stoops E, Lewczuk P, Shaw LM, Zegers I, Schimmel H, Zetterberg H, Blennow K. CSF Aβ₁₋₄₂ - an excellent but complicated Alzheimer's biomarker - a route to standardisation. *Clin Chim Acta*. 2016
- Kukull WA, Higdon R, Bowen JD, McCormick WC, Teri L, Schellenberg GD, van BG, Jolley L, Larson EB. Dementia and Alzheimer disease incidence: a prospective cohort study. *Arch Neurol*. 2002; 59:1737–1746. [PubMed: 12433261]
- Lim J, Kou J, Song M, Pattanayak A, Jin J, Lalonde R, Fukuchi K. MyD88 deficiency ameliorates beta-amyloidosis in an animal model of Alzheimer's disease. *Am J Pathol*. 2011; 179:1095–1103. [PubMed: 21763676]
- Liu MY, Wang S, Yao WF, Zhang ZJ, Zhong X, Sha L, He M, Zheng ZH, Wei MJ. Memantine improves spatial learning and memory impairments by regulating NGF signaling in APP/PS1 transgenic mice. *Neuroscience*. 2014; 273:141–151. [PubMed: 24846616]
- Lombardi VR, Garcia M, Rey L, Cacabelos R. Characterization of cytokine production, screening of lymphocyte subset patterns and in vitro apoptosis in healthy and Alzheimer's Disease (AD) individuals. *J Neuroimmunol*. 1999; 97:163–171. [PubMed: 10408971]
- Lueg G, Gross CC, Lohmann H, Johnen A, Kemmling A, Deppe M, Groger J, Minnerup J, Wiendl H, Meuth SG, Duning T. Clinical relevance of specific T-cell activation in the blood and cerebrospinal fluid of patients with mild Alzheimer's disease. *Neurobiol Aging*. 2015; 36:81–89. [PubMed: 25277040]

- Lupton MK, Proitsi P, Lin K, Hamilton G, Daniilidou M, Tsolaki M, Powell JF. The role of ABCA1 gene sequence variants on risk of Alzheimer's disease. *J Alzheimers Dis.* 2014; 38:897–906. [PubMed: 24081377]
- Maia LF, Kaeser SA, Reichwald J, Hruscha M, Martus P, Staufenbiel M, Jucker M. Changes in amyloid-beta and Tau in the cerebrospinal fluid of transgenic mice overexpressing amyloid precursor protein. *Sci Transl Med.* 2013; 5:194re2.
- Mawuenyega KG, Sigurdson W, Ovod V, Munsell L, Kasten T, Morris JC, Yarasheski KE, Bateman RJ. Decreased clearance of CNS beta-amyloid in Alzheimer's disease. *Science.* 2010; 330:1774. [PubMed: 21148344]
- McManus RM, Higgins SC, Mills KH, Lynch MA. Respiratory infection promotes T cell infiltration and amyloid-beta deposition in APP/PS1 mice. *Neurobiol Aging.* 2014; 35:109–121. [PubMed: 23993702]
- Monsonogo A, Imitola J, Petrovic S, Zota V, Nemirovsky A, Baron R, Fisher Y, Owens T, Weiner HL. Abeta-induced meningoencephalitis is IFN-gamma-dependent and is associated with T cell-dependent clearance of Abeta in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A.* 2006; 103:5048–5053. [PubMed: 16549802]
- Monsonogo A, Zota V, Karni A, Krieger JJ, Bar-Or A, Bitan G, Budson AE, Sperling R, Selkoe DJ, Weiner HL. Increased T cell reactivity to amyloid beta protein in older humans and patients with Alzheimer disease. *J Clin Invest.* 2003; 112:415–422. [PubMed: 12897209]
- Morales-Corraliza J, Wong H, Mazzella MJ, Che S, Lee SH, Petkova E, Wagner JD, Hemby SE, Ginsberg SD, Mathews PM. Brain-wide insulin resistance, tau phosphorylation changes, and hippocampal neprilysin and amyloid-beta alterations in a monkey model of Type 1 diabetes. *J Neurosci.* 2016; 36:4248–4258. [PubMed: 27076423]
- Mulay V, Wood P, Manetsch M, Darabi M, Cairns R, Hoque M, Chan KC, Reverter M, Alvarez-Guaita A, Rye KA, Rentero C, Heeren J, Enrich C, Grewal T. Inhibition of mitogen-activated protein kinase Erk1/2 promotes protein degradation of ATP binding cassette transporters A1 and G1 in CHO and HuH7 cells. *PLoS ONE.* 2013; 8:e62667. [PubMed: 23634230]
- Nordestgaard LT, Tybjaerg-Hansen A, Nordestgaard BG, Frikke-Schmidt R. Loss-of-function mutation in ABCA1 and risk of Alzheimer's disease and cerebrovascular disease. *Alzheimers Dement.* 2015; 11:1430–1438. [PubMed: 26079414]
- Pietrzak RH, Lim YY, Neumeister A, Ames D, Ellis KA, Harrington K, Lautenschlager NT, Restrepo C, Martins RN, Masters CL, Villemagne VL, Rowe CC, Maruff P. Amyloid-beta, anxiety, and cognitive decline in preclinical Alzheimer disease: a multicenter, prospective cohort study. *JAMA Psychiatry.* 2015; 72:284–291. [PubMed: 25629787]
- Ridolfi E, Barone C, Scarpini E, Galimberti D. The role of the innate immune system in Alzheimer's disease and frontotemporal lobar degeneration: an eye on microglia. *Clin Dev Immunol.* 2013; 2013:939786. [PubMed: 23970926]
- Rios JA, Cisternas P, Arrese M, Barja S, Inestrosa NC. Is Alzheimer's disease related to metabolic syndrome? A Wnt signaling conundrum. *Prog Neurobiol.* 2014; 121:125–146. [PubMed: 25084549]
- Rodgers JM, Robinson AP, Rosler ES, Lariosa-Willingham K, Persons RE, Dugas JC, Miller SD. IL-17A activates ERK1/2 and enhances differentiation of oligodendrocyte progenitor cells. *Glia.* 2015; 63:768–779. [PubMed: 25557204]
- Rodriguez JJ, Butt AM, Gardenal E, Parpura V, Verkhratsky A. Complex and differential glial responses in Alzheimer's disease and ageing. *Curr Alzheimer Res.* 2016; 13:343–358. [PubMed: 26923267]
- Schrijvers EM, Witteman JC, Sijbrands EJ, Hofman A, Koudstaal PJ, Breteler MM. Insulin metabolism and the risk of Alzheimer disease: the Rotterdam Study. *Neurology.* 2010; 75:1982–1987. [PubMed: 21115952]
- Sebastiao I, Candeias E, Santos MS, de Oliveira CR, Moreira PI, Duarte AI. Insulin as a bridge between type 2 diabetes and Alzheimer disease - How anti-diabetics could be a solution for dementia. *Front Endocrinol (Lausanne).* 2014; 5:110. [PubMed: 25071725]

- Shalit F, Sredni B, Brodie C, Kott E, Huberman M. T lymphocyte subpopulations and activation markers correlate with severity of Alzheimer's disease. *Clin Immunol Immunopathol.* 1995; 75:246–250. [PubMed: 7768042]
- Sharma S, Verma S, Kapoor M, Saini A, Nehru B. Alzheimer's disease like pathology induced six weeks after aggregated amyloid-beta injection in rats: increased oxidative stress and impaired long-term memory with anxiety-like behavior. *Neurol Res.* 2016; 38:838–850. [PubMed: 27431920]
- Sierksma AS, Prickaerts J, Chouliaras L, Rostamian S, Delbroek L, Rutten BP, Steinbusch HW, Van den Hove DL. Behavioral and neurobiological effects of prenatal stress exposure in male and female APPswe/PS1dE9 mice. *Neurobiol Aging.* 2013; 34:319–337. [PubMed: 22738723]
- Sil S, Ghosh T. Cox-2 plays a vital role in the impaired anxiety like behavior in colchicine induced rat model of Alzheimer disease. *Behav Neurol.* 2016; 2016:1501527. [PubMed: 26880859]
- Smith EE, Greenberg SM. Beta-amyloid, blood vessels, and brain function. *stroke.* 2009; 40:2601–2606. [PubMed: 19443808]
- Smolders J, Remmerswaal EB, Schuurman KG, Melief J, van Eden CG, van Lier RA, Huitinga I, Hamann J. Characteristics of differentiated CD8(+) and CD4 (+) T cells present in the human brain. *Acta Neuropathol.* 2013; 126:525–535. [PubMed: 23880787]
- Song X, Dai D, He X, Zhu S, Yao Y, Gao H, Wang J, Qu F, Qiu J, Wang H, Li X, Shen N, Qian Y. Growth factor FGF2 cooperates with Interleukin-17 to repair intestinal epithelial damage. *Immunity.* 2015; 43:488–501. [PubMed: 26320657]
- Spangenberg EE, Lee RJ, Najafi AR, Rice RA, Elmore MR, Blurton-Jones M, West BL, Green KN. Eliminating microglia in Alzheimer's mice prevents neuronal loss without modulating amyloid-beta pathology. *Brain.* 2016; 139:1265–1281. [PubMed: 26921617]
- Spetsieris PG, Ko JH, Tang CC, Nazem A, Sako W, Peng S, Ma Y, Dhawan V, Eidelberg D. Metabolic resting-state brain networks in health and disease. *Proc Natl Acad Sci USA.* 2015; 112:2563–2568. [PubMed: 25675473]
- Starkstein SE, Jorge R, Petracca G, Robinson RG. The construct of generalized anxiety disorder in Alzheimer disease. *Am J Geriatr Psychiatry.* 2007; 15:42–49. [PubMed: 17194814]
- Taniuchi N, Niidome T, Goto Y, Akaike A, Kihara T, Sugimoto H. Decreased proliferation of hippocampal progenitor cells in APPswe/PS1dE9 transgenic mice. *NeuroReport.* 2007; 18:1801–1805. [PubMed: 18090315]
- Togo T, Akiyama H, Iseki E, Kondo H, Ikeda K, Kato M, Oda T, Tsuchiya K, Kosaka K. Occurrence of T cells in the brain of Alzheimer's disease and other neurological diseases. *J Neuroimmunol.* 2002; 124:83–92. [PubMed: 11958825]
- Verdile G, Keane KN, Cruzat VF, Medic S, Sabale M, Rowles J, Wijesekara N, Martins RN, Fraser PE, Newsholme P. Inflammation and oxidative stress: The molecular connectivity between insulin resistance, obesity, and Alzheimer's disease. *Mediators Inflamm.* 2015; 2015:105828. [PubMed: 26693205]
- Wahrle SE, Jiang H, Parsadanian M, Hartman RE, Bales KR, Paul SM, Holtzman DM. Deletion of Abca1 increases Abeta deposition in the PDAPP transgenic mouse model of Alzheimer disease. *J Biol Chem.* 2005; 280:43236–43242. [PubMed: 16207708]
- Wahrle SE, Jiang H, Parsadanian M, Kim J, Li A, Knoten A, Jain S, Hirsch-Reinshagen V, Wellington CL, Bales KR, Paul SM, Holtzman DM. Overexpression of ABCA1 reduces amyloid deposition in the PDAPP mouse model of Alzheimer disease. *J Clin Invest.* 2008; 118:671–682. [PubMed: 18202749]
- Wang J, Tanila H, Puolivali J, Kadish I, van Groen T. Gender differences in the amount and deposition of amyloidbeta in APPswe and PS1 double transgenic mice. *Neurobiol Dis.* 2003; 14:318–327. [PubMed: 14678749]
- Wong AD, Ye M, Levy AF, Rothstein JD, Bergles DE, Searson PC. The blood-brain barrier: an engineering perspective. *Front Neuroeng.* 2013; 6:7. [PubMed: 24009582]
- Yang J, Kou J, Lim JE, Lalonde R, Fukuchi KI. Intracranial delivery of interleukin-17A via adeno-associated virus fails to induce physical and learning disabilities and neuroinflammation in mice but improves glucose metabolism through AKT signaling pathway. *Brain Behav Immun.* 2016; 53:84–95. [PubMed: 26562537]

- Yu A, Duan H, Zhang T, Pan Y, Kou Z, Zhang X, Lu Y, Wang S, Yang Z. IL-17A promotes microglial activation and neuroinflammation in mouse models of intracerebral haemorrhage. *Mol Immunol*. 2016; 73:151–157. [PubMed: 27107665]
- Zenaro E, Pietronigro E, Della BV, Piacentino G, Marongiu L, Budui S, Turano E, Rossi B, Angiari S, Dusi S, Montresor A, Carlucci T, Nani S, Tosadori G, Calciano L, Catalucci D, Berton G, Bonetti B, Constantin G. Neutrophils promote Alzheimer's disease-like pathology and cognitive decline via LFA-1 integrin. *Nat Med*. 2015; 21:880–886. [PubMed: 26214837]
- Zhang J, Ke KF, Liu Z, Qiu YH, Peng YP. Th17 cell-mediated neuroinflammation is involved in neurodegeneration of abeta1-42-induced Alzheimer's disease model rats. *PLoS ONE*. 2013; 8:e75786. [PubMed: 24124514]
- Ziv Y, Ron N, Butovsky O, Landa G, Sudai E, Greenberg N, Cohen H, Kipnis J, Schwartz M. Immune cells contribute to the maintenance of neurogenesis and spatial learning abilities in adulthood. *Nat Neurosci*. 2006; 9:268–275. [PubMed: 16415867]

Highlights

- IL-17A overexpression decreases cerebral amyloid angiopathy in Alzheimer model mice
- IL-17A overexpression decreases CSF and hippocampal A β in Alzheimer model mice
- IL-17A overexpression reduces endothelial ABCA1 expression via ERK activation
- IL-17A overexpression does not exacerbate neuroinflammation in Alzheimer model mice

Figure 1. Overexpression of IL-17A in AD mouse brain and its effect on IPGTT

(A) The levels of IL-17A in plasma and CSF were determined by ELISA at the indicated time points. Control indicates plasma IL-17A levels in mice 11 months after PBS and AAV5-LacZ injection (undiscernible). (B and C) Forty-two weeks after injection, mice were fasted for 16 h and injected i.p. with 1.5 g glucose/kg body weight and blood glucose and insulin were measured before and after glucose injection at the indicated time points. (Control mice, $n = 9$; IL-17A-overexpressing mice, $n = 8$. * $P < 0.05$, # $P < 0.01$ and ## $P < 0.001$)

Figure 2. Effects of IL-17A overexpression on T-maze

The number of alternations (A) and the latencies (B) before responding with 1 min cut-off are shown as mean $\pm$ SEM. * $P < 0.05$ relative to chance levels at 50% (Control mice, $n = 9$; IL-17A-overexpressing mice, $n = 8$)

Figure 3. Effects of IL-17A overexpression on mice in open-field

The percentages of peripheral time (A), central time (B), resting time (C), moving slow time (D) and moving fast time (E) together with total travelling distances (F) in a 5-min session are shown as mean $\pm$ SEM for 3 days. (Control mice, $n = 9$; IL-17A-overexpressing mice, $n = 8$)

Figure 4. Effects of IL-17A overexpression on cognitive function in Morris water maze
 (A) Total escape latencies (s), (B) total path lengths (cm) per day, and (C) percentage of time in all quads in probe test are shown as mean $\pm$ SEM. (Control mice, n = 9; IL-17A-overexpressing mice, n = 8)

Figure 5. Effect of IL-17A overexpression on microglia activation by IBA1 staining

(A) Forty-four weeks after injection, mice were euthanized at 52 weeks of age and brain sections were subjected to immunohistochemistry using anti-IBA1 antibody. Scale bars 500 μ m. (B) The ratio of microglial cell body to cell size in hippocampus and (C) the total number of microglial cells in hippocampus were determined. (Control mice, $n = 9$; IL-17A-overexpressing mice, $n = 8$)

Figure 6. Effect of IL-17A overexpression on A β load and cerebral amyloid angiopathy in the brain

(A) Fibrillar A β deposits in the brain were visualized by thioflavin S staining. Scale bars 500 μ m. (B) Average percentages of areas showing thioflavin S positive staining measured by morphometry in the cortex and hippocampus are shown. (C) The levels of A β 40 and A β 42 in CSF were measured by ELISA. (D, E, F, and G) The levels of soluble and insoluble A β 40 and A β 42 in cortex and hippocampus were measured by ELISA. Data are shown as mean $\pm$ SEM. (H) Cerebral amyloid angiopathy (CAA) in the right neocortices in control mice and IL-17A-overexpressing mice was visualized by thioflavin S fluorescence and blood vessels positive for thioflavin S fluorescence were enumerated. (I) The numbers of blood vessels positive for thioflavin S fluorescence per square millimeter are shown (mean $\pm$ SEM). Scale bars 100 μ m. (Control mice, n = 9; IL-17A-overexpressing mice, n = 8. # P < 0.01)

Figure 7. Effect of IL-17A on ABCA1 expression in the hippocampus
(A) Hippocampal lysates were analyzed by western blotting using ABCA1 antibody and (B) bar graph represents the results of densitometric analysis. (Control mice, n=9; IL-17A-overexpressing mice, n = 8, ## P < 0.001)

Figure 8. Effects of IL-17A on ERK1/2 and ABCA1 expression in bEnd.3 cells

(A) bEnd.3 cells were treated with 0 (Control), 1, 10 and 100 ng/ml IL-17A for 24 h. The cell lysates were analyzed by western blotting using ABCA1 antibody and bar graph represents the results of densitometric analysis. (B) bEnd.3 cells were treated with 100 ng/ml IL-17A for 0 (Control), 5 min, 15 min and 30 min. The cell lysates were analyzed by western blotting using phospho-ERK1/2 and total ERK1/2 antibodies and bar graph represents the results of densitometric analysis. (C) bEnd.3 cells were pretreated with DMSO (vehicle) or phospho-ERK1/2 inhibitor, U0126 (25 μ M), for 1 h followed by treatment with or without IL-17A (100 ng/ml) for 30 min. The cell lysates were analyzed by western blotting using phospho-ERK1/2 and total ERK1/2 antibodies and bar graph represents the results of densitometric analysis. (D) bEnd.3 cells were pretreated with DMSO or U0126 (25 μ M) for 1h followed by treatment with or without IL-17A (100 ng/ml) for 24 h. The cell lysates were analyzed by western blotting using ABCA1 antibody and bar graph represents the results of densitometric analysis. The statistical results, * $P < 0.05$ and ## $P < 0.001$, are from three independent experiments as shown in A, B, C and D.

Table 1Effects of IL-17A on exploratory activity in plus-maze (mean $\pm$ SEM; * P<0.05)

Measures	Control mice	IL-17A-overexpressing mice
Day1		
Open arms		
Entries	6.89 $\pm$ 1.15	8.13 $\pm$ 1.57
Duration (s)*	30.47 $\pm$ 7.27	67.83 $\pm$ 11.92
Enclosed arms		
Entries	14.22 $\pm$ 1.96	15.00 $\pm$ 1.51
Duration (s)	228.54 $\pm$ 10.20	190.37 $\pm$ 12.8
Open/total ratio (%)		
Entries	0.15 $\pm$ 0.02	0.17 $\pm$ 0.02
Duration (s)	0.10 $\pm$ 0.02	0.23 $\pm$ 0.04
Day2		
Open arms		
Entries	1.89 $\pm$ 0.40	1.13 $\pm$ 0.37
Duration (s)	17.09 $\pm$ 4.23	8.45 $\pm$ 2.67
Enclosed arms		
Entries	10.11 $\pm$ 1.37	10.38 $\pm$ 1.77
Duration (s)	260.65 $\pm$ 6.20	268.59 $\pm$ 5.81
Open/total ratio (%)		
Entries	0.08 $\pm$ 0.01	0.04 $\pm$ 0.01
Duration (s)	0.06 $\pm$ 0.01	0.03 $\pm$ 0.01