Published in final edited form as:

Trends Microbiol. 1997 February ; 5(2): 52–58.

Cell invasion by the vertebrate stages of *Plasmodium*

Photini Sinnis* and

Dept of Medical and Molecular Parasitology, New York University Medical Center, New York, NY 10016, USA

B. Kim Lee Sim

EntreMed, 9610 Medical Center Drive, Rockville, MD 20850, USA

Abstract

Protozoans of the genus *Plasmodium* are the causative agents of malaria; they have a complex life cycle involving vertebrate and arthropod hosts and have three distinct invasive stages. Although the invasive stages probably invade cells using similar mechanisms, each stage has a different host cell specificity and utilizes different receptors to enter cells.

Of the many forms the malaria parasite takes during its complex life cycle, three stages must invade host cells; the ookinete, the sporozoite and the merozoite (see Fig. 1 for an overview of the life cycle of the malaria parasite). In this review, we will discuss sporozoite invasion of hepatocytes and erythrocyte invasion by merozoites (see Box 1 for a glossary of terms used). Unlike most viral and bacterial intracellular pathogens, which trigger their own uptake by the host cell, *Plasmodium* invasion of cells appears to be an active process in which the parasite utilizes an actin-based motility system to enter the target cell (Ref. 1 and reviewed in Ref. 2). Although the invasive stages of the parasite are morphologically and biochemically different from one another, they share a highly conserved structural organization and apical organelles called micronemes and rhoptries. The function of these conserved apical structures has not been fully elucidated but they are likely to be required for host cell invasion. Although these common structures may confer a similar overall pattern of target cell invasion, the details differ. Each invasive stage of the parasite has a different target cell specificity, probably governed by specific receptor–ligand interactions. Much of the work on the sporozoite stage of the parasite has centered around the challenge faced by the parasite in finding its target. In contrast, merozoites are usually released in close proximity to a potential host cell and most of the work in this field has focused on the process of invasion after initial attachment of the parasite to the erythrocyte.

Hepatocyte invasion by sporozoites

Injection of two to ten *Plasmodium* sporozoites can initiate infection^{3,4}. Although it is not known how many parasites are injected by a mosquito in the field, laboratory studies suggest that the median number of injected parasites during a blood meal is 15 (Ref. 5). In addition to being efficient, sporozoite invasion of hepatocytes is a rapid process, occurring minutes

* tel: +1 212 263 5346, fax +1 212 263 8179, sinnip01@mcrcr6.med.nyu.edu.

after intravenous injection⁶. The speed and efficiency of hepatocyte infection by sporozoites suggest that the parasites go to the liver directly.

The Kupffer cell hypothesis

The mechanism by which the parasites are arrested in the liver is not known. The endothelial cell lining of the liver sinusoids is unique in that it has open fenestrations, but the diameter of the fenestrae (0.1 μm)⁷ is smaller than that of a sporozoite (1 μm). To circumvent this difficulty, it was thought that the parasites initially interact with Kupffer cells (which are found in the sinusoids) before invading hepatocytes.

Morphological studies performed to address the role of Kupffer cells are difficult to interpret because parasites are found in hepatocytes, sinusoids and Kupffer cells minutes after injection^{6,8}. When Kupffer cells are eliminated using liposome-encapsulated dichloromethylene diphosphonate, there are 4–5-fold more exoerythrocytic forms (EEF) developing in the liver compared with controls, indicating that macrophages are not required for sporozoite invasion of hepatocytes⁹. Studies *in vitro* show that sporozoites can interact with, invade and sometimes kill macrophages^{1,10}. It is possible, therefore, that sporozoites may interact with Kupffer cells *in vivo*, although the nature of this interaction requires further investigation.

An alternative theory is that sporozoites bind to and pass through hepatic endothelial cells¹¹. Although there are no data to demonstrate the transendothelial passage of sporozoites, the electron microscopic techniques used to localize parasites in the liver may not be able to distinguish between the presence of parasites in liver sinusoids and the binding of parasites to hepatic endothelia. The use of new microscopic techniques, as well as increased knowledge of organ-specific endothelial cell markers, may help to answer this question.

Hepatocyte-adhesive domains on sporozoite surface proteins

Alternatively, sporozoites may invade hepatocytes directly, without initially interacting with Kupffer cells or endothelial cells. This hypothesis is supported by recent studies demonstrating the specific binding of two sporozoite surface proteins, the circumsporozoite protein (CS; reviewed in Ref. 12) and the sporozoite surface protein 2 (SSP2; Ref. 13) or thrombospondin-related adhesion protein (TRAP; Ref. 14), to the basolateral domain of hepatocytes^{15,16}. Immunoelectron microscopy of liver sections incubated with either CS or SSP2/TRAP shows that both proteins bind to hepatocyte microvilli within the space of Disse, the portion of the cell exposed to the circulation. Neither protein binds to Kupffer cells nor to endothelial cells. In addition, antibodies to either CS or SSP2/TRAP inhibit sporozoite invasion of HepG2 cells^{15,17,18}, a hepatoma cell line permissive for *Plasmodium berghei* sporozoite invasion and development *in vitro*.

CS and SSP2/TRAP have only one region of identity: a known cell-adhesive motif with a high degree of homology to the type I thrombospondin repeats¹⁴. In CS, this motif is called region II-plus^{19,20}; it is conserved in all CS and SSP2/TRAP proteins sequenced to date (Table 1) and it is required for binding of CS and SSP2/TRAP to hepatocytes^{15,17}.

Initial studies have shown that CS and SSP2FFRAP bind to sulfated glycoconjugates in a region II-plus- dependent fashion^{17,21}. Subsequent experiments have demonstrated that CS binds to heparan sulfate proteoglycans (HSPGs)²². Although the identity of the cell-surface sulfated glycoconjugate that binds to SSP2/TRAP is not known, it may also be an HSPG because SSP2FFRAP binding to liver sections is eliminated after treatment with heparitinase¹⁶.

Structural studies on the parasite ligand have shown that the downstream, positively charged residues, as well as the interspersed hydrophobic amino acids of region II-plus, are required for binding to HSPGs (Ref. 20). In addition, only multimers of CS, or of peptides representing region II-plus, bind in a stable fashion to the hepatocyte receptors. It is likely that the lysines and arginines of region II-plus form ionic bonds with the negatively charged sulfate molecules of the HSPG glycosaminoglycan chains (GAGs).

It is not clear why the parasite should have two surface proteins with the same cell-adhesive motif. Perhaps the fine specificity of the receptor for each protein differs because heparan sulfate consists of repeating disaccharides that undergo secondary modifications, giving rise to an enormous amount of structural diversity throughout the length of each GAG chain. CS and SSP2/TRAP, therefore, may have different binding specificities based on conservative changes in region II-plus and/or amino acid residues in neighboring regions. Alternatively, CS and SSP2/TRAP may function at different stages of hepatocyte invasion. A different functional role for each protein is suggested by their different patterns of expression on the surface of the parasite: CS forms a dense coat whereas SSP2FFRAP gives a punctate staining pattern by immunofluorescence and immunoelectron microscopy^{18,23}.

A role for CS binding to hepatic HSPGs *in vivo*

The density and structure of CS on the parasite surface make it an ideal ligand for trapping the parasite in the liver. When radiolabeled CS is intravenously injected into mice, it is rapidly cleared from the blood circulation and found in the liver²⁴. Competition experiments with the physiological ligands for hepatic HSPGs, namely lipoprotein remnant particles and lactoferrin (reviewed in Ref. 25), show that CS clearance is delayed in mice preinjected with lactoferrin and in mice with high circulating levels of remnant lipoproteins²⁶.

To determine whether or not sporozoites are captured in the liver by the same mechanism as CS, mice with high circulating levels of remnant lipoproteins were used for experiments *in vivo*. Forty hours after injection with sporozoites, mice with high remnant lipoprotein levels had 8–10-fold fewer parasites developing in the liver than control mice²⁶, suggesting that the binding between the abundant hepatocyte HSPGs and the dense CS coat of the parasite is critical for the arrest of the parasite in the liver.

As mentioned earlier, one confounding issue is that the hepatocyte microvilli are within the space of Disse behind the fenestrated endothelial cells. Perhaps the long GAG chains of the HSPGs protrude through the endothelial fenestrae into the sinusoids and capture circulating particles, such as lipoprotein remnants and *Plasmodium* sporozoites (Fig. 2). Lyon and Gallagher have shown that the bulk of hepatic HSPG sulfation is along the distal third of the molecule, the part that would be in the sinusoidal lumen if this hypothesis is correct²⁷. An

alternative possibility is that the fenestrae are not rigid structures and their diameter can increase during blood circulation, allowing for direct contact between sporozoites and hepatocyte microvilli.

Other sporozoite surface proteins

Sporozoite arrest in the liver is the first step of hepatocyte invasion; once in the liver, sporozoites must enter hepatocytes and develop into EEF. In contrast to what is known about merozoite invasion of red cells, little is known about how sporozoites invade hepatocytes. Thus far, only one other sporozoite surface protein has been cloned and sequenced: the 78-kDa sporozoite threonine- and asparagine-rich protein STARP (Ref. 28). STARP has a high degree of structural conservation in different *Plasmodium falciparum* isolates, as well as a high degree of homology with the gene from the closely related *Plasmodium reichenowi*, suggesting that it is important in the life cycle of the parasite²⁹. Whether this protein is involved in sporozoite invasion of hepatocytes and, if so, what role it plays is not yet known.

Erythrocyte invasion by *Plasmodium*

When merozoites are released from erythrocytes they are in close proximity to their target cells: uninfected erythrocytes. If the target erythrocyte is in direct contact with the red cell from which the merozoite emerges, the parasite may only be extracellular briefly. Finding host cells is more difficult for *Plasmodium vivax* merozoites, which primarily, if not exclusively, invade reticulocytes³⁰⁻³². However, even for *P. vivax*, the extra-cellular journey of a merozoite is much less complex than that of a sporozoite, which must travel from the site of inoculation in the skin to the hepatocyte.

Erythrocyte invasion by malaria parasites is a multistep process that involves attachment and reorientation, apical end binding and junction formation, parasitophorous vacuole membrane (PVM) formation and internalization (reviewed in Ref. 2) (Fig. 3). The apical organelles (micronemes and rhoptries), as well as other organelles called dense granules, seem to be involved in several of these steps. How this process is controlled is not understood, but it may occur as a defined sequence, in which one event occurs as a consequence of the previous step, or as a cascade catalyzed by a single trigger.

Attachment and orientation

After release from ruptured schizonts, merozoites come into contact with erythrocytes and reorient so that ligands expressed at the apical end of the parasite are able to interact with erythrocyte receptors. Reorientation appears to involve attachment, dissociation and reattachment as the merozoite appositions itself in the correct orientation on the erythrocyte membrane (reviewed in Ref. 2). The interaction between the erythrocyte membrane and the merozoite surface must be of low affinity because the attachment is reversible. The specific receptors and ligands that mediate this interaction have not been identified; however, it seems likely that the re-orientation process involves an abundant surface protein of merozoites, and one possible candidate is merozoite surface protein 1 (MSP-1)^{33,34}. Full-length MSP-1 from supernatants of *P. falciparum* erythrocytic stage cultures binds specifically to the sialic acid residues of glycoporphins on erythrocytes, but this binding is of

low affinity³⁵. However, others have found that MSP-1 is processed at about the time of merozoite release from erythrocytes and a 19-kDa cysteine-rich glycosyl-phosphatidylinositol (GPI)-linked glycosylated fragment is left on the merozoite surface as it enters the erythrocyte³⁶. How then does full-length MSP-1 participate in invasion? Possibly, sequential binding and proteolysis produce a gradient of contact that facilitates directionally oriented movement towards the apical pole³⁷. Although it is not clear if MSP-1 is involved in attachment and orientation of the merozoite to the erythrocyte, antibodies raised against this protein inhibit merozoite invasion of erythrocytes³⁶ and confer protection^{38,39}, suggesting that MSP-1 participates in the invasion process.

Apical end binding and junction formation

To date, all of the ligands involved in apical end binding of the merozoite to the erythrocyte have been localized to the micronemes. These ligands are the 175-kDa erythrocyte-binding antigen (EBA-175) of *P. falciparum*⁴⁰⁻⁴² and the Duffy binding proteins of *P. vivax*^{43,44} and *Plasmodium knowlesi*^{45,46}. Another group of proteins, the reticulocyte-binding proteins of *P. vivax*⁴⁷, is seen at the apical end of the merozoite but its precise localization in the merozoite has not yet been described.

Interestingly, the CS and SSP2/TRAP proteins of sporozoites, which both bind to hepatocytes, have been localized to the micronemes of sporozoites^{18,48,49}. Although there are no obvious similarities between the gene structure of the merozoite and sporozoite ligands, the co-localization of these binding proteins in the micronemes suggests a common subcellular trafficking and sequestration route. As both CS and SSP2/TRAP are found on the surface of the sporozoite, the micronemes may function as a constitutive intermediate compartment in the trafficking of surface proteins. However, neither EBA-175 nor the Duffy binding proteins have been visualized on the surface of merozoites. Visualizing these ligands on the surface of *P. falciparum* and *P. vivax* merozoites may not be easy because trafficking to the surface may depend on the viability of merozoites and the isolation of viable merozoites of these species is not yet feasible. However, the Duffy binding protein of *P. knowlesi*, whose merozoites can be isolated in a viable state, has not been visualized on the merozoite surface. Another possibility is that the micronemes function as a regulated secretory granule, releasing their contents at a specific time during invasion. Both CS and SSP2/TRAP are found on the surface of sporozoites before they contact target cells, in contrast to EBA-175 and the Duffy binding proteins, which are not found on the merozoite surface and may only be secreted after contact with target cells. Perhaps the signals regulating protein secretion by the micronemes vary depending on the stage of the parasite.

The binding of the merozoite ligand to the erythrocyte receptor leads to the formation of a tight junction that can be visualized as an electron-dense zone beneath the erythrocyte membrane (Ref. 50 and reviewed in Ref. 2). The formation of a tight junction appears to be the committed step in the invasion process. The tight junction starts as a localized patch and transforms into a circumferential interaction between the apical end of the merozoite and the invaginated erythrocyte membrane. This moves as a band around the merozoite and fuses as the posterior end of the merozoite enters the erythrocyte, closing the orifice not unlike an iris diaphragm. The merozoite is now within the erythrocyte surrounded by the PVM (Fig. 3).

Parasitophorous vacuole formation

As mature erythrocytes are incapable of receptor-mediated endocytosis⁵¹, the driving force behind PVM formation must be provided by the merozoites. During PVM formation, the contents of the apical organelles are released onto the erythrocyte membrane. Rhoptries are a pair of electron-dense, teardrop-shaped organelles connected by ducts leading to the apical end of the parasite. Several rhoptry proteins have been identified and characterized⁵²⁻⁵⁵, and one of these, a 110-kDa protein, is part of a rhoptry protein complex that is secreted into the erythrocyte membrane during invasion⁵⁶. However, it is not known whether the PVM is formed as a result of the insertion of apical organelle proteins into the erythrocyte membrane, which could lead to its deformation, or if the membrane is deformed by the active invasion of the merozoite, which then leads to the release of rhoptry contents.

The protein composition of the PVM is distinct from that of the erythrocyte membrane. Rhoptry proteins are inserted into the PVM (Ref. 56) and several major erythrocyte proteins are excluded from the PVM (Refs 57,58). However, in experiments using fluorescent lipophilic tracers, it has been found that the lipid composition of the PVM is indistinguishable from that of the normal erythrocyte membrane^{2,59}.

After rhoptry discharge, dense granules from the apical and lateral surfaces of the merozoite discharge their contents into the PVM (Refs 50,60), and the trailing end of the invaginated erythrocyte membrane fuses. Dense granules are spherical organelles that are larger than a microneme but smaller than the rhoptry⁶⁰. A parasite protein that has been localized to the dense granules is the 155-kDa ring-infected erythrocyte surface antigen (RESA). RESA is released from the dense granules^{49,61} and then, by as yet undefined mechanisms, is transported across the PVM and binds spectrin in the erythrocyte cytoskeleton^{62,63}.

Parasite proteases are thought to play a role at various steps of the invasion process⁶⁴⁻⁶⁶, perhaps by local disruption of the erythrocyte membrane. Elegant studies by Braun Breton and co-workers have shown that in *Plasmodium chabaudi*, the parasite p68 serine protease cleaves band 3 protein in the erythrocyte membrane, leading to successful invasion⁶⁷. Cleavage of band 3 may modify the erythrocyte cytoskeleton so that it can deform during merozoite entry.

Receptor and ligands

The important historical observations and data that have implicated particular proteins as receptors and ligands for invasion will not be described here. The initial observation that *P. knowlesi* and *P. vivax* do not invade Duffy-negative erythrocytes has led to the identification of the Duffy blood group antigen as a putative invasion receptor for these parasites (reviewed in Ref. 2). In the case of *P. falciparum*, invasion studies with mutant and enzymatically treated erythrocytes *in vitro* have demonstrated that the sialic acid residues of glycophorins are receptors for invasion⁶⁸⁻⁷⁰.

More recently, parasite ligands for the Duffy blood group antigens and glycophorin A have been discovered using an erythrocyte-binding assay^{40,45} and supernatants from labeled parasite cultures. The 140-kDa and 135-kDa Duffy binding proteins of *P. vivax*⁴³ and *P. knowlesi*⁴⁵, respectively, and EBA-175 of *P. falciparum*⁴⁰ were identified in this way. There

are striking similarities between the gene structures of the Duffy binding proteins and EBA-175 and they have been proposed to be members of an erythrocyte-binding family of proteins⁷¹. They all have three exons encoding transmembrane and cytoplasmic domains and they share 5' and 3' cysteine-rich domains that contain conserved cysteine motifs.

The 5' cysteine-rich region of these erythrocyte-binding proteins, designated region II, has been identified as the receptor-binding domain^{42,72}. Within region II, the cysteines and some aromatic residues are strictly conserved and probably constitute the erythrocyte-binding motif, while the intervening sequences determine the specificity of the receptor binding⁷¹. In the case of EBA-175, binding is specific for sialic acids linked 2,3 on O-linked tetrasaccharides⁷³ and requires both sialic acids and the protein backbone of glycophorin A for binding to erythrocytes⁴².

Conservation of functional domains in evolutionarily distant species was previously described for the CS protein, in which the only regions conserved among species were those shown to bind hepatocytes^{15,74}. As the Duffy binding proteins of *P. knowlesi* appear to be required for junction formation during invasion, it is likely that erythrocyte-binding proteins in other *Plasmodium* species have the same role.

The specificities of the erythrocyte-binding proteins do not account for all of the host cell preferences observed *in vivo*. *P. vivax* merozoites preferentially invade reticulocytes^{30,31,47}, even though the Duffy antigen is expressed on both reticulocytes and mature erythrocytes. The reasons for this preferential invasion of reticulocytes is not understood; however, two erythrocyte-binding proteins of 250 kDa and 280 kDa that preferentially bind reticulocytes have recently been identified⁴⁷. These proteins are expressed at the invasive, apical end of merozoites and bind both Duffy-positive and Duffy-negative erythrocytes. Whether these proteins are the parasite ligands responsible for the preferential invasion of reticulocytes remains to be determined.

There are undoubtedly many more, as yet unidentified, parasite proteins involved in the invasion process. In addition, there is evidence that these parasites use alternative pathways to invade erythrocytes. For example, the *P. vivax* Duffy binding protein does not bind to Saimiri monkey erythrocytes, even though these erythrocytes are invaded by *P. vivax* merozoites⁴³. *P. falciparum* can also use alternative pathways for entering erythrocytes. All strains of *P. falciparum* studied to date primarily use sialic acid residues on glycophorin A as receptors because neuraminidase treatment of erythrocytes greatly affects the invasion efficiencies of these strains *in vitro*⁶⁸. However, certain strains of *P. falciparum* can be selected to invade and develop in neuraminidase-treated erythrocytes vs and it has been observed that some strains of *P. falciparum* (Dd2 and FCR3/A2) can invade after trypsin treatment of erythrocytes⁷⁶; trypsin cleaves glycophorin A. These alternative pathways of invasion possibly indicate a redundancy required to give a selective advantage to parasites for survival in a polymorphic human population.

Conclusions

Invasion of host cells by malaria parasites is a tightly controlled multistep process that occurs over a short period of time in specific host compartments (i.e. blood-stream, liver sinusoids and mosquito midgut). In the case of erythrocyte invasion by merozoites, this process has been visualized and involves a defined sequence of events, which includes recognition and attachment, reorientation, junction formation, PVM formation and internalization. Although sporozoite entry into hepatocytes has not been visualized in this detail, the process may involve some of the same steps⁷⁷.

Microscopic observations, as well as biochemical studies of merozoite entry into host cells, have shown that the micronemes contain proteins required for invasion, and that the contents of the rhoptries and dense granules are released during invasion. However, there are no known apical organelle proteins that are identical, or even similar, in different stages of the same *Plasmodium* species. Perhaps this strategy enables the parasite to better avoid the immune response of the host. Despite these differences, within a given stage of the parasite, but among evolutionarily distant species, there is conservation of protein domains that have been shown to function during cell invasion, for example, region II-plus of CS and SSP2/TRAP and region II of the erythrocyte-binding proteins of *P. vivax*, *P. knowlesi* and *P. falciparum*. Identifying the receptors and ligands involved in parasite invasion is important; however, experiments aimed at understanding the overall mechanisms of invasion are also of interest and perhaps of wider significance. It is hoped that future work will elucidate some of the common mechanisms utilized by apicomplexan parasites for entry into host cells.

Acknowledgments

P.S. is supported by an NIH Physician Scientist Award (K11 AI-01175).

References

1. Vanderberg J, Chew S, Stewart MJ. J Protozool. 1990; 37:528–536. [PubMed: 2086782]
2. Ward, G., Chitnis, CE., Miller, LH. Strategies for Intracellular Survival of Microbes. Russell, D., editor. Saunders; 1994. p. 155-190.
3. Khusmith S, Sedegah M, Hoffman SL. Infect Immun. 1994; 62:2979–2983. [PubMed: 8005684]
4. Ungureanu E, et al. Trans R Soc Trop Med Hyg. 1976; 70:482–483. [PubMed: 320727]
5. Ponnudurai T, et al. Trans R Soc Trop Med Hyg. 1991; 85:175–180. [PubMed: 1887464]
6. Shin SCJ, Vanderberg JP, Terzakis JA. J Protozool. 1982; 29:448–454. [PubMed: 6752394]
7. Wisse E, et al. Hepatology. 1985; 5:683–692. [PubMed: 3926620]
8. Meis JFGM, et al. Parasitology. 1983; 86:231–242. [PubMed: 6343960]
9. Vreden SGS, et al. Infect Immun. 1993; 61:1936–1939.
10. Seguin MC, Ballou WR, Nacy CA. Immunol. 1989; 143:1716–1722.
11. Vanderberg JP. Parasitol Today. 1995; 11:24.
12. Nussenzweig, V., Nussenzweig, RS. Advances in Immunology. Dixon, FJ., editor. Academic Press; 1989. p. 283-334.
13. Hedstrom RC, et al. Bull WHO. 1990; 68:152–157. [PubMed: 2094580]
14. Robson KJH, et al. Nature. 1988; 335:79–82. [PubMed: 3045563]
15. Cerami C, et al. Cell. 1992; 70:1021–1035. [PubMed: 1326407]
16. Robson KJH, et al. EMBO J. 1995; 14:3883–3894. [PubMed: 7664729]

17. Muller HM, et al. EMBO J. 1993; 12:2881–2889. [PubMed: 8392935]
18. Rogers WO, et al. Proc Natl Acad Sci U S A. 1992; 89:9176–9180. [PubMed: 1409621]
19. Dame JB, et al. Science. 1984; 225:593–599. [PubMed: 6204383]
20. Sinnis P, et al. J Exp Med. 1994; 180:297–306. [PubMed: 8006589]
21. Pancake SJ, et al. J Cell Biol. 1992; 6:1351–1357.
22. Frevert U, et al. J Exp Med. 1993; 177:1287–1298. [PubMed: 8478608]
23. Cowan G, et al. Lancet. 1992; 339:1412–1413. [PubMed: 1350818]
24. Cerami C, et al. J Exp Med. 1994; 179:695–701. [PubMed: 8294876]
25. Mahley, RW., et al. Biology of α_2 -Macroglobulin, its Receptor, and Related Proteins. Borth, W., et al., editors. Vol. 737. New York Academy of Sciences; 1994. p. 39–52.
26. Sinnis P, et al. J Exp Med. 1996; 184:945–954. [PubMed: 9064354]
27. Lyon M, Deakin JA, Gallagher JT. J Biol Chem. 1994; 269:11208–11215. [PubMed: 8157650]
28. Fidock DA, et al. Mol Biochem Parasitol. 1994; 64:219–232. [PubMed: 7935600]
29. Fidock DA, et al. Mol Biochem Parasitol. 1994; 67:255–267. [PubMed: 7870130]
30. Kitchen SF. Am J Trop Med Hyg. 1938; 18:347–359.
31. Mons B, et al. Exp Parasitol. 1988; 66:183–188. [PubMed: 3294025]
32. Barnwell JW, et al. J Exp Med. 1989; 169:162–167.
33. Holder AA, et al. Mol Biochem Parasitol. 1985; 14:293–303. [PubMed: 3887156]
34. Miller LH, et al. Mol Biochem Parasitol. 1993; 59:1–14. [PubMed: 8515771]
35. Perkins ME, Rocco LJ. J Immunol. 1988; 141:3190–3196. [PubMed: 2459245]
36. Blackman MJ, et al. Exp Med. 1990; 172:379–382.
37. Barnwell JW, Galinski MR. Res Immunol. 1991; 142:666–671. [PubMed: 1817296]
38. Siddiqui WA, et al. Proc Natl Acad Sci U S A. 1987; 84:3014–3018. [PubMed: 3554251]
39. Etlinger HM, et al. Infect Immun. 1991; 59:3498–3503. [PubMed: 1894356]
40. Camus D, Hadley TJ. Science. 1985; 230:553–556. [PubMed: 3901257]
41. Sim BKL, et al. J Cell Biol. 1990; 111:1877–1884. [PubMed: 2229177]
42. Sim BKL, et al. Science. 1994; 264:1941–1944. [PubMed: 8009226]
43. Werthheimer SP, Barnwell JW. Exp Parasitol. 1989; 69:340–350. [PubMed: 2680568]
44. Fang X, et al. Mol Biochem Parasitol. 1991; 44:125–132.
45. Haynes JD, et al. Exp Med. 1988; 367:1873–1881.
46. Adams JH, et al. Cell. 1990; 63:141–153. [PubMed: 2170017]
47. Galinski M, et al. Cell. 1992; 69:1213–1226. [PubMed: 1617731]
48. Fine E, et al. Am J Trop Med Hyg. 1984; 30:220–226.
49. Aikawa M, et al. Bull WHO. 1990; 68:165–171.
50. Bannister LH, et al. Parasitology. 1975; 71:483–491. [PubMed: 1202413]
51. Zweig S, Singer SJ. J Cell Biol. 1979; 80:487–491. [PubMed: 457755]
52. Petersen MG, et al. Mol Cell Biol. 1989; 9:3151–3154.
53. Ridley RG, et al. Mol Biochem Parasitol. 1990; 41:125–134. [PubMed: 2200961]
54. Brown HJ, Coppel RL. Mol Biochem Parasitol. 1991; 49:99–110. [PubMed: 1775161]
55. Saul A, et al. Mol Biochem Parasitol. 1992; 50:139–150. [PubMed: 1542308]
56. Sam-Yellowe TY, Shio H, Perkins ME. J Cell Biol. 1988; 106:1507–1513. [PubMed: 2453514]
57. Atkinson CT, et al. J Cell Biol. 1987; 45:192–199.
58. Dluzewski AR, et al. J Cell Sci. 1989; 92:691–699. [PubMed: 2532219]
59. Ward GE, Miller LH, Dvorak JA. J Cell Sci. 1993; 106:237–248. [PubMed: 8270628]
60. Torii M, et al. Infect Immun. 1989; 57:3230–3233. [PubMed: 2777381]
61. Culvenor JG, Day KP, Anders RF. Infect Immun. 1991; 59:1183–1187. [PubMed: 1997422]
62. Foley M, et al. Mol Biochem Parasitol. 1991; 46:137–148. [PubMed: 1852169]
63. Ruangjirachuporn W, et al. Exp Parasitol. 1991; 73:62–72. [PubMed: 2055301]
64. Barale JC, et al. Res Immunol. 1991; 142:672–681. [PubMed: 1817297]

65. Braun Breton C, Pereira da Silva LH. *Parasitol Today*. 1993; 9:9296–9300.
66. McKerrow JH, et al. *Annu Rev Microbiol*. 1993; 47:821–853. [PubMed: 8257117]
67. Braun Breton C, et al. *Proc Natl Acad Sci U S A*. 1992; 89:9647–9651.
68. Pasvol G. *Philos Trans R Soc London*. 1984; 307:189–200. [PubMed: 6151683]
69. Hadley, TJ., Miller, LH. Protein Blood Group Antigens of the Human Red Cell. Agree, PC., Cartron, JC., editors. Johns Hopkins University Press; 1992. p. 228-245.
70. Haynes, JD. Current Opinion in Haematology. Adamson, JW., editor. Current Science; 1993. p. 79-89.
71. Adams JH, et al. *Proc Natl Acad Sci U S A*. 1992; 89:7085–7089. [PubMed: 1496004]
72. Chitnis CE, Miller LH. *J Exp Med*. 1994; 180:497–506. [PubMed: 8046329]
73. Orlandi PA, et al. *J Cell Biol*. 1992; 116:901–911. [PubMed: 1310320]
74. Aley SB, et al. *J Exp Med*. 1986; 164:1915–1921. [PubMed: 3023519]
75. Dolan SA, Miller LH, Wellems TE. *J Clin Invest*. 1990; 86:618–624. [PubMed: 2200806]
76. Dolan SA, et al. *Mol Biochem Parasitol*. 1994; 64:55–63. [PubMed: 8078523]
77. Russell DG. *Parasitology*. 1983; 87:199–209. [PubMed: 6646806]

Box 1**Glossary**

CS: Circumsporozoite protein.

Duffy blood group antigens: A group of antigens on the erythrocyte surface used for blood typing.

EBA-175: 175-kDa erythrocyte-binding antigen of *Plasmodium falciparum*.

EEF: Exoerythrocytic forms of *Plasmodium*.

GAGs: Glycosaminoglycan chains of HSPG.

HepG2 cells: A hepatoma cell line permissive for *Plasmodium berghei* sporozoite invasion and development.

HSPGs: Heparan sulfate proteoglycans.

Kupffer cells: Actively phagocytic cells that line the liver sinusoids.

MSP-I: Merozoite surface protein 1.

PVM: Parasitophorous vacuole membrane.

Region II: Receptor-binding domain of erythrocyte-binding proteins of merozoites.

Region II-plus: A cell-adhesive motif that occurs in both CS and SSP2/TRAP with homology to the type I thrombospondin repeats and which binds to sulfated glycoconjugates.

RESA: 155-kDa ring-infected erythrocyte surface antigen.

Space of Disse: The space between hepatocytes and the sinusoid lining that is continuous with the sinusoid lumen.

SSP2/TRAP: Sporozoite surface protein 2 or thrombospondin-related adhesion protein.

Questions for future research

- What are the signaling events that trigger release of the contents of apical organelles during invasion?
- Once in the circulation of the vertebrate host, the sporozoite is probably cleared by hepatocytes via binding of region II-plus motifs in CS (and perhaps SSP2/TRAP) to hepatocyte HSPGs. However, sporozoites are normally injected intradermally, but not intravenously, by mosquitoes. How do the sporozoites get from the site of injection into the circulation?
- Although blood smears from malarious patients and *Plasmodium falciparum* cultures *in vitro* frequently reveal several multiply infected erythrocytes, different stages of the parasite are never seen in the same erythrocytes. This suggests that a merozoite can not invade an already infected erythrocyte. In what way has the intracellular parasite changed its host erythrocyte, such that a merozoite can no longer enter?
- Sporozoites and merozoites have surface molecules with transmembrane domains and GPI-anchored surface molecules. Do these differently anchored proteins on the surface of the parasite have distinct roles in invasion and motility?
- In what way is the membrane of the parasitophorous vacuole different from the host cell membrane, and how do these differences ensure the survival of the intracellular parasite?

Fig. 1.

Life cycle of the malaria parasite. (1) Sporozoites are injected into a vertebrate host during the blood meal of a female anopheline mosquito and rapidly invade hepatocytes. One sporozoite can develop into 20 000 merozoites, which rupture from the hepatocyte, enter the bloodstream and invade erythrocytes. (2) In the asexual erythrocytic cycle, merozoites invade erythrocytes and mature from ring stages to schizonts within these cells in 48–72 h, the time varying with the species of malaria parasite. (3) Some erythrocytic stages differentiate into gametocytes, which are infective for mosquitoes. Fertilization occurs in the mosquito midgut, and within 24 h zygotes transform into ookinetes, which penetrate the midgut and form oocysts. Sporozoites rupture from these oocysts and invade the salivary glands of the mosquito from where they are injected into a vertebrate host.

Fig. 2.

Diagrammatic representation of direct interaction between sporozoites and hepatocytes.

Heparan sulfate proteoglycans (HSPGs) are shown as branched structures on the hepatocyte membrane, protruding through the open fenestrae of the endothelial cells into the sinusoidal lumen. Sporozoites, shown as black half-moon structures, are captured by these HSPGs and then invade hepatocytes.

Fig. 3. Schematic representation of the major morphological events of merozoite invasion. On top, the merozoite is depicted in the attachment and orientation phase, followed by binding, junction formation and rhoptry discharge. The merozoite is then shown entering the parasitophorous vacuole as the junction between the merozoite and the erythrocyte moves towards the posterior end of the merozoite. The parasite is then completely intracellular and flattens with its cytoplasm remaining thick only at the edges, giving the appearance of a 'ring'.

Table 1

Region II-plus sequences of CS and homologous sequences of SSP2/TRAP from several species of malaria parasites^a

Malaria parasite	Region II-plus sequences ^b
Circumsporozoite protein	
<i>Plasmodium falciparum</i>	EWSPCSVTCGNGIQV RIK
<i>Plasmodium vivax</i>	EWTPCSVTCGVGV RVR RRR
<i>Plasmodium malariae</i>	EWSPCSVTCGSGIR ARRK
<i>Plasmodium knowlesi</i>	EWTPCSVTCGNGV RI RRK
<i>Plasmodium cynomolgi</i>	EWSPCSVTCGKRV RM RRK
<i>Plasmodium brasilianum</i>	EWSPCSVTCGKRV RM RRK
<i>Plasmodium berghei</i>	EWSPCSVTCGSGIR ARRK
<i>Plasmodium yoelii</i>	EWSQCNVTCGSGIR V RRK
SSP2/TRAP	
<i>Plasmodium falciparum</i>	EWSPCSVTCGKG TR SR KR
<i>Plasmodium yoelii</i>	EWSECSTTCDEG RK IRRR

^a Abbreviations: CS, circumsporozoite protein; SSP2, sporozoite surface protein 2; TRAP, thrombospondin-related adhesion protein.

^b The highly conserved cysteines and the downstream basic residues that are essential for binding to heparan sulfate proteoglycans are marked in bold.