bmj.com news roundup

Full versions of these stories are available at: bmj.com/content/vol330/issue7491/#NEWS_ROUNDUP

Low dose aspirin helps prevent ischaemic stroke in women

A large study examining the benefits of low dose aspirin in the primary prevention of cardiovascular disease in women found a significant benefit only for stroke. The findings, which were presented at the American College of Cardiology meeting in Florida this week, will be published in the *New England Journal of Medicine* on 31 March.

In the study researchers randomised 39 876 healthy women over 45 years to receive 100 mg of aspirin every other day, or placebo, and monitored them for 10 years.

Compared with placebo, aspirin had no significant effect on the risk of fatal or non-fatal heart attacks (1.02; 0.84 to 1.25; $P=0.83$) or death from cardiovascular causes (0.95; 0.74 to 1.22; $P=0.68$).

Researchers found, however, that there was a 17% reduction in the risk of stroke in the aspirin group, compared with the placebo group (0.83; 0.69 to 0.99; $P=0.04$), owing to a 24% reduction in the risk of ischaemic stroke (0.76; 0.63 to 0.93; $P=0.009$) that was offset by only a small and non-significant increase in the risk of haemorrhagic stroke (1.24; 0.82 to 1.87; $P=0.31$).

Scott Gottlieb *New York*

Japanese study is more evidence that MMR does not cause autism

A Japanese research study has provided the strongest proof yet that the measles, mumps, and rubella (MMR) vaccination does not cause autism, by showing that rates of autism in Japan continued to rise even after the triple vaccine was withdrawn.

The latest study, of 31 426 children in the Japanese city of Yokohama, examined the incidence of autism between 1988 and 1996, a period when uptake of the MMR vaccination steadily declined before being

Doctors question whether all blood transfusions are effective and necessary

Doctors have questioned the value of spending a large chunk of the health budget to ensure blood for transfusions is safe while there is a severe lack of evidence that the procedure is effective.

The cost of supplying blood to the United Kingdom grew from £250m (\$480m; €365m) in 1995 to £500m in 2004, a conference held last week at the Royal College of Physicians of Edinburgh was told. The increase is due mainly to new tests and processes to reduce the risk of transmitting infection.

There was particular concern about the introduction of nucleic acid testing to supplement existing tests for hepatitis C which cost between \$5.8m and \$8.4m for each quality adjusted life year (QALY) resulting from the avoidance of hepatitis C.

Prion reduction filtration to reduce the risk of transmitting variant Creutzfeldt-Jakob disease could add a further £100m to the United Kingdom's blood service bill.

The emphasis on safety had not, however, been matched by an equal emphasis on effectiveness. Randomised controlled trials looking at the triggers for transfusion involved fewer than 2000 people. Half of these patients were in one trial that showed no benefit from transfusion (*New England Journal of Medicine* 1999;340:409-17). In this study mortality was higher in patients under 55 who were given more blood (see above).

Richard Smith *Edinburgh*

withdrawn in 1993 and replaced by single vaccines (published online in the *Journal of Child Psychology and Psychiatry*, DOI: 10.1111/j.1469-7610.2005.01425.x).

Yet the incidence of autism continued to rise, from 48 cases per 10 000 children born in 1988 to 117.2 per 10 000 born in 1996.

The controversy over the vaccine was triggered by a 1998 research paper by gastroenterologist Andrew Wakefield (*Lancet* 1998;351:637-41).

Andrew Cole *London*

GPs' college issues guidance on hepatitis A and B vaccination

Injecting drug users should be vaccinated against hepatitis A, and all users of illegal drugs should be vaccinated against hepatitis B, the Royal College of General Practitioners recommended this week. The vaccinations should be carried out without waiting for prevaccination tests for hepatitis A or B.

It is estimated that 21% of injecting drug users in England and Wales have evidence of past or current hepatitis B infection. Poor uptake of vaccination in drug users has led experts now to advise that vaccination is done before waiting for test results for hepatitis B.

All drug users should be vaccinated with hepatitis B vaccine because non-injectors often become injectors, says the College. The accelerated schedule for hepatitis B vaccination should be used, under which the vaccination is given at 0, 7, and 21 days.

Robert Short *London*

Guidance for Hepatitis A and B Vaccination of Drug Users in Primary Care and Criteria for Audit is available at www.rcgp.org.uk/drug/

Treatment for heart disease must improve in deprived areas

The government has made major steps in the treatment of heart disease in England and Wales, according to a report by the NHS inspection body, the Healthcare Commission. But it still needs to do more to target people most at risk of heart disease, particularly those in areas of social or economic deprivation, where rates of heart disease are still higher than normal.

The report looks at progress on the national service framework for heart disease, which was launched in 2000. People who have a heart attack are getting treatment more quickly now. Eighty five per cent of patients were given thrombolytic drugs in 2004 within the target time of 30 minutes of arriving at the hospital; in 2002 the figure was 59%. Of patients who were discharged from hospital in 2004 after a heart attack, nearly 94% receive statins, compared to 84% in 2002.

Lynn Eaton *London*

Getting to the Heart of it, Coronary Heart Disease in England: A Review of Progress Towards National Standards is available at www.healthcarecommission.org.uk