UK and US governments must monitor Iraq casualties

Owen Dyer *London*

An international group of public health experts has called on the US and UK governments to commission an independent inquiry into Iraqi war related casualties.

"We believe that the joint US-UK failure to make any effort to monitor Iraqi casualties is, from a public health perspective, wholly irresponsible," says the public statement, signed by 23 experts in public health and epidemiology from the United Kingdom, the United States, Canada, Spain, and Australia. "We have waited too long for this information."

The British government's reliance on Iraqi Ministry of Health figures is "unacceptable," argue the signatories. These figures "are likely seriously to underestimate casualties," as they cover only April 2004 to October 2004, include only violent deaths reported through the health system, and "do not allow for reliable attribution between different causes of death and injury."

The Iraqi Ministry of Health lists 3853 civilian deaths and 15 517 injuries during the six months, casualties of both military and insurgent actions.

Professor Sheila Bird, of the

Medical Research Council's biostatistics unit at Cambridge, one of the signatories, said the statement was a response to the government's lack of action since November last year, when the *Lancet* published research showing that Iraq had about 98 000 excess deaths (95% confidence interval 8000 to 194 000) since the April 2003 invasion (2004; 364:1857).

"There were a number of comments from the Foreign Office in the press in November, suggesting that they would work to improve the methodology of data collection. But nothing has happened since," she said.

The foreign secretary, Jack Straw, told parliament in November that the government did not accept the central conclusion of the *Lancet* study, which "was carried out under exceptionally difficult conditions which in particular restricted the size of the samples surveyed."

But the public health experts' statement retorts that "the obvious answer to removing uncertainties that remain is to commission a larger study with full official support and assistance, but scientific independence."

The British and American governments contend that they

have no legal responsibility to count civilian casualties. Mr Straw told parliament, "The *Lancet* study suggests that there is an obligation deriving from article 27 of the Fourth Geneva Convention for the multinational force itself to have a reckoning of the number of civilian casualties it has caused. There is nothing in article 27, or elsewhere in the Fourth Geneva Convention, to support this suggestion."

A Foreign Office spokesman told the *BMJ*, "We continue to feel that the Iraqi Ministry of Health figures are the best available in an uncertain situation, being based on an actual head count instead of extrapolation. In the current security climate, more accurate research is not

feasible." Should the security situation improve, he said, a final tally would be a matter for the Iraqi government.

Professor Klim McPherson, a public health epidemiologist at Oxford University, was the instigator and first signatory of the statement. He said: "Basically, this is a response to the government's continuing procrastination. A group of us sent a letter to the prime minister in December under the auspices of the Count the Casualties campaign, an initiative of Medact and Iraq Body Count, which essentially got the brush-off, so we decided to write publicly." (See p 550.) □

The full statement is in News on bmj.com

An Iraqi girl screams after her parents are killed by US soldiers who fired on their car when it failed to stop during a dusk patrol

Low cost measures could save lives of three million babies a year

Rebecca Coombes *London*

The lives of nearly three million of the four million newborn babies who die worldwide each year could be saved by low cost, low tech interventions, a series of articles published by the *Lancet* shows.

Ninety nine per cent of the four million deaths occur in poor countries, although almost all published research relates to the 1% in rich countries.

The first paper in the series

of articles, all published in the 3 March issue (www.thelancet.com), looks at when, where, and why newborn babies are dying and finds that 75% of neonatal deaths occur in the first week of life, with the highest risk of death on the first day of life. Two thirds of deaths occur in 10 countries, which include Afghanistan, Ethiopia, and Pakistan.

The main causes of neonatal deaths are infections (36%), premature birth (28%), and asphyxia (23%). Neonatal tetanus, a preventable illness that is virtually unknown in the developed world, kills half a million babies each year.

A key message of the *Lancet* series is that interim approaches are already available in the community that would save newborns. For example, between 60% and 80% of neonatal deaths

are in babies with a low birth weight, and many could be saved with simple support such as keeping them warm and breast feeding.

Researchers estimate that a package of 16 low tech interventions could be provided to 90% of women and babies in poor countries at a cost of just \$1 (£0.5; €0.8) extra per inhabitant per year.

The 16 interventions set out in the papers include tetanus vaccinations for pregnant women, promoting clean delivery, prompt and exclusive breast feeding, extra care for low weight babies, and antibiotics for neonatal infection. None of the interventions involves expensive drugs or complex technology.

Dr Zulfiqar Bhutta, professor of paediatrics and child health at

Aga Khan University, Karachi, said it was vital to improve care practices in the home, where most neonatal deaths occur. More than half of women in Africa and south Asia give birth at home without a skilled attendant.

"We know the interventions are simple, so how come they are not there? You have to look at where the deaths occur. At home there is a sense of fatalism around newborns that nothing can be done. We need to get the message across that small babies can be saved. Health systems may be bad, but families are not mobilised sufficiently to go out and seek help," he said.

A major barrier to progress in reducing mortality among newborns has been the false belief that expensive, high tech solutions are needed to bring death rates down, he said. □