Published in final edited form as:

Depress Anxiety. 2017 August ; 34(8): 761–769. doi:10.1002/da.22669.

Transdiagnostic Emotional Vulnerabilities Linking Obsessive-Compulsive and Depressive Symptoms in a Community-Based Sample of Adolescents

Gregory S. Chasson, Ph.D.^{a,*}, Mariel S. Bello, B.S.^b, Alexandria M. Luxon, M.A.^c, Trevor A. A. Graham, B.A.^d, and Adam M. Leventhal, Ph.D.^{b,d}

^aIllinois Institute of Technology, Department of Psychology, Chicago, IL, USA

^bUniversity of Southern California, Department of Psychology, Los Angeles, CA, USA

^cTowson University, Department of Psychology, Towson, MD, USA

^dUniversity of Southern California Keck School of Medicine, Department of Preventive Medicine, Los Angeles, CA, USA

Abstract

Background—Transdiagnostic emotional vulnerabilities are suspected to underlie psychopathologic comorbidity but have received little attention in adolescent emotional pathology literature. We examined distress tolerance, anxiety sensitivity, and anhedonia as concomitant transdiagnostic mechanisms that account for (i.e., statistically mediate) the covariance between adolescent obsessive-compulsive disorder (OCD) and major depressive disorder (MDD) symptoms.

Method—Data on MDD, OCD, and the three aforementioned transdiagnostic vulnerabilities were collected from a community-based sample of 3,094 9th graders in a large metropolitan area and analyzed using mixed effects modeling to evaluate mediation effects.

Results—Individually and when controlling for each other, all three transdiagnostic vulnerabilities mediated the relation between OCD and MDD symptoms both before and after adjusting for demographics. Conclusions: Distress tolerance, anxiety sensitivity, and anhedonia may be unique mechanisms accounting for comorbidity between OCD and MDD symptoms in youth. Longitudinal evaluation of these candidate transdiagnostic emotional vulnerabilities in adolescent OCD-MDD comorbidity is warranted.

Keywords

Obsessive-Compulsive Disorder; Depression; Anxiety; Anhedonia; Comorbidity

Recent perspectives in psychopathological science propose that cross-cutting dimensions and biological pathways underlie and account for the presentation and co-presentation of

*Corresponding Author: Greg Chasson, Ph.D. Illinois Institute of Technology, Department of Psychology, 3105 S. Dearborn, Pritzker Science Center, Chicago, IL 60616, gchasson@iit.edu, Phone: 312.567.5936, Fax: 312.567.3493.

Note: Work was performed at all author institutions. Authors report no conflict of interest or financial disclosures for this project.

various mental disorders (Cuthbert & Insel, 2013; Dozois, Seeds, & Collins, 2009; Sauer-Zavala et al., 2012). In one such conceptualization, the underlying cause of various emotional symptoms and disorders, as well as their comorbidity, may be underpinned by a smaller set of *reactive* transdiagnostic vulnerabilities (i.e., characteristic patterns of maladaptive responses to emotional stimuli and states) (e.g., Leventhal & Zvolensky, 2015). By enhancing or diminishing the response to emotion beyond normative functioning, reactive vulnerabilities can increase risk of emotional disturbances manifested in various ways (e.g., sadness, fear, withdrawal, disgust). By doing so, a small set of reactive vulnerabilities can precede and confer risk for various individual or combinations of clinical phenotypic expressions, including but not limited to MDD and OCD, and therefore be a root cause of psychopathological comorbidity.

A transdiagnostic emotional vulnerabilities framework may be useful for conceptualizing the high degree of overlap between obsessive-compulsive disorder (OCD) symptoms and major depressive disorder (MDD) symptoms (Watson, 2009). To this end, three putative vulnerabilities, highlighted as part of extant transdiagnostic emotion models (e.g., Leventhal & Zvolensky, 2015), are evaluated in the current study as mechanisms accounting for the overlap between OCD and MDD: distress tolerance (i.e., ability to withstand distressing states), anxiety sensitivity (i.e., fear of anxiety-related sensations), and anhedonia (i.e., inability to experience pleasure). Evidence from various clinical and non-clinical participant populations link poor distress tolerance with both OCD (Cougles et al. 2011; Cougles et al. 2012; Cougles et al. 2013; Macatee et al. 2013) and depressive (Ellis et al. 2013; McHugh et al. 2014; Ali et al. 2015; Buckner et al. 2007; Gorka et al. 2012, Magidson et al. 2013, Tull & Gratz 2013; Brandt et al. 2013, O’Cleirigh & Ironson 2007) symptoms. Much like distress tolerance, anxiety sensitivity has also been implicated in clinical and non-clinical samples with OCD (Calamari et al. 2008; Olatunji & Wolitzky-Taylor 2009; Raines et al. 2014; Storch et al. 2013; Wheaton et al. 2012), as well as depression in clinical and non-clinical samples (Otto et al. 1995; Rector et al. 2007, Taylor et al. 1996; Tull & Gratz 2008; Zavos et al. 2012). Moreover, anhedonia demonstrates a similar relationship with depression and OCD, as it is a diagnostic criterion for the former (APA 2013), and it was recently shown to positively correlated with OCD severity after controlling for depression in an online sample of adults with OCD (Abramovitch et al. 2014). Although evidence of the link between anhedonia and OCD is limited, anhedonia may represent an important vulnerability for OCD given the emerging evidence implicating the reward pathway in the pathophysiology of OCD (Abramovich et al., 2014), such as abnormal activation patterns in the ventral striatum and insula (e.g., Figeet et al., 2011).

Very little work has empirically tested whether reactive emotional vulnerabilities are transdiagnostic mechanisms of comorbidity (c.f., Wolitzky-Taylor et al., 2015; 2016; Zvolensky, Farris, Leventhal, & Schmidt, 2014), and no existing study has investigated anhedonia, distress tolerance, and anxiety sensitivity in tandem as mechanisms accounting for the relationship between OCD and MDD symptoms. To address this gap, the current cross-sectional correlational report in community-based sample of high school freshman, tested the hypothesis that all three transdiagnostic vulnerabilities (i.e., mediators) in isolation (i.e., each evaluated as a unique mediator in the absence of the other proposed mediators) and in combination (i.e., all mediators tested simultaneously as predictors in a single

mediation model) would account for a significant proportion of the relation between OCD symptoms and depressive symptoms. By testing these hypotheses, this study is positioned to provide evidence of possible mechanisms of co-occurrence of two debilitating psychiatric syndromes in a vulnerable developmental period (i.e., mid-adolescence), which could advance theoretical models of the etiology of this comorbidity and inform treatments for OCD and MDD phenomena.

Method

Participants and Procedures

For the purposes of this study, a community-based sample of 9th grade students was collected from ten public high schools across the greater Los Angeles metropolitan area as part of a larger longitudinal investigation of adolescent health and substance use (the current study uses responses from the first timepoint of data collection). These schools were chosen to sufficiently represent a diverse demographic: the percent of students who were eligible for free lunch within each school (i.e., students whose parental income $\geq$ 185% of the national poverty level) was 31.1% ($SD = 19.7$, range: 8.0% – 68.2%) among the ten schools. Students were ineligible if they were enrolled in special education (e.g., learning disabilities) or English as a Second Language Programs; a total of 4,100 students were deemed eligible to participate in the study. Assent to participate was collected from 3,874 (94.5%) students; of those, 3,383 (82.5%) provided written parental consent to participate in the study. After excluding students with substantial missing data on key variables from the analyses ($n = 289$), a total of 3,094 participants were used for the study at hand. For the sample, mean age was 14.57($SD = .40$), mostly female (54.3%), and diverse (e.g., 45.3% Hispanic, 16.5% Asian, 16% non-Hispanic White) (see Table 1 for more demographic details).

Paper-and-pencil surveys were distributed in Fall 2013 during two separate one-hour in-class survey sessions less than two weeks apart. Each school that participated was compensated \$2,500; students who completed the surveys were entered into a raffle for a gift card worth \$50 as a means of compensation. The study was approved by the University of Southern California Institutional Review Board.

Measures

Covariates and Sample Descriptive Measures—To describe the sample population and include potential covariates, we assessed for demographic characteristics (e.g., age, gender, race/ethnicity [coded as nominal variable to reflect the 8 categories listed in Table 1]). Median neighborhood income level was also computed utilizing self-reported zip code information based on 2010 U.S. Census data (United States Census Bureau/American FactFinder, 2013).

Anhedonia—The Snaith-Hamilton Pleasure Scale (SHAPS; Snaith et al., 1995) is a 14-item self-report measure in which participants rate the degree of agreement with statements regarding hedonic experience based on how they had been feeling in the past few days. Items use a 4-point Likert scale (rated 1-4) that are scored based on the algorithm from Snaith et al. (1995), in which item responses were dichotomized so that strongly agree and

agree equaled zero and strongly disagree and disagree equaled one. Higher scores reflecting more anhedonia. Prior research has found the SHAPS to exhibit sufficient internal consistency and construct validity in adolescent populations (Leventhal et al., 2015).

Anxiety Sensitivity—The Childhood Anxiety Sensitivity Index (CASI; Silverman et al., 1991) includes 18 items assessing fear or concern in response to anxiety-related physiological sensations, mental states, and social situations. Responses are based on a 3-point scale and are summed, with higher scores indicating higher anxiety sensitivity. The CASI has demonstrated adequate two-week test-retest reliability and satisfactory criterion-related validity in non-clinical adolescent samples (Weems et al., 2001; Silverman, Ginsburg, & Goedhart, 1999).

Distress Tolerance—The 15-item Distress Tolerance Scale (DTS; Simons & Gaher, 2005) includes statements regarding typical responses indicative of intolerance of distressing emotional states. The measure requires averaging the responses to each item, which each use a 5-point Likert scale, with higher average scores reflecting lower levels of distress tolerance. The DTS has been shown to have excellent psychometric properties, including high test-retest reliability and appropriate convergence with other relevant measures assessing affective distress and regulation (Simons & Gaher, 2005).

Obsessive Compulsive Disorder Symptoms—The OCD symptoms subscale of the Revised Children's Anxiety and Depression Scale (RCADS; Chorpita et al., 2000) was used to measure OCD symptoms. This subscale consists of 6 items based on a 4-point Likert scale (scored 0-3) indicating relative frequency of occurrence. Item results are added together, yielding a sum score, with higher scores indicating higher OCD symptom frequency. Prior work has demonstrated the RCADS to display good reliability and sufficient convergent and discriminant validity in adolescent samples (Ferdinand et al., 2006; Chorpita et al., 2000). A sum on the OCD subscale of greater than 5 considered in the clinical range, yielding a specificity of .65 and sensitivity of .70 for a diagnosis of OCD among adolescents (Chorpita, Moffitt, & Gray, 2005).

Depressive Symptoms—The Center for Epidemiologic Studies Depression Scale (CESD; Radloff, 1977) is a 20-item measure of depressive symptoms during the preceding week. Responses are based on a 4-point Likert scale and are summed, with higher numbers indicating higher frequency. The CESD has sufficient psychometric characteristics, and has been validated in samples of high school students as well as ethnically-diverse adolescent populations (Edman et al., 1999; Radloff, 1991). A score greater than 16 is considered in the clinical range (Radloff, 1977) with a score higher than 26 falling in the severe range.

Analytic Approach

Hypotheses were tested using mixed effects modeling (df 's based on Satterthwaite), with the first level of the model including all covariates (i.e., age, income, gender, and minority status) and mediator variables and the second level containing a school identity variable to account for participant nesting within schools.

We examined the direct effects (depressive symptoms regressed on OCD symptoms controlling for the mediators), indirect effects (association between OCD symptoms and depressive symptoms through the mediators), and total effects (depressive symptoms regressed on OCD symptoms not controlling for the mediators). We calculated the relative indirect effect (i.e., ratio of the indirect effect and total effect as an index of the proportion of the total effect accounted for by mediators; Preacher & Kelley, 2011) as an effect size. The mediational analyses were performed by first investigating each mediator in isolation from each other (Figure 1). We then calculated a multiple mediator model in which all three mediators were simultaneously included to control for overlapping variance in mediational paths and to determine which pathways may exhibit empirically unique relations relative to the others (Figure 2). All models were tested both unadjusted and after adjusting for demographic covariates. Results are reported as unstandardized path coefficients and relative indirect effects with bootstrapped 95% CIs in Tables 2–3, with standardized coefficients reported in Figures 1–2. Interpretation of the standardized path coefficients was based on Cohen's (1988) rule of thumb for small (.1), medium (.30), and large (.50).

Mediation analyses can be used for determining a sequential causal model, but the current study makes use of this approach as a tool for estimating the degree to which the transdiagnostic factors statistically account for the covariance between OCD and MDD symptoms (MacKinnon, Fairchild, & Fritz, 2007). There is precedent in the research literature for using mediation in this way to evaluate putative transdiagnostic factors (e.g. Wolitzky-Taylor et al. 2015; Wolitzky-Taylor et al., 2016). Missing data in the final sample were handled using multiple imputation. Inflated error rates were addressed by applying a False Discover Rate procedure— $\alpha_{\text{critical}} = .028$ (Benjamini & Hochberg, 1995).

Results

Preliminary Analysis and Total Effects

Descriptive statistics and Cronbach's alphas for the sample are provided in Table 1. Based on abovementioned cutoff scores, 33% of the sample scored in the clinical range for OCD symptom severity, and 33% and 16% in the clinical and severe ranges for MDD symptom severity, respectively. There were significant total effects for the positive relation of OCD and depressive symptoms in models unadjusted, $b(95\% \text{ bootstrapped CI}) = 1.39(1.29, 1.49)$, $p < .0001$, and adjusted by covariates, $b(95\% \text{ CI}) = 1.33(1.24, 1.43)$, $p < .0001$. Standardized path coefficients indicate a medium-to-large effect for OCD on depressive symptoms (see Figure note).

Emotional Vulnerabilities as Mediators

Greater OCD symptoms were associated with poorer distress tolerance, anxiety sensitivity, and anhedonic symptoms (Tables 2–3), with medium effect sizes for distress tolerance and anxiety sensitivity, and a small effect size for anhedonia (see Figure 1). Individual analyses of each mediator separate from one another ($a \times b$ paths) indicated that the OCD-MDD association was significantly mediated by each transdiagnostic mediator in the expected direction (see Tables 2–3). Distress tolerance and anxiety sensitivity demonstrated a similar

magnitude of effect size as indicated by the standardized indirect effect (see Figure 1) and relative indirect effect (see Tables 2 and 3).

Similarly, to test whether the mediational paths were empirically unique or redundant, the three mediators were entered into the same model in tandem. Similar to the models with a single mediator, when the mediators controlled for each other in the analyses, all three were significant, and the effect sizes for anxiety sensitivity and distress tolerance were similar in magnitude (see Tables 2 and 3 for full results). The standardized total indirect effect was small-to-medium in size (see Figure 2).

Supplemental Mediation Analyses Reversing the Ordering of MDD and OCD Symptoms

The transdiagnostic vulnerabilities framework applied here proposes that emotional vulnerabilities precede the onset of both MDD and OCD symptoms and that the ordering of OCD and MDD in the statistical modeling may be arbitrary. To address the possible bi-directionality of OCD and MDD symptoms, we tested the *alternate* ordering in which the abovementioned adjusted models were re-tested with OCD symptoms as the dependent variable and depressive symptoms as the independent variable. The pattern of findings paralleled those reported in the primary analysis—all three factors were significant mediators in the multiple-mediator models and the magnitudes of relative indirect effects were stronger for distress tolerance and anxiety sensitivity compared to anhedonia (results available upon request to the first author).

Discussion

In a community-based sample of adolescents, this study provides new evidence indicating that anhedonia, anxiety sensitivity, and distress tolerance each accounted for a significant portion of the relation between OCD and depressive symptoms (i.e., indirect effect was 55% and 50% of the total effect for adjusted and unadjusted analyses, respectively). The results were similar when statistical models included adjustments for demographics. Altogether, the evidence provides support that all three vulnerabilities may reflect transdiagnostic mechanisms of OCD-MDD comorbidity in adolescents. Given that each of these mediators remained robust after controlling for each other in the multiple mediation model, each of these putative vulnerabilities may represent unique causal pathways linking OCD and MDD symptoms.

The results of the current investigation align with transdiagnostic vulnerability models of emotional comorbidity (e.g., Leventhal & Zvolesnky, 2015; Fairholme, Boisseau, Ellard, Ehrenreich, & Barlow, 2010), which purport that OCD and MDD symptoms are not associated *merely* because OCD causes MDD or vice versa. Transdiagnostic models propose that in a significant portion of comorbid cases these vulnerabilities precede OCD and MDD symptoms and increase the risk of both disorders. The order onset of OCD relative to MDD may be arbitrary in cases whereby the comorbidity originates from a common transdiagnostic mechanism. Consistent with this model, when OCD and MDD symptoms were swapped as independent and dependent variables in a supplemental analysis, result patterns were comparable. In addition, the *b* path models from the original and swapped analyses suggest that these three emotional vulnerabilities independently associate with

depressive symptoms (while controlling for OCD symptoms), as well as OCD symptoms (while controlling for depressive symptoms). Given the cross-sectional and non-experimental nature of the study design, however, such alternate directional results provide only indirect support for the ordering of the vulnerabilities relative to MDD and OCD.

A transdiagnostic vulnerabilities model is not the only explanation for the current findings. The relation between OCD symptoms and depression remained after controlling for the mediators in the models, suggesting partial mediation. In addition, according to standardized path coefficients, most effects were small (e.g., anhedonia) or small-to-medium (e.g., total indirect effect). Thus, other factors may also explain the relation between OCD and MDD symptoms, in addition to the three transdiagnostic processes studied herein. Future studies should examine other factors that may account for the link between OCD and MDD, including psychological constructs (e.g., intolerance of uncertainty), biological factors (e.g., genetic variants in the serotonergic system), and environmental risk factors (e.g., trauma).

In contrast to a transdiagnostic vulnerabilities model, the causal cascade model whereby one disorder causes another through a reactive vulnerability mediating mechanism is another plausible account of why reactive vulnerabilities may account for OCD-MDD covariance. For example, individuals with OCD symptoms may experience high anxiety sensitivity (e.g., Calamari et al. 2008), triggering safety behaviors or avoidance (i.e., Mowrer, 1947) due to low distress tolerance (e.g., Cougle et al. 2011). In turn, these safety behaviors may cause the individual to withdrawal and therefore reduce positive reinforcement, eliciting high anhedonia (e.g., Hatzigiakoumis et al., 2011), which is a risk factor for depression (e.g., Lewinsohn, 1974). This is one plausible causal account of the comorbidity between OCD and MDD symptoms in addition to the transdiagnostic vulnerability model.

In addition to providing novel evidence of transdiagnostic processes, this study also adds to emerging literature on the role of reactive emotional vulnerabilities psychopathology that is primarily limited to adult samples. Results in this adolescent sample are consistent with and build on previous adult investigations linking OCD to distress tolerance (Cougle et al. 2011, Cougle et al. 2012, Cougle et al. 2013; Macatee et al. 2013; Robinson & Freeston 2014) and anxiety sensitivity (Calamari et al. 2008; Olatunji & Wolitzky-Taylor 2009; Raines et al. 2014; Storch et al. 2013; Wheaton et al. 2012), as well as studies associating depression with distress tolerance (e.g., Ellis et al. 2013; McHugh et al. 2014) and anxiety sensitivity (Otto et al. 1995; Rector et al. 2007, Taylor et al. 1996; Tull & Gratz 2008; Zavos et al. 2012). Our results advance the literature by documenting distress tolerance and anxiety sensitivity as possible explanation of the comorbidity between OCD and MDD symptoms in adolescents.

Only recently has evidence emerged for a link between anhedonia and OCD (Abramovitch et al., 2014). The current study extends this finding to a large, non-clinical community-based high school sample and suggests anhedonia may have a mediating role in the relation between OCD symptoms and depression that is empirically distinct from mediation by the other transdiagnostic factors. This finding is consistent with emerging research on the pathophysiology of OCD implicating brain reward pathways, such as the ventral striatum and nucleus accumbens (e.g., Figeet et al., 2011), that may underpin anhedonia (e.g., Nestler & Carlezon, 2006).

From a clinical perspective, the current results suggest that practitioners providing evidence-based psychosocial treatments for OCD and depression could benefit from targeting these transdiagnostic vulnerabilities. Specifically targeting low distress tolerance, anxiety sensitivity, and anhedonia is not a defined feature of exposure and response prevention, the gold standard psychosocial treatment for OCD (Jenike, 2004). Comorbid depressive symptoms reduce the efficacy of behavior treatments for OCD (Abramowitz & Foa, 2000; Foa et al.; 1983), and individuals with OCD and depressive symptoms tend to have higher levels of anxiety, more psychiatric comorbidities in general, increased rates of unemployment, and more functional disability (Abramowitz, Storch, Keeley, & Cordell, 2007; Hong et al. 2004; Ricciardi & McNally, 1995; Tükel, Meteris, Koyuncu, Tecer, & Yazici, 2006; Tükel, Polat, Özdemir, Skut, & Turksöy, 2002). Thus, incorporating such transdiagnostic vulnerabilities into a behavioral case conceptualization for OCD, particularly in the presence of comorbid depressive symptoms, may facilitate patient recovery.

One limitation of the study is the cross-sectional and correlational design, which limits causal inference about the mediating effects of the transdiagnostic vulnerability factors. This complicates drawing conclusions about temporal patterns, such as whether or not OCD or MDD are better modeled as predictors or outcomes. This cross-sectional limitation also precludes ruling out anhedonia as just a diagnostic feature of depression and not a vulnerability factor, although research and theory has supported the latter (Lewinsohn, 1974; Loas, 1996). As another limitation, the self-report nature of the data may suggest that a portion of the effects in the study stemmed from shared method variance. Similarly, there were no clinical diagnostic tools included in data collection to corroborate self-reports. The current study included a measure of anhedonia that measures feelings in a short and recent window of time (i.e., the last few days). Although the measure has demonstrated good temporal stability estimates (ICC $r = .70$; Franken et al., 2007), anhedonia as operationalized in this study may reflect more of a state construct than a trait construct. Future research would also benefit from enhanced generalizability by including a wider range of adolescent ages from other geographic regions, as well as a clinically diagnosed MDD and OCD sample. Studying this research question in a clinical sample, for example, could help account for other possible mechanisms linking these two conditions, such as medication usage and past treatments.

Conclusion

This is the first study to demonstrate jointly and independently the mediating effect of distress tolerance, anxiety sensitivity, or anhedonia in the relation between OCD symptoms and depression in a large, community-based adolescent sample. This is a group that is rarely studied and an age range that may be particularly pertinent for examining risk factors for the development of MDD and OCD, given that puberty is a common age at onset for MDD and most cases of OCD onset in adolescence or early adulthood (APA, 2013). Thus, future research corroborating results from the current study may suggest avenues for preventing MDD and OCD comorbidity by targeting important transdiagnostic emotional vulnerabilities.

Acknowledgments

This research was supported by funds from National Institute on Drug Abuse (grant number R01-DA033296). The funding source had no role in the study design, collection, analysis or interpretation of the data, writing the manuscript, or the decision to submit the paper for publication. None of the authors report a conflict of interest related to submission of this manuscript.

References

- Abramovitch A, Pizzagalli DA, Reuman L, Wilhelm S. Anhedonia in obsessive-compulsive disorder: Beyond comorbid depression. *Psychiatry Research*. 2014; 216:223–229. DOI: 10.1016/j.psychres.2014.02.002 [PubMed: 24564999]
- Abramowitz JS, Foa EB. Does major depressive disorder influence outcome of exposure and response prevention for OCD? *Behavior Therapy*. 2000; 31:795–800. DOI: 10.1016/S0005-7894(00)80045-3
- Abramowitz JS, Storch EA, Keeley M, Cordell E. Obsessive-compulsive disorder with comorbid major depression: What is the role of cognitive factors? *Behaviour Research and Therapy*. 2007; 45:2257–2267. DOI: 10.1016/j.brat.2007.04.003 [PubMed: 17521605]
- Ali B, Seitz-Brown CJ, Daughters SB. The interacting effect of depressive symptoms, gender, and distress tolerance on substance use problems among residential treatment-seeking substance users. *Drug and Alcohol Dependence*. 2015; 148:21–26. DOI: 10.1016/j.drugalcdep.2014.11.024 [PubMed: 25578252]
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 5th. 2013. Major Depressive Disorder.
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 5th. 2013. Obsessive Compulsive Disorder.
- Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society: Series B (Methodological)*. 1995; 57:289–300. DOI: 10.2307/2346101
- Brandt CP, Zvolensky MJ, Bonn-Miller MO. Distress tolerance, emotion dysregulation, and anxiety and depressive symptoms among HIV+ individuals. *Cognitive Therapy and Research*. 2013; 37:446–455. DOI: 10.1007/s10608-012-9497-9
- Buckner JD, Keough ME, Schmidt NB. Problematic alcohol and cannabis use among young adults: The roles of depression and discomfort and distress tolerance. *Addictive Behaviors*. 2007; 32:1957–1963. DOI: 10.1016/j.addbeh.2006.12.019 [PubMed: 17258398]
- Calamari JE, Rector NA, Woodard JL, Cohen RJ, Chik HM. Anxiety sensitivity and obsessive-compulsive disorder. *Assessment*. 2008; 15:351–363. DOI: 10.1177/1073191107312611 [PubMed: 18310595]
- Chorpita BF, Moffitt CE, Gray J. Psychometric properties of the Revised Child Anxiety and Depression Scale in a clinical sample. *Behaviour Research & Therapy*. 2005; 43:309–322. DOI: 10.1016/j.brat.2004.02.004 [PubMed: 15680928]
- Cohen, J. *Statistical power analysis for the behavioral sciences* (2nd edition). Hillsdale, NJ: Lawrence Erlbaum Associates; 1988.
- Chorpita BF, Yim L, Moffitt C, Umemoto LA, Francis SE. Assessment of symptoms of DSM-IV anxiety and depression in children: A revised child anxiety and depression scale. *Behaviour Research And Therapy*. 2000; 38:835–855. DOI: 10.1016/S0005-7967(99)00130-8 [PubMed: 10937431]
- Cougle JR, Timpano KR, Fitch KE, Hawkins KA. Distress tolerance and obsessions: An integrative analysis. *Depression and Anxiety*. 2011; 28:906–914. DOI: 10.1002/da.20846 [PubMed: 21681873]
- Cougle JR, Timpano KR, Goetz AR. Exploring the unique and interactive roles of distress tolerance and negative urgency in obsessions. *Personality and Individual Differences*. 2012; 52:515–520. DOI: 10.1016/j.paid.2011.11.017

- Cougle JR, Timpano KR, Sarawgi S, Smith CM, Fitch KE. A multi-modal investigation of the roles of distress tolerance and emotional reactivity in obsessive-compulsive symptoms. *Anxiety, Stress, & Coping*. 2013; 26:478–492. DOI: 10.1080/10615806.2012.697156
- Cuthbert, BN., Insel, TR. Toward precision medicine in psychiatry: The NIMH research domain criteria project. In: Charney, DS.Buxbaum, JD.Sklar, P.Nestler, EJ.Charney, DS.Buxbaum, JD., Nestler, EJ., editors. *Neurobiology of mental illness*. 4th. New York, NY, US: Oxford University Press; 2013. p. 1076-1088.
- Dozois DA, Seeds PM, Collins KA. Transdiagnostic approaches to the prevention of depression and anxiety. *Journal Of Cognitive Psychotherapy*. 2009; 23:44–59. DOI: 10.1891/0889-8391.23.1.44
- Edman JL, Danko GP, Andrade N, McArdle JJ, Foster J, Glipa J. Factor structure of the CES-D (Center for Epidemiologic Studies Depression Scale) among Filipino-American adolescents. *Social Psychiatry And Psychiatric Epidemiology*. 1999; 34:211–215. DOI: 10.1007/s001270050135 [PubMed: 10365627]
- Ellis AJ, Vanderlind WM, Beevers CG. Enhanced anger reactivity and reduced distress tolerance in major depressive disorder. *Cognitive Therapy and Research*. 2013; 37:498–509. DOI: 10.1007/s10608-012-9494-z
- Fairholme, CP., Boisseau, CL., Ellard, KK., Ehrenreich, JT., Barlow, DH. Emotions, emotion regulation, and psychological treatment: A unified perspective. In: Kring, AM.Sloan, DM.Kring, AM., Sloan, DM., editors. *Emotion regulation and psychopathology: A transdiagnostic approach to etiology and treatment*. New York, NY, US: Guilford Press; 2010. p. 283-309.
- Ferdinand RF, van Lang NJ, Ormel J, Verhulst FC. No distinctions between different types of anxiety symptoms in pre-adolescents from the general population. *Journal Of Anxiety Disorders*. 2006; 20:207–221. DOI: 10.1016/j.janxdis.2004.12.003 [PubMed: 16464705]
- Figeo M, Vink M, de Geus F, Vulink N, Veltman DJ, Westenberg H, Denys D. Dysfunctional reward circuitry in obsessive-compulsive disorder. *Biological Psychiatry*. 2011; 69(9):867–874. DOI: 10.1016/j.biopsych.2010.12.003 [PubMed: 21272861]
- Foa EB, Grayson JB, Steketee GS, Doppelt HG, Turner RM, Latimer PR. Success and failure in the behavioral treatment of obsessive-compulsives. *Journal Of Consulting And Clinical Psychology*. 1983; 51:287–297. DOI: 10.1037/0022-006X.51.2.287 [PubMed: 6841773]
- Franken IHA, Rassin E, Muris P. The assessment of anhedonia in clinical and non-clinical populations: Further validation of the Snaith–Hamilton Pleasure Scale (SHAPS). *Journal of Affective Disorders*. 2007; 99:83–89. DOI: 10.1016/j.jad.2006.08.020 [PubMed: 16996138]
- Gorka SM, Ali B, Daughters SB. The Role of Distress Tolerance in the Relationship Between Depressive Symptoms and Problematic Alcohol Use. *Psychology of Addictive Behaviors*. 2012; 26:621–626. DOI: 10.1037/a0026386 [PubMed: 22121919]
- Hatzigiakoumis DS, Martinotti G, Giannantonio MD, Janiri L. Anhedonia and Substance Dependence: Clinical Correlates and Treatment Options. *Frontiers in Psychiatry*. 2011; 2:10. <http://doi.org/10.3389/fpsy.2011.00010>. [PubMed: 21556280]
- Hong JP, Samuels J, Bienvenu OI, Cannistraro P, Grados M, Riddle MA, Nestadt G. Clinical correlates of recurrent major depression in obsessive-compulsive disorder. *Depression And Anxiety*. 2004; 20:86–91. DOI: 10.1002/da.20024 [PubMed: 15390212]
- Jenike MA. Obsessive-Compulsive Disorder. *The New England Journal Of Medicine*. 2004; 350:259–265. DOI: 10.1056/NEJMcp031002 [PubMed: 14724305]
- Leventhal AM, Unger JB, Audrain-McGovern J, Sussman S, Volk HE, Strong DR. Measuring anhedonia in adolescents: A psychometric analysis. *Journal Of Personality Assessment*. 2015; 97:506–514. DOI: 10.1080/00223891.2015.1029072 [PubMed: 25893676]
- Leventhal AM, Zvolensky MJ. Anxiety, depression, and cigarette smoking: A transdiagnostic vulnerability framework to understanding emotion–smoking comorbidity. *Psychological Bulletin*. 2015; 141:176–212. DOI: 10.1037/bul0000003 [PubMed: 25365764]
- Lewinsohn, PM. A behavioral approach to depression. In: Friedman, RJ.Katz, MM.Friedman, RJ., Katz, MM., editors. *The psychology of depression: Contemporary theory and research*. Oxford, England: John Wiley & Sons; 1974.
- Loas G. Vulnerability to depression: A model centered on anhedonia. *Journal Of Affective Disorders*. 1996; 41:39–53. DOI: 10.1016/0165-0327(96)00065-1 [PubMed: 8938204]

- Macatee RJ, Capron DW, Schmidt NB, Cogle JR. An examination of low distress tolerance and life stressors as factors underlying obsessions. *Journal of Psychiatric Research*. 2013; 47:1462–1468. DOI: 10.1016/j.jpsychires.2013.06.019 [PubMed: 23870797]
- MacKinnon DP, Fairchild AJ, Fritz MS. Mediation analysis. *Annual Review of Psychology*. 2007; 58:593–614. DOI: 10.1146/annurev.psych.58.110405.085542
- Magidson JF, Linstead AR, Seitz-Brown CJ, Anderson KE, Lindberg B, Wilson A, Daughters SB. Rumination mediates the relationship between distress tolerance and depressive symptoms among substance users. *Cognitive Therapy and Research*. 2013; 37:456–465. DOI: 10.1007/s10608-012-9488-x [PubMed: 23729916]
- McHugh RK, Kertz SJ, Weiss RB, Baskin-Sommers AR, Hearon BA, Björgvinsson T. Changes in distress intolerance and treatment outcome in a partial hospital setting. *Behavior Therapy*. 2014; 45:232–240. DOI: 10.1016/j.beth.2013.11.002 [PubMed: 24491198]
- Mowrer OH. On the dual nature of learning—a re-interpretation of 'conditioning' and 'problem-solving.'. *Harvard Educational Review*. 1947; 17:102–148.
- Nestler EJ, Carlezon WJ. The Mesolimbic Dopamine Reward Circuit in Depression. *Biological Psychiatry*. 2006; 59(12):1151–1159. DOI: 10.1016/j.biopsych.2005.09.018 [PubMed: 16566899]
- O'Leirigh CO, Ironson G. Does distress tolerance moderate the impact of major life events on psychosocial variables and behaviors important in the management of HIV? *Behavior Therapy*. 2007; 38:314–323. DOI: 10.1016/j.beth.2006.11.001 [PubMed: 17697855]
- Olatunji BO, Wolitzky-Taylor KB. Anxiety sensitivity and the anxiety disorders: A meta-analytic review and synthesis. *Psychological bulletin*. 2009; 135:974–999. DOI: 10.1037/a0017428 [PubMed: 19883144]
- Otto MW, Pollack MH, Fava M, Uccello R, Rosenbaum JF. Elevated anxiety sensitivity index scores in patients with major depression: Correlated and changes with antidepressant treatment. *Journal of Anxiety Disorders*. 1995; 9:117–123. DOI: 10.1016/0887-6185(94)00035-2
- Preacher KJ, Kelley K. Effect size measures for mediation models: Quantitative strategies for communicating indirect effects. *Psychological Methods*. 2011; 16:93–115. DOI: 10.1037/a0022658 [PubMed: 21500915]
- Raines AM, Capron DW, Bontempo AC, Dane BF, Schmidt NB. Obsessive compulsive symptom dimensions and suicide: The moderating role of anxiety sensitivity cognitive concerns. *Cognitive Therapy and Research*. 2014; 38:660–669. DOI: 10.1007/s10608-014-9622-z
- Radloff LS. The CES-D Scale: A self-report depression scale for research in the general population. *Applied Psychological Measurement*. 1977; 1:385–401. DOI: 10.1177/014662167700100306
- Radloff LS. The use of the Center for Epidemiologic Studies Depression Scale in adolescents and young adults. *Journal Of Youth And Adolescence*. 1991; 20:149–166. DOI: 10.1007/BF01537606 [PubMed: 24265004]
- Rector NA, Szacun-Shimizu K, Leybman M. Anxiety sensitivity within the anxiety disorders: disorder-specific sensitivities and depression comorbidity. *Behaviour Research and Therapy*. 2007; 25:1967–1975. DOI: 10.1016/j.brat.2006.09.017
- Ricciardi JN, McNally RJ. Depressed mood is related to obsessions, but not to compulsions, in obsessive-compulsive disorder. *Journal Of Anxiety Disorders*. 1995; 9:249–256. DOI: 10.1016/0887-6185(95)00006-A
- Robinson LJ, Freeston MH. Emotion and internal experience in obsessive compulsive disorder: Reviewing the role of alexithymia, anxiety sensitivity, and distress tolerance. *Clinical Psychology Review*. 2014; 34:256–271. DOI: 10.1016/j.cpr.2014.03.003 [PubMed: 24682109]
- Sauer-Zavala S, Boswell JF, Gallagher MW, Bentley KH, Ametaj A, Barlow DH. The role of negative affectivity and negative reactivity to emotions in predicting outcomes in the unified protocol for the transdiagnostic treatment of emotional disorders. *Behaviour Research And Therapy*. 2012; 50:551–557. DOI: 10.1016/j.brat.2012.05.005 [PubMed: 22738907]
- Silverman WK, Fleisig W, Rabian B, Peterson RA. Child Anxiety Sensitivity Index. *Journal Of Clinical Child Psychology*. 1991; 20:162–168. DOI: 10.1207/s15374424jccp2002_7
- Silverman WK, Ginsburg GS, Goedhart AW. Factor structure of the Childhood Anxiety Sensitivity Index. *Behaviour Research And Therapy*. 1999; 37:903–917. DOI: 10.1016/S0005-7967(98)00189-2 [PubMed: 10458052]

- Simons JS, Gaher RM. The Distress Tolerance Scale: Development and validation of a self-report measure. *Motivation And Emotion*. 2005; 29:83–102. DOI: 10.1007/s11031-005-7955-3
- Snaith RP, Hamilton M, Morley S, Humayan A, Hargreaves D, Trigwell P. A scale for the assessment of the hedonic tone: The Snaith-Hamilton Pleasure Scale. *The British Journal Of Psychiatry*. 1995; 167:99–103. DOI: 10.1192/bjp.167.1.99 [PubMed: 7551619]
- Storch EA, Wu MS, Small BJ, Crawford EA, Lewin AB, Horng B, Murphy TK. Mediators and moderators of functional impairment in adults with obsessive– compulsive disorder. *Comprehensive Psychiatry*. 2013; doi: 10.1016/j.comppsy.2013.10.014
- Taylor S, Koch WJ, Woody S, McLean P. Anxiety sensitivity and depression: How are they related. *Journal of Abnormal Psychology*. 1996; 105:474–479. DOI: 10.1037/0021-843X.105.3.474 [PubMed: 8772020]
- Tükel R, Meteris H, Koyuncu A, Tecer A, Yazici O. The clinical impact of mood disorder comorbidity on obsessive-compulsive disorder. *European Archives Of Psychiatry And Clinical Neuroscience*. 2006; 256:240–245. DOI: 10.1007/s00406-006-0632-z [PubMed: 16683062]
- Tükel R, Polat A, Özdemir Ö, Aksüt D, Türksoy N. Comorbid conditions in obsessive-compulsive disorder. *Comprehensive Psychiatry*. 2002; 43:204–209. DOI: 10.1053/comp.2002.32355 [PubMed: 11994838]
- Tull MT, Gratz KL. Further examination of the relationship between anxiety sensitivity and depression: The mediating role of experiential avoidance and difficulties engaging in goal-directed behavior when distressed. *Anxiety Disorders*. 2008; 22:199–210. DOI: 10.1016/j.janxdis.2007.03.005
- Tull MT, Gratz KL. Major depression and risky sexual behavior among substance dependent patients: The moderating roles of distress tolerance and gender. *Cognitive Therapy and Research*. 2013; 37:483–497. DOI: 10.1007/s10608-012-9490-3 [PubMed: 24634552]
- United States Census Bureau / American FactFinder. 2010. Retrieved 1 January 2013, from <http://factfinder2.census.gov> 0030-1299.2005.13727.x
- Watson D. Differentiating the mood and anxiety disorders: A quadripartite model. *Annual Review of Clinical Psychology*. 2009; 5:221–247. DOI: 10.1146/annurev.clinpsy.032408.153510
- Weems CF, Berman SL, Silverman WK, Saavedra LM. Cognitive errors in youth with anxiety disorders: The linkages between negative cognitive errors and anxious symptoms. *Cognitive Therapy And Research*. 2001; 25:559–575. DOI: 10.1023/A:1005505531527
- Wheaton MG, Mahaffey B, Timpano KR, Berman NC, Abramowitz JS. The relationship between anxiety sensitivity and obsessive-compulsive symptom dimensions. *Journal of Behavior Therapy and Experimental Psychiatry*. 2012; 43:891–896. DOI: 10.1016/j.jbtep.2012.01.001 [PubMed: 22321579]
- Wolitzky-Taylor K, Guillot CR, Pang RD, Kirkpatrick MG, Zvolensky MJ, Buckner JD, Leventhal AM. Examination of anxiety sensitivity and distress tolerance as transdiagnostic mechanisms linking multiple anxiety pathologies to alcohol use problems in adolescents. *Alcoholism: Clinical And Experimental Research*. 2015; 39(3):532–539. DOI: 10.1111/acer.12638
- Wolitzky-Taylor K, McBeth J, Guillot CR, Stone MD, Kirkpatrick MG, Zvolensky MJ, Leventhal AM. Transdiagnostic processes linking anxiety symptoms and substance use problems among adolescents. *Journal Of Addictive Diseases*. 2016; 35(4):266–277. DOI: 10.1080/10550887.2016.1207969
- Zavos HMS, Rijdsdijk FV, Eley TC. A longitudinal, genetically informative, study of associations between anxiety sensitivity, anxiety and depression. *Behavior Genetics*. 2012; 42doi: 10.1007/s10519-012-9535-0
- Zvolensky MJ, Farris SG, Leventhal AM, Schmidt NB. Anxiety sensitivity mediates relations between emotional disorders and smoking. *Psychology Of Addictive Behaviors*. 2014; 28(3):912–920. DOI: 10.1037/a003745 [PubMed: 25222178]

Figure 1. Standardized Path Coefficients for Individual Mediation Models

Note. Separate mediational model sets were conducted for each transdiagnostic mediator. a = path from the independent variable (i.e., OCD symptoms) to the mediator. b = path from the mediator (i.e., distress tolerance, anxiety sensitivity, anhedonia) to the dependent variable (i.e., depression) after controlling for the effect of the independent variable. $a \times b$ = the indirect effect of the independent variable on the dependent variable that occurs through the mediator, which equals the product of the “a” path and “b” path. c' = the direct effect of the independent variable on the dependent variable that is not carried through the mediator, which equals the effect of the independent variable on the dependent variable after controlling for the mediator. The total effect = the sum of the indirect effect ($a \times b$) and direct effect (c'), which equals the effect of the independent variable on the dependent variable not adjusting for the mediators. Unadj = unadjusted $B(95\% \text{ CI})$; Adj = adjusted $B(95\% \text{ CI})$. Unadjusted C path β ($95\% \text{ CI}$) = .45 (.42, .48). Adjusted C path β ($95\% \text{ CI}$) = .43 (.40, .46).

Figure 2. Standardized Path Coefficients for Combined Mediation Models

Note. A single mediational model set was conducted, which included all mediators simultaneously modeled. *a* = path from the independent variable (i.e., OCD symptoms) to the mediator. *b* = path from the mediators (i.e., distress tolerance, anxiety sensitivity, anhedonia entered as simultaneous predictors) to the dependent variable (i.e., depression) after controlling for the effect of the independent variable. $a \times b$ = the indirect effect of the independent variable on the dependent variable that occurs through a mediator controlling for the mediation occurring through the other two variables, which equals the product of the “*a*” path and “*b*” path for that specific mediator. c' = the direct effect of the independent variable on the dependent variable that is not carried through the mediators, which equals the effect of the independent variable on the dependent variable after controlling for all three mediators. The total effect = the sum of the indirect effect ($a \times b$) of the three mediators and the direct effect (c'), which equals the effect of the independent variable on the dependent variable. For the “*a*” path estimates, please see Figure 1. Unadj = unadjusted *B*(95% CI); Adj = adjusted *B*(95% CI). Unadjusted C path β (95% CI) = .45 (.42, .48). Adjusted C path β (95% CI) = .43 (.40, .46).

Table 1

Sample Characteristics and Variable Descriptive Statistics

Variable: <i>M(SD)</i> or %	Overall Sample (<i>N</i> = 3,094)	α
Age	14.57 (.40), range 12.83–16.24]	
Gender		
Female	54.3	
Male	45.7	
Race/Ethnicity^a		
American Indian or Alaskan Native	1.0	
Asian	16.5	
Black or African American	4.8	
Native Hawaiian or Pacific Islander	3.4	
Hispanic or Latino	45.3	
White	16.0	
Multiracial	5.6	
Other	5.7	
Median Income by Zipcode	\$69,469 (\$13,433)	
Symptoms		
RCADS OCD Subscale	4.48 (3.86) [range 0–18]	.82
CESD	14.66 (11.92) [range 0–60]	.81
DTS	2.58 (0.85), [range 1–5]	.91
CASI	30.48 (7.04), [range 18–54]	.88
SHAPS	1.71 (2.40), [range 0–14]	.89

Note.

^aMissing race/ethnicity data for 1.7% of participants. RCADS OCD Subscale = Revised Children's Anxiety Scale Obsessive-Compulsive Disorder Subscale. Center for Epidemiologic Studies Depression Scale. DTS = Distress Tolerance Scale. CASI = Childhood Anxiety Sensitivity Index. SHAPS = Snaith-Hamilton Pleasure Scale. Data from 9th grade students in Los Angeles, California, USA collected in Fall 2013. α = Cronbach's α internal consistent estimate.

Unadjusted Effects of OCD Symptom Severity on Depressive Symptom Severity and Mediation by Distress Tolerance, Anxiety Sensitivity, and Anhedonia

Table 2

	Component Paths			Mediation: OCD symptoms → Mediator(s) → MDD symptoms		
	OCD symptoms → Mediator(s) Controlling for OCD symptoms	Mediator(s) → MDD symptoms Controlling for OCD symptoms		Indirect effect AB Path (95% CI)	Direct effect C' Path (95% CI)	Relative Indirect Effect (95% CI)
Individual Analyses						
Distress Tolerance	.10 (.10, .11) [‡]	3.95 (3.46, 4.44) [‡]		.40 (.36, .43) [‡]	.98 (.87, 1.09) [‡]	.287 (.25, .34)
Anxiety Sensitivity	.87 (.81, .93) [‡]	.47 (.41, .53) [‡]		.41 (.35, .47) [‡]	.98 (.87, 1.09) [‡]	.294 (.25, .35)
Anhedonia	.06 (.04, .08) [‡]	1.54 (1.40, 1.69) [‡]		.11 (.06, .13) [‡]	1.30 (1.20, 1.39) [‡]	.066 (.04, .10)
Combined Analyses						
Distress Tolerance	–	2.66(2.16, 3.15) [‡]		.27(.21, .32) [‡]	–	.194 (.16, .24)
Anxiety Sensitivity	–	.38 (.32, .44) [‡]		.33 (.28, .39) [‡]	–	.237 (.20, .29)
Anhedonia	–	1.60(1.46, 1.74) [‡]		.10 (.06, .13) [‡]	–	.072 (.04, .10)

Note. $N = 3,094$. Data from 9th grade students in Los Angeles, California, USA collected in fall 2013. OCD = Obsessive Compulsive Disorder symptom severity. MDD = depressive symptom severity. A path = unstandardized path coefficient from the respective mediator regressed on OCD symptoms. B Path = unstandardized path coefficient from depression severity regressed on the mediator(s). C' Path = unstandardized path coefficient for OCD symptoms for the model in which depression severity was simultaneously regressed on the mediator(s) and OCD symptoms (i.e., relation of OC to depression after controlling for mediator(s)). C Path = unstandardized path coefficient from depression severity regressed on OCD symptoms—total effect, C-path (95% CI) = 1.39 (1.29, 1.49)[‡]. Relative indirect effect is an effect size commonly interpreted as the proportion of the total effect accounted for by mediation (Preacher & Kelley, 2011).

[‡] $p < .0001$.

Adjusted Effects of OCD Symptom Severity on Depressive Symptom Severity and Mediation by Distress Tolerance, Anxiety Sensitivity, and Anhedonia

Table 3

	Component Paths			Mediation: OCD symptoms → Mediator(s) → MDD symptoms		
	OCD symptoms → Mediator(s) Controlling for OCD symptoms			Direct effect		
	A Path (95% CI)	B Path (95% CI)	AB Path (95% CI)	C' Path (95% CI)	Relative Indirect Effect (95% CI)	
Individual Analyses						
Distress Tolerance	.10 (.09, .11) [†]	3.49 (3.01, 3.98) [†]	.35 (.30, .41) [†]	.98 (.88, 1.09) [†]	.262 (.23, .32)	
Anxiety Sensitivity	.84 (.78, .89) [†]	.38 (.32, .44) [†]	.32 (.26, .37) [†]	1.02 (.91, 1.12) [†]	.240 (.20, .30)	
Anhedonia	.06 (.04, .09) [†]	1.60 (1.46, 1.74) [†]	.10 (.06, .13) [†]	1.24 (1.14, 1.33) [†]	.072 (.04, .11)	
Combined Analyses						
Distress Tolerance	–	2.54(2.06, 3.02) [†]	.25 (.20, .31) [†]	–	.191 (.16, .24)	
Anxiety Sensitivity	–	.29(.23, .35) [†]	.24 (.19, .30) [†]	–	.183(.14, .24)	
Anhedonia	–	1.62(1.49, 1.76) [†]	.10 (.06, .13) [†]	–	.073 (.05, .11)	

Note. $N = 3,094$. Data from 9th grade students in Los Angeles, California, USA collected in fall 2013. OCD = Obsessive Compulsive Disorder symptom severity. MDD = depressive symptom severity. A path = unstandardized path coefficient from the respective mediator regressed on OCD symptoms plus covariates. B Path = unstandardized path coefficient from depression severity regressed on the mediator(s) plus covariates. C' Path = unstandardized path coefficient for OCD symptoms for the model in which depression severity was simultaneously regressed on the mediator(s), OCD symptoms (i.e., relation of OC to depression after controlling for mediator(s)) plus covariates. C Path = unstandardized path coefficient from depression severity regressed on OCD symptoms plus covariates —total effect; C-path (95% CI) = 1.33 (1.24, 1.43)[†]. Covariates for the models include gender, minority status (non-Hispanic White vs. not non-Hispanic White), age, and income. Relative indirect effect is an effect size commonly interpreted as the proportion of the total effect accounted for by mediation (Preacher & Kelley, 2011).

[†] $p < .0001$.