Published in final edited form as:

Dev Psychopathol. 2014 November ; 26(4 Pt 2): 1241–1254. doi:10.1017/S0954579414000996.

Sensitizing effect of early adversity on depressive reactions to later proximal stress: Moderation by 5-HTTLPR and CRHR1 in a 20-year longitudinal study

Lisa R. Starr,

University of Rochester

Constance Hammen,

University of California, Los Angeles

Christopher C. Conway,

University of California, Los Angeles

Elizabeth Raposa, and

University of California, Los Angeles

Patricia A. Brennan

Emory University

Abstract

Previous research supports gene-environment interactions for polymorphisms in corticotrophin hormone receptor (CRHR1) and the serotonin transporter (5-HTTLPR) genes in predicting depression, but has rarely considered genetic influences on stress sensitization processes, whereby early adversities (EA) increase depressive reactivity to proximal stressors later in life. The current study tested a gene-environment-environment interaction ($G \times E \times E$; specifically gene-EA-proximal stress interaction) model of depression in a 20-year longitudinal study. Participants were assessed prospectively for EA up to age 5 and recent chronic stress and depressive symptoms at age 20 and genotyped for CRHR1 SNP rs110402 and 5-HTTLPR. EA predicted stronger associations between recent chronic stress and depression, and the effect was moderated by genes. CRHR1 A alleles and 5-HTTLPR short alleles were associated with greater stress sensitization (i.e., greater depressive reactivity to chronic stress for those also exposed to high levels of EA). Results are consistent with the notion that EA exposure results in neurobiological and cognitive-emotional consequences (e.g., altered hypothalamic-pituitary-adrenal [HPA] axis functioning) leading to emotional distress in the face of recent stressors among those with certain genetic characteristics, although further research is needed to explore explanatory mechanisms.

Keywords

serotonin transporter gene; 5-HTTLPR; corticotrophin hormone receptor gene (CRHR1); depression; early adversity; stress

Environmental circumstances play an indisputable role in the pathogenesis of depression, including both as distal predictors of long-term risk and short-term precipitants of depressive episodes. However, environmental events do not uniformly influence depression risk, and thus much of contemporary research in the psychopathology of depression has focused on neurobiological and psychosocial mechanisms by which stress and adversity phenomena result in depressive experiences in some people but not others. The numerous, interacting, and mutually-influencing factors that likely contribute to depressive vulnerability span diverse systems—including biological (genetic, neural, and neuroendocrine), psychological (personality traits, cognitions, coping styles), developmental (timing of stressor), and environmental (characteristics and contexts of the stressors)—and also involve the interplay between multiple risk and protective factors.

Research using a multiple-levels-of-analysis framework may help elucidate how such varied systems collectively influence depression risk. The study of multilevel dynamics involves identifying the transactions among biological, environmental, and psychosocial processes influencing stage-salient development, leading to diverse outcomes (Masten, 2007). Biological and environmental risk factors are not viewed as solitary forces operating in isolation, but instead as interactive pieces in intricate systems that cut across multiple spheres to influence outcomes. To capture such complexity, progressively more integrative models and methodologies are needed (e.g., Cicchetti, 2013; Hammen, Rudolph, & Abaied, in press; Masten, 2007), particularly those incorporating long-term longitudinal designs that shed light on how experiences occurring across crucial developmental periods impact distal outcomes. As one of the founding principles of developmental psychopathology, the multiple levels perspective has transformed research on risk and protective factors for psychopathology across the life span (Cicchetti & Dawson, 2002; Cicchetti & Toth, 2009).

The present study examines risk factors for depression across multiple levels of analysis in a sample of youth at risk for depression. We examine interactive effects of multiple systems, including genetic and environmental, consistent with an abundance of research supporting gene-environment interactions ($G \times E$) in predicting depression (e.g., Karg, Burmeister, Shedden, & Sen, 2011). Moreover, multiple levels of analysis are further construed to include multiple levels of environmental factors. Rather than view “stress” as a unitary construct, we address contributions of both early childhood adverse conditions and chronic stress more proximal to depressive experiences and their potential interactive relationship. Specifically, we examine whether early adverse experiences during developmentally salient early childhood periods affect depressive reactivity to proximal stressors later in life (an environment-environment interaction [$E \times E$]), and whether this “stress sensitization” pattern in turn is moderated by genetic vulnerability (gene-environment-environment interaction [$G \times E \times E$]). We focus on two candidate genes relevant to sensitivity to stress—a polymorphism in the promoter region of the serotonin transporter gene (5-HTTLPR), and a single nucleotide polymorphism (SNP; rs110402) in the corticotrophin releasing hormone receptor 1 gene (CRHR1).

Early Adversity and Depression

Exposure to adverse conditions in early childhood is strongly predictive of adolescent and adult depressive disorders, as well as anxiety, substance use, and disruptive behavioral disorders, in the U.S. (Kessler, Davis, & Kendler, 1997; Kessler & Magee, 1993; McLaughlin, Green, Gruber, Sampson, Zaslavsky, & Kessler, 2012) and internationally (Kessler et al., 2010). Adversities reflecting family functioning may be especially strongly associated with internalizing disorders (McLaughlin et al., 2012).

Exposure to early adversities may increase risk for depression in part by modifying reactivity to proximal stressors. The *stress sensitization* hypothesis, or the idea that prior childhood adversity exposure predicts greater likelihood of reacting to recent stressors with depression, has been supported in several studies. Hammen, Henry, and Daley (2000) showed that among young women who developed a depressive episode in a longitudinal study of the transition to adulthood, those reporting childhood adversities developed a depressive episode following lower levels of recent stressors than women without childhood adversity exposure. Harkness, Bruce, and Lumley (2006) also showed that adolescents with a history of childhood maltreatment reported a lower severity of stressful life events prior to first depressive episode onset than reported by those without abuse. Similarly, in an epidemiological sample of adults McLaughlin, Conron, Koenen, and Gilman (2010) found that past-year stressful life events were more predictive of major depression among those with higher levels of early childhood adversity than among those without adversity.

Genetic Predictors of Sensitivity to Stress

The current study examined genetic moderation of stress sensitization. There are several plausible reasons to expect that some individuals would be genetically predisposed to become more reactive to stressors following early adversity exposure. First, the mechanisms presumed to underlie stress sensitization may be influenced by genetic factors. Although both biological and psychosocial processes likely contribute to effects of early adversity exposure on depression and stress sensitization (Cicchetti, 2013), one important mechanism may be disruptions in the development of neuroregulatory processes. For example, hypothalamic-pituitary-adrenal (HPA) axis functioning has been shown to be sensitive to early adversity exposure (e.g., Cicchetti & Rogosch, 2001; Gunnar & Quevedo, 2007). Exposure to stressful conditions (e.g., family conflict, harsh discipline, maltreatment) in childhood is linked to abnormalities in basal levels of cortisol (Gonzalez, Jenkins, Steiner, & Fleming, 2009; Taylor, Karlamangla, Friedman, & Seeman, 2011) and cortisol reactivity to acute stressors (Wisner Fries, Shirtcliff, & Pollak, 2008), which are in turn consistent with abnormalities in the HPA axis observed in depressed individuals (Pariante & Lightman, 2008; Posener et al., 2000). Thus, disruptions in the HPA axis development, as well as their effects on neural circuitry, may promote sensitivity to stressful life events and result in increased risk for later depression. As such, researchers have suggested that genes that regulate HPA axis functioning may influence the degree to which biological systems are altered by early stress exposure (Gillespie, Phifer, Bradley, & Ressler, 2009).

In addition, as outlined below, recent $G \times E$ research has identified specific candidate genes that interact with environmental stressors (including early childhood adversity and recent acute and chronic stress) to predict depressive outcomes. The current study extends these findings by examining whether genes interact with early adversity exposure to predict depressive reactivity to proximal stressors in a $G \times E \times E$ pattern. We focused on two genes that both play an established role in HPA axis functioning and have been shown to interact with stress to predict depression.

Corticotrophin Releasing Hormone Receptor Gene (CRHR1)

Given the central role of corticotrophin releasing hormone in HPA axis functioning (Heim & Nemeroff, 2001), polymorphisms in CRHR1 are considered promising candidates for $G \times E$ research in depression. Several studies have examined a three-allele CRHR1 haplotype (or the individual variants that constitute it) involving SNPs at rs7209436, rs110402, and rs242924 as a possible moderator of adversity-depression associations. The first $G \times E$ study on this topic found that a TAT haplotype at these loci attenuated the association between retrospectively-reported child abuse—assessed using the Childhood Trauma Questionnaire (CTQ; Bernstein & Fink, 1998)—and adult depressive symptoms (Bradley et al., 2008). A second investigation replicated the TAT haplotype buffering effect in one sample in which child abuse was again measured with the CTQ, but not in another sample that assessed abuse with a combination of prospective and retrospective indices (Polanczyk et al., 2009). In other words, in these two studies the A allele rs110402—the main focus of the present analyses—(in combination with T alleles at the other two loci) was associated with reduced vulnerability to depression following stress exposure. In contrast, other studies have observed that the TAT haplotype intensifies the association of childhood maltreatment with trait neuroticism and blunted diurnal cortisol rhythms, two phenotypes known to underlie risk for depressive disorders (Cicchetti, Rogosch, & Oshri, 2011; DeYoung, Cicchetti, & Rogosch, 2011). In line with these latter findings, a recent study demonstrated $G \times E$ effects where the TAT haplotype was associated with elevated risk for alcoholism (Ray et al., 2013), and several studies have directly linked the TAT haplotype to internalizing outcomes, including depression (Papiol et al., 2007; Ishitobi et al., 2012; Wasserman, Wasserman, & Sokolowski, 2010). Additional investigations have failed to find evidence for $G \times E$ involving any of these SNPs and childhood adversity (Lewis, Collishaw, Harold, Rice, & Thapar, 2012), or concluded that the pattern of interaction varies for males versus females (Heim et al., 2009). No studies to date have examined the role of CRHR1 variation in predicting the association of proximal stressors to depression, though one investigation demonstrated greater risk of suicidal responses to stress among rs110402 A allele carriers (Wasserman, Sokolowski, Rozanov, & Wasserman, 2008). It is evident that CRHR1 $G \times E$ findings to date have been inconsistent, and there is some debate over whether interaction findings are dependent on the measure used to define childhood stressors (i.e., CTQ; Heim et al., 2009; Lewis et al., 2012), the nature of the stressors (e.g., severity of early adversity, sexual versus physical abuse; DeYoung et al., 2011; Heim et al., 2009) or characteristics of the sample such as ethnicity. Together, these studies suggest that CRHR1 variants play a role in modulating depressive responses to stress, but more research is needed to discern the exact pattern of risk.

Serotonin Transporter Gene

In addition to CRHR1 genotypes thought to moderate the effects of childhood adversity exposure on depression outcomes, another theoretically plausible candidate gene is a length polymorphism in the promoter region of the serotonin transporter gene (5-HTTLPR). Caspi and colleagues (2003) initially provided support for a $G \times E$ effect for 5-HTTLPR in depression, demonstrating that short (S) allele carriers show heightened depression rates following stress exposure compared to long (L) allele homozygotes. The decade following the publication of Caspi et al.'s (2003) findings has seen a tremendous growth of research on this gene in the context of depression. Although there have been several notable failures to replicate Caspi et al.'s (2003) $G \times E$ pattern (see Risch et al., 2009), the majority of longitudinal studies find that S allele carriers experience higher rates of depression following exposure to naturally-occurring stressors (Caspi, Hariri, Holmes, Uher, & Moffitt, 2010; Uher & McGuffin, 2010). A recent quantitative review demonstrated that 5-HTTLPR $G \times E$ is a fairly robust phenomenon when only studies that assess stressful environmental conditions with gold-standard interview methods or objective criteria (e.g., legally documented cases of abuse) are included (Karg et al., 2011). Additionally, there is evidence that 5-HTTLPR interacts more strongly with early childhood adversity, as compared to stressful experiences that are more proximal to depression onset (Karg et al., 2011), perhaps due to the chronicity of biological and psychological dysfunction set into motion by early stress exposure (Brown & Harris, 2008).

Further, a growing body of experimental research demonstrates that 5-HTTLPR plays a role in regulating neural, endocrine, and attentional reactivity to stressful conditions (e.g., Munafò, Brown, & Hariri, 2008; Pergamin-Hight, Bakermans-Kranenburg, van IJzendoorn, & Bar-Haim, 2012). For instance, several independent teams of investigators have observed that 5-HTTLPR S allele carriers, relative to L allele homozygotes, exhibit heightened cortisol responses to social stressors in the laboratory (Gotlib, Joormann, Minor, & Hallmayer, 2008; Way & Taylor, 2010). All told, existing evidence makes a compelling case that 5-HTTLPR genotype plays a crucial role in stress regulation.

The notion that the two genes, CRHR1 and 5-HTTLPR, may influence depressive sensitivity to both early adversities and later proximal stressors in a $G \times E \times E$ relationship has been proposed (e.g., Homberg & van den Hove, 2012) but rarely tested. As previously noted, research on CRHR1's role in depression has exclusively examined early adversities and not proximal stressors. The larger, more extensive research literature on 5-HTTLPR has separately examined the interactive effects of both early and proximal stressors with 5-HTTLPR genotype, but has rarely differentiated between their effects. In one recent exception, Grabe et al. (2012) tested a three-way interaction between 5-HTTLPR genotype, child abuse, and adult trauma exposure. They found support for the $G \times E \times E$ effect, with S allele carriers exposed to both child abuse and adult traumas showing the highest levels of depression. These results offer compelling preliminary support for our model; however, the Grabe et al. (2012) study was limited by its cross-sectional design, use of a retrospective, self-report measure of child abuse (i.e., CTQ), and exclusive focus on adult trauma rather than more normative proximal stressors. Thus, our study seeks to both replicate and extend these findings.

Timing of Early Adversity

Several researchers have proposed the existence of sensitive periods of neural development, during which relevant brain circuitry is maximally plastic and therefore stressful conditions have the greatest potential to disrupt development of stress regulation systems (see Heim & Binder, 2012). Although research has yet to pinpoint the age at which this sensitive period occurs, some evidence suggests that stressors occurring at younger ages, such as preschool age and earlier, may have stronger negative effects on biological markers of stress regulation than those occurring at later ages (e.g., Carpenter et al., 2004, 2009; Heim, Newport, Mlezko, Miller, & Nemeroff, 2008; Rogeness & McClure, 1996). Stress exposure during this early time period may also disrupt important socioemotional development tasks, such as attachment formation (Bowlby, 1980), which have lasting implications for stress regulation and mental health (e.g., van IJzendoorn, Schuengel, & Bakermans-Kranenburg, 1999). As such, some evidence suggests that adversities experienced in early childhood may be particularly predictive of mental health outcomes (Dunn, McLaughlin, Slopen, Rosand, & Smoller, 2013; Keiley, Howe, Dodge, Bates, & Pettit, 2001; Gunnar & van Dulmen, 2007). In line with these findings, in the current study, we define early adversities as major stressors occurring during the first five years of life.

The Present Study

The present study examines the effect of early childhood adversity exposure on depressive responses to stressful conditions in young adulthood, and whether the effects are further modified by genotypes of a SNP of the CRHR1 gene (rs110402), and by a polymorphism in the promoter region of the serotonin transporter gene (5-HTTLPR). We hypothesize $G \times E \times E$ interactions involving each of the two candidate genes, with greater early adversity exposure increasing depressive responses to chronic stress at age 20, depending on genetic vulnerability.

Chronic, ongoing stress, rather than acute life events, will be the focus for empirical and conceptual reasons: the $G \times E$ literature seems to support a stronger association of depression with chronic conditions such as maltreatment than acute recent events (Brown & Harris, 2008; Karg et al., 2011; Moffitt, Caspi, & Rutter, 2005), and ongoing stress presents a more stable and reliable picture of stress experience at a particular time point (e.g., Moffitt et al., 2005; Rutter, 2005). The data are from a longitudinal sample of youth at risk for depression, studied since mothers' pregnancy and selected from a large cohort to represent children of depressed or never-depressed women. Depressive outcomes were assessed at age 20. The transition to adulthood is a developmentally active period marked by selection into relationship, social, occupational and academic roles that are increasingly challenging and potentially stressful, and for some, a period of risk for depressive and anxiety disorders and other forms of maladaptation. We assessed both self-reported depressive symptoms and also a broader index of internalizing pathology to test the likelihood that results apply not only to depressive experiences but more generally to the common depression-anxiety mix. Childhood adversity exposure is a multiple-variable scale capturing commonly correlated risk factors affecting the family, such as parental marital instability, parental mental illness or criminality, poverty, and high levels of stressful life events. The measure is based on

contemporaneous data collected from the mothers at 4 time points from pregnancy to child age 5. Measures of youth chronic stress at age 20 included reliable and valid interview assessments across multiple domains, and depressive symptoms were collected at youth age 20. Several previous studies examining CRHR1 and 5-HTTLPR have revealed gender-specific effects (Heim et al., 2009; Du, Bakish, & Hrdina, 2000; Hammen, Brennan, Keenan-Miller, Hazel, & Najman, 2010) and internalizing problems are substantially more common among females (Kessler, McGonagle, Swartz, Blazer, & Nelson, 1993); therefore, gender is a covariate in analyses of the $G \times E \times E$ hypotheses.

Method

Participants

Participants were drawn from an initial sample of 815 youth who completed procedures at age 15 for a study of youth at risk due to maternal depression, of whom 705 took part in a follow-up study at age 20. Between ages 22 to 25, all youth who could be located were asked to participate in collection of DNA samples, of whom 512 consented. All individuals included in the present analyses had to have been part of the original data collection through age 5, the ages 15 and 20 follow-ups, and the DNA collection. The current study is based on 381 (149 males and 232 females) who were genotyped for 5-HTTLPR, and 444 (182 males, 262 females) who were genotyped for the rs110402 SNP of CRHR1. Characteristics and data collection procedures are described in the Genotyping section below.

The original sample of 815 was recruited from the Mater University Study of Pregnancy (MUSP; Keeping et al., 1989) consisting of all women who were pregnant and gave birth at the Mater Misericordiae Mother's Hospital in Brisbane, Queensland, Australia in the early 1980s. The original birth cohort study of youth and mother health and development consisted of over 7,000 families, and included measures of mothers' circumstances and depressive symptoms during pregnancy, immediately post-partum, at six months after birth, and 5 years after birth. Mothers' self-reported depressive symptoms on the Delusions-Symptoms-States Inventory (DSSI; Bedford & Foulds, 1978) were used by investigators (CH and PB) to identify women with varying levels of severity and chronicity of depressive symptoms—or no depression—for a study of children of 815 women. Full details of the original sample selection and ascertainment are reported in Hammen and Brennan (2001). Mothers' depression status was confirmed by the Structured Clinical Interview for DSM-IV (SCID; First, Spitzer, Gibbon, & Williams, 1995) when the children turned 15 years of age; 357 (44%) had diagnoses of major depressive disorder and/or dysthymic disorder in the child's lifetime; 458 (56%) were never depressed. The youth sample at age 15 consisted of 403 females and 412 males. The families were predominately lower and middle income, primarily Caucasian (91.4%; 3.6% Asian, 5% "other" or not reported).

Those participants in the age 20 follow-up who were not included in the DNA sampling encompassed those who could not be contacted, no longer lived in the geographical area, declined to participate, had significant medical problems, or were deceased. There were no differences between those who participated and those who did not on depression history or maternal depressive status, but participants were more likely to be female ($\chi^2 = 21.29, p < .001$).

Measures

Youth depressive and internalizing symptoms—The Beck Depression Inventory-II (BDI; Beck, Steer, & Brown, 1996) is a widely used measure of depressive symptoms with excellent psychometric properties, including reliability and validity, as well as strong sensitivity and specificity for detecting depression in community samples (Beck, Steer, & Garbin, 1988; Lasa, Ayuso-Mateos, Vazquez-Barquero, Diez-Manrique, & Dowrick, 2000). Cronbach's alpha was .92 at the age 20 assessment. The BDI was employed instead of depression diagnoses because (a) relatively few participants met full criteria for current major depressive episode (MDE) at age 20 ($n = 17$, 3.8% of the sample), (b) BDI is a valid measure of risk for diagnosable depression (Beck et al., 1988) and is also, in the current sample, predictive of current and future impairment in functioning, and (c) the BDI provides continuous symptom data, which allow greater statistical power for detecting interaction effects. The broadband internalizing scores included the internalizing subscales of the Young Adult Self-Report. These scales are part of the Achenbach System of Empirically Based Assessments (ASEBA; Achenbach, 2009). The YSR internalizing scale is based on the sum of the scores of the Withdrawn, Somatic Complaints, and Anxious/Depressed scales, and the YASR included the sum of Withdrawn and Anxious/Depressed subscales. The scales have well-established psychometric evidence of reliability and validity (e.g., Achenbach, 2009).

Early adversity (EA)—A contemporaneous composite measure of family adversity in the first five years of the child's life was derived from information provided by the mother at one or more of the pregnancy, birth, 6-month and 5-year assessments (except for maternal Axis I diagnosis). An EA composite was formed from procedures previously reported in Hazel, Hammen, Brennan, and Najman (2008). Variables included as adversities were maternal Axis I diagnosis prior to age 5 years, financial hardship, child chronic illness, parental discord, maternal stressful life events and mothers' separation from partners. *Maternal Axis I diagnoses* (omitting specific phobia) between the child's birth and age 5 years were assessed using the Structured Clinical Interview for DSM-IV (First et al., 1995) at the age 15 years interview, covering the mother's lifetime. The most common diagnoses were major depressive disorder, dysthymic disorder, and social phobia. Kappa for depressive disorders within the child's first 5 years in a reliability sample was 0.81 ($p < .01$). *Financial hardship* was assessed by calculating the mean of maternal ratings of total family income at the prenatal, 6-month and 5-year data collections. *Childhood illness* was assessed at the age 5 years assessment by asking mothers to endorse whether the child had had any of 15 illnesses or injuries (e.g. asthma) lasting 3 or more months that impaired the child's activities at least 'some'. *Maternal stressful life events* were assessed using checklists of nine interpersonal, health or occupational problems that might have occurred in the last 6 months prior to the prenatal and post-natal data collections. The numbers of reported events at the two assessments were highly correlated ($r = .59$) and were summed to reduce the checklists to a single measure of perinatal stressful life events. *Parental discord* was evaluated as maternal relationship satisfaction with her romantic partner, assessed using the mean of the eight-item satisfaction scale of the Dyadic Adjustment Scale (DAS; Spanier, 1976) collected during pregnancy and at birth, 6 months and 5 years (α 's ranged from 0.85 to 0.97). If on the age 5 years questionnaire a mother reported that she had been divorced or separated from a partner or that she had changed partners over the last 5 years the child was

considered to have experienced a *parental partner separation*. The continuous variables, income, DAS and maternal life events, were recoded as present/absent using the 33rd percentile as the cut-off point for each measure to identify the third of the sample experiencing the most adverse conditions. The specific cut-off was chosen as a consistent point across measures that would balance the need for sufficient numbers for meaningful analyses with selection of a moderately adverse level of each variable. A summary measure of early childhood adversity was formed by counting the number of adversities for each child, resulting in a range of 0–6 adversities experienced (distribution: zero adversities = 23%, one = 28%, two = 18%, three = 16%, four = 8%, five or more = 7%).

Youth chronic stress—The UCLA Life Stress Interview (e.g., Hammen & Brennan, 2001) was used to obtain a summary measure of chronic stress at the age 20 interview, covering the past 6 months. It is a semi-structured interview for ongoing conditions adapted for younger and older adolescents, and was developed from earlier versions for adults (e.g., Hammen et al., 1987). The age 20 version included developmentally appropriate domains: social life, close friendship, romantic relation (dating activity and interest), relations with family members, financial, work, academic activity, health of self, health of close family members. Each scale included parallel versions to capture all participants' status; for example, the academic scale included versions for those continuing further education and for those not pursuing additional education. Trained advanced graduate student interviewers probed each area with the youth, using standard general probes and semi-structured follow-up queries where needed. Each domain was scored by the interviewer on a 5-point scale with behaviorally anchored descriptors to capture objective, factual conditions. Across all domains, the anchors at the "1" end of the scale represented superior (exceptional) functioning, and "5" represented severe difficulties. In order to make the ratings as objective as possible, the interviewer elicited specific information and examples, and each scale value was indicated by specific behavioral information. For example, the "social life" segment queried the number of people in the youth's regular social circle, how close and trusting their relationships are, what kind and frequency of activities do they engage in, how much they are sought out by others, frequency of disputes and conflicts, and how they are resolved, leading to scoring exemplified by levels of the following scale of "social life":

1-Exceptional social life –many good friends, very popular and engages in frequent social activities outside school, gets along well with others, no conflict.

3-Average social life but has some conflicts with peers or difficulty making and keeping friends.

5-Severe social problems with no friends, totally isolated from peers or frequent conflicts and fights, rejected by peers.

Reliabilities were based on independent judges' ratings of audiotaped interviews. Mean intraclass correlation across the domains was .81 at age 20. Evidence supporting the convergent and predictive validity of the chronic stress measure in the current sample is reported in Hammen, Brennan, & Keenan-Miller (2008). For the current study, summary measures for chronic stress levels were computed as a sum across all stress domains.

Genotyping

Two to five years after the age 20 follow-up (mean interval of 3.32 years, $SD = 1.02$) participants provided DNA samples for genetic analysis and completed questionnaires. Participants were mailed blood collection kits and consent forms, and had blood samples drawn at local facilities, which were picked up by courier and delivered to the Genetic Epidemiological Laboratory of the Queensland Institute of Medical Research (QIMR). The Institutional Review Boards of the University of Queensland, UCLA, and Emory University approved this research.

5-HTTLPR—Because of budgetary and procedural constraints, genotyping was restricted to a single full plating of 384 samples; 384 participants were selected randomly for genotyping from the pool of participants who provided blood-based DNA samples. Three samples produced invalid readings, leading to a final sample of 381 participants. Although selection for genotyping was random, by chance more female participants were selected than males ($\chi^2 = 16.49$, $p < .001$); otherwise, the genotyped and nongenotyped participants did not differ by their own depression status at previous follow-ups or by maternal depression status.

Genotyping was conducted at QIMR using agarose gel analysis of polymerase chain reaction products spanning the central portion of the repeats in the 5-HTTLPR. Polymerase chain reaction utilized Qiagen enzyme and buffer, with 30% deazaguanine and with 10 cycles of Touchdown protocol beginning at 67 °C and finishing at 62 °C with a further 32 cycles. Samples were subject to independent duplicate polymerase chain reaction with primer set 1 (acgttgatgTCCTG CATCC CCCAT, cgttgatgGCAGGGGGGATACTGCGA, lower case sequence is nontemplated) that gave products of 198 and 154 bp for Long and Short versions, respectively, and primer set 2 (acgttgatgTCCTGCATCC CCCAT, acgttgatgGGGGATGCTG GAAGGGC) for products of 127 and 83 bp. Gel analyses were conducted in triplicate for most samples. At least two matching independent results were required for inclusion. Final call rate was 96.4%. To estimate accuracy, duplicate samples were genotyped for 764 individuals in a different study in the same laboratory, following above procedures, with discordance rates of 0.45%. In the current sample, frequency of genotypes was L/L = 122 (32%), L/S = 178 (47%), and S/S = 81 (21%), with proportions in Hardy-Weinberg equilibrium, $\chi^2(1, 381) = 1.61$, $p < .20$. Based on evidence suggesting that the L form variants designated as L_g function similarly to the S allele (Wendland, Martin, Kruse, Lesch, & Murphy, 2006), 21 L_g variants were reclassified as s forms. Following this reclassification, updated allele frequencies were L/L = 101 (27%), S/L = 189 (50%), S/S = 91 (24%).

CRHR1—Aliquots of blood-based DNA were shipped to UCLA for processing at the Social Genomics Core of the USC/UCLA Biodemography Center. The *CRHR1* rs110402 and rs242924 polymorphisms were genotyped because of their frequent use in previous $G \times E$ studies; they were highly correlated with each other ($r = .99$), and constitute two-thirds (along with rs7209436) of the TAT haplotype commonly cited in the literature. The rs110402 and rs242924 polymorphisms were assayed by a commercial TaqMan Genotyping Assay (Applied Biosystems, Foster City, CA) performed on an iCycler real-time PCR instrument (BioRad, Hercules, CA) following the manufacturer's specified protocol, as

described in Cole et al. (2010). Test-retest reliability of duplicated specimens yielded a total genotyping error rate < 1%. Because of the near identity of the polymorphisms, primary analyses were conducted on the rs110402 snp. Genotype distributions were in Hardy-Weinberg equilibrium, $\chi^2 = 1.28$ ($p > .05$), and were distributed as follows: G/G, $n = 140$ (31%); A/G, $n = 208$ (47%); and A/A, $n = 96$ (22%).

Results

Descriptive Statistics and Main Effects

Table 1 displays descriptive statistics for study variables for the full sample and divided by genotype. Table 2 lists bivariate correlations for study variables. Neither 5-HTTLPR nor CRHR1 genotype was significantly associated with depressive symptoms at age 20, EA, chronic stress at age 20, gender, or maternal depression status, all $ps > .05$. Age 20 BDI showed significant, positive bivariate correlations with both EA and recent chronic stressors, which were also correlated with each other, all $ps < .001$. Both EA and chronic stress variables remained significant when entered simultaneously into a regression predicting age 20 depressive symptoms (chronic stress $b = 0.86$, $SE = 0.07$, $p < .001$; EA $b = 0.47$, $SE = 0.20$, $p = .020$).

Interaction of Early Adversity and Chronic Stress in Predicting Depressive Symptoms

All subsequent analyses controlled for gender. We first examined whether experiencing adversities in early childhood increased susceptibility to proximal chronic stress, using multiple linear regression with age 20 BDI as the outcome. Both predictor variables were mean-centered. We entered main effects for EA and chronic stress in the first step, and their interaction in the second step. The interaction term was significant ($b = 0.010$, $SE = 0.04$, $p = .014$, $R^2 = .25$, R^2 due to interaction = .01), with stronger associations between chronic stress and BDI at high levels of EA (one SD above the mean; $b = 0.97$, $SE = 0.08$, $p < .001$) than at low levels (one SD below the mean; $b = 0.67$, $SE = 0.10$, $p < .001$).

Interactive Effects between CRHR1 Genotype, Early Adversity, and Proximal Chronic Stress

Using the same statistical approach described above, we next separately tested two-way interactions between CRHR1 rs110402 and (a) adversity by age 5, and (b) proximal chronic stress, predicting age 20 depressive symptoms. Genotype was coded as a continuous scale reflecting number of A alleles (G/G = 0, A/G = 1, A/A = 2, an approach used elsewhere; e.g., Bradley et al., 2008), but applying a dichotomous coding scheme (presence versus absence of A allele) yielded similar results. Number of A alleles significantly interacted with EA ($b = 0.95$, $SE = 0.37$, $p = .011$, $R^2 = .07$, $R^2 = .02$), with no effect of EA on depressive symptoms for G homozygotes ($b = -.043$, $SE = 0.50$, $\beta = -.08$, $p = .395$) but significant effects for A carriers (with similar effects for A/G heterozygotes, $b = 1.86$, $SE = 0.42$, $\beta = .32$, and A homozygotes, $b = 1.42$, $SE = 0.48$, $\beta = .32$, $ps < .001$). CRHR1 genotype also interacted with recent chronic stress to predict age 20 depressive symptoms ($b = 0.29$, $SE = 0.12$, $p = .016$, $R^2 = .25$, $R^2 = .01$). Chronic stress significantly predicted greater depressive symptoms across all genotypes (all $ps < .001$), but at a significantly higher magnitude for A

carriers (A/G $b = 1.11$, $SE = 0.14$, $\beta = .52$; A/A $b = 1.07$, $SE = 0.18$, $\beta = .57$) than for G homozygotes ($b = 0.62$, $SE = 0.14$, $\beta = .36$).

Results should be interpreted in light of the final step, a three-way interaction (CRHR1 genotype $\times$ EA $\times$ Chronic Stress) predicting depression. The three-way interaction term was significant ($b = 0.24$, $SE = 0.07$, $p = .001$, $R^2 = .29$, $R^2 = .02$), and is illustrated in Figure 1. Conditional effects were evaluated using the PROCESS macro for SPSS (Hayes, 2013). Among A homozygotes, EA potentiated the effect of recent stressors on depressive symptoms ($b = 0.35$, $SE = 0.10$, $p < .001$). In contrast, there was no interaction between EA and chronic stress for G homozygotes ($b = -0.13$, $SE = 0.08$, $p = .11$). In the alternative comparison, the G $\times$ E between CRHR1 and chronic stress was significant only for those with higher levels of childhood adversities.

Substituting broadband YASR internalizing symptoms for depressive symptoms as the dependent variable produced comparable results, with a significant three-way interaction between CRHR1 genotype, chronic stress, and EA ($b = 0.21$, $SE = 0.09$, $p = .014$, $R^2 = .18$, $R^2 = .01$) predicting age 20 internalizing symptoms, showing a similar pattern of decomposition. Analyses were repeated using the highly correlated ($r = .99$) alternate CRHR1 SNP rs242924. Results were virtually identical, with no substantive differences in significance or magnitude.

Interactive Effects between 5-HTTLPR Genotype, Early Adversity, and Proximal Chronic Stress

We next separately tested two-way interactions of 5-HTTLPR genotype with (a) EA and (b) proximal chronic stress in predicting age 20 depressive symptoms. Genotype was defined as the number of S alleles (L/L = 0, S/L = 1, S/S = 2), consistent with the approach of others (e.g., Caspi et al., 2003), but coding genotype dichotomously (0 = L/L, 1 = S/L or S/S) produced similar results. Number of S alleles did not interact with EA to predict depressive symptoms ($p = .767$) but did interact with chronic stress to predict age 20 depressive symptoms ($b = 0.35$, $SE = 0.14$, $p = .011$, $R^2 = .26$, $R^2 = .01$). In line with expectations, associations between chronic stress and depressive symptoms were stronger for S-carriers (S/L: $b = 1.07$, $SE = .013$, $\beta = .54$; S/S: $b = 1.20$, $SE = 0.24$, $\beta = .49$, $p < .001$) than for L homozygotes ($b = 0.56$, $SE = 0.17$, $\beta = .33$, $p = .001$).

Finally, we tested a three-way interaction between 5-HTTLPR genotype, EA, and recent chronic stress, predicting age 20 depressive symptoms. In a hierarchical linear regression, we first entered gender as a covariate, then all main effects, then all two-way interactions, then the three-way interaction term. The three-way interaction coefficient was significant ($b = 0.22$, $SE = 0.08$, $p = .009$, $R^2 = .28$, $R^2 = .01$), as illustrated in Figure 2. The conditional effects of the chronic stress by EA interaction at different short allele counts were evaluated using the PROCESS macros for SPSS (Hayes, 2013). This two-way interaction was significant for S-homozygotes ($b = 0.29$, $SE = 0.11$, $p < .010$), with stronger effects of chronic stress on depressive symptoms for those with high EA (90th percentile; $b = 1.90$, $SE = 0.29$, $p < .001$) than for those with low adversity (10th percentile; $b = 0.74$, $SE = 0.30$, $p = .013$). The interaction between chronic stress and EA was not significant for L homozygotes ($b = -0.14$, $SE = 0.09$, $p = .112$). Stated otherwise, the G $\times$ E between 5-HTTLPR and recent

chronic stress was moderated by history of early adverse experiences: it was significant for those with a history of high early adversities ($b = 0.79$, $SE = 0.21$, $p < .001$), with significant reactivity to chronic stressors increasing with the number of S alleles, whereas there was no significant G×E for those with low adversities ($b = -0.08$, $SE = 0.22$, $p = .720$).

The three-way interaction between EA, age 20 chronic stress, and 5-HTTLPR genotype was also significant in predicting age 20 broadband internalizing symptoms ($b = 0.19$, $SE = 0.09$, $p = .032$, $R^2 = .18$, $R^2 = .01$), and the pattern of interaction was equivalent to that described above.

All genetic data analyses were repeated controlling for race (Caucasian versus non-Caucasian), with no impact on significance of results. Excluding non-Caucasians from analyses also did not alter results.

Discussion

The current study found support for the hypothesis that specific genetic polymorphisms predict greater depression reactivity to proximal stress among those with high levels of exposure to adverse conditions in early childhood. The multiple levels formulation of $G \times E \times E$ is consistent with a model of the impact of EA exposure on shaping developing neural, biological, and psychosocial processes toward greater likelihood of emotional reactivity to later stressors (sensitization), guided by genetic vulnerability. The results extend prior $G \times E$ findings on 5-HTTLPR and CRHR1, suggesting that some of the inconsistencies in prior findings with these genotypes may be due, at least in part, to unknown and unmeasured variation in exposure to distal or proximal stress.

With a few exceptions, previous $G \times E$ studies focusing on these genotypes have examined only the impact of stressors occurring during a single developmental period. Assessing multiple levels of stressors at different time points, the present study specifically examined sensitization processes, in which early childhood stress exposure predicted the strength of depressive reactivity in the face of current stressors, and the extent to which they were intensified by genetic factors. To date, no studies of CRHR1 $G \times E$ have examined naturally occurring proximal stressors on depression, although their typical focus on EA exposure may imply the assumption of an underlying sensitization reaction to later stress which has not generally been measured or reported. The larger research literature on 5-HTTLPR and stress has variously included recent *or* distal stressors including maltreatment in childhood (or both, as in Caspi et al., 2003; see description of multiple studies in Karg et al., 2011). However, we are aware of only one study that has specifically included the sensitization hypothesis of an interaction between early and later stressors. Grabe et al. (2012) found support for the hypothesis that 5-HTTLPR genotype moderated the interaction between early childhood maltreatment and adult traumas in the prediction of depressive outcomes. The current study both replicated these findings using several superior methodological elements (long-term longitudinal design, prospective assessment of adversities and stressors, gold-standard interview measure of proximal stress) and extended this conceptualization to a second genotype implicated in stress reactivity and depression etiology.

Our findings add an additional layer of complexity to traditional $G \times E$ models, but the complete nature of the relationship between genetic vulnerability, environmental risk, and depression is likely far more intricate still. First, multiple environmental risk factors tend to correlate with each other. Different types of early adversities tend to cluster together (e.g., Kessler, Davis, & Kendler, 1997), and the dysfunctional family dynamics accompanying various forms of adverse childhood experiences also promote vulnerability through cognitive and socioemotional pathways not tested here (e.g., Cicchetti, 2013). Similarly, there tends to be continuity between adversities early in life and stress exposure at later developmental stages; in previous research we showed that children at risk due to maternal depression tended to have high rates of continuing acute and chronic stress and depression over a 20-year follow-up (Hammen, Hazel, Brennan, & Najman, 2012). Thus, the experiences of early childhood and those of later adolescence do not occur within a vacuum, and disentangling their individual influences on depression in interaction with genetic factors can be difficult.

Secondly, genetic vulnerability is not independent of environmental risk, as previous research has supported a likely role of gene-environment correlations. Many of the ingredients of early adversity—exposure to maladaptive parenting, maltreatment, family violence, marital instability, and parental mental illness, among others, have heritable components (e.g., Kendler & Baker, 2006). Children inherit genes and also, in a different manner, environments, affecting not only what they are exposed to but different coping and resource characteristics. One marker of genetic risk for depression—having a depressed parent—also represents an environmental risk factor (e.g., Gotlib & Colich, in press) and is often accompanied by the multiple early adversities (marital instability, stress, economic disadvantage and child ill health) included in the current study. Genetic vulnerability for depressive reactivity to stress—specifically 5-HTTLPR genotype—also predicts a tendency to generate or select into stressful contexts following depression (Starr, Hammen, Brennan, & Najman, 2012). Thus, children at high genetic risk for depression experience more, generate more, and may be more biologically sensitive to stress, a cascade of forces conjointly and interactively promoting depressive vulnerability.

Indeed, the findings may be particularly relevant to populations at risk for depression due to maternal depression. Mothers' depressive experiences in the early lives of their children are fairly common, and are often accompanied by the multiple correlated risk factors of comorbidity, marital instability, stress, economic disadvantage and child ill health included in the current study. We may tend to think of the children of depressed mothers as having a predisposed vulnerability to depression, but another perspective is to view them as having a vulnerability to negative emotional reactivity to stress based on genetically influenced traits and heightened exposure to environmental risk factors, which then increases their risk for depression and other internalizing pathology.

Finally, the individual genotypes investigated here likely operate in conjunction with a variety of other genes. Indeed, they may interact with each other, as a recent study supported a $G \times G \times E$ effect between early maltreatment, CRHR1, and 5-HTTLPR in predicting depressive symptoms (Ressler et al., 2010). We did not test interactive effects of both genes in the current study, as testing a four-way interaction ($G \times G \times E \times E$) would not have been feasible in light of our moderate sample size, but it is plausible that such an effect could

exist. It is safe to assume that many other genes also contribute, with possible candidates including FKBP5 (Gillespie et al., 2009), COMT (Mandelli et al., 2007), and BDNF (Comasco, Aslund, Orelund, & Nilsson, 2013). In sum, in addition to $G \times E$ and $G \times E \times E$ interactions, environmental risk aggregates with other environmental risk, genetic risk interacts with other genetic risk, and genetic risk directly influences environmental risk. Consequently, many additional genetic and psychosocial factors are undoubtedly in play well beyond the few specific elements modeled in the current study. Although modeling numerous additive and multiplicative effects can be a challenge, strategies of multiple levels of analysis are essential.

There are several issues that require further study and clarification. First, our results suggested that $G \times E \times E$ effects are not only limited to depression, but also apply to more broadly defined internalizing symptoms. Future research should more closely evaluate specificity and explore other non-internalizing outcomes. Second, the effects of EA timing on stress regulation systems need to be resolved, particularly in relation to $G \times E$ (or more complicated forms of gene-environment interplay) effects. Research suggests that sensitive periods may occur, during which neural systems are most susceptible to environmental influences, but has yet to precisely identify the age frame in which this occurs (Heim & Binder, 2012). Indeed, doing so is methodologically complicated, as it is unusual for stress exposure to occur in a single isolated time period (Hammen et al., 2012). Further, there may not be a single, discrete sensitive period; different brain structures mature at varying rates and may be differentially affected by stressors occurring at different ages (Andersen et al., 2008) and there may be multiple periods of plasticity (Heim & Binder, 2012). Moreover, EA may also influence stress sensitization through other mechanisms, such as disruption of psychosocial developmental tasks, which may also occur at different ages. Although our decision to limit our definition of EA to events in the first five years of life is supported by available research (e.g., Carpenter et al., 2004; Dunn et al., 2013; Heim et al., 2008), further research is needed to better understand when environmental events have their most potent effects for those with genetic vulnerabilities.

Third, the literature remains inconsistent over the risk versus protective properties of the different alleles of this CRHR1 SNP. In our study, the A allele at rs110402 (and the T allele of rs242924) conferred greater risk for depression in interaction with environmental stressors, conflicting with some previous studies describing the A allele as protective (Bradley et al., 2008; Polanczyk et al. [E-Risk]; 2009), but in line with other studies that also suggest that the A allele elevates risk, both directly (Ishitobi et al., 2012; Papiol et al., 2007; Wasserman et al., 2009), and in interaction with negative environmental conditions (Cicchetti et al., 2011; DeYoung et al., 2011; Ray et al., 2013). Variation in findings may be related to how stress was defined in each study, reinforcing the notion that environmental stressors are complex phenotypes whose specific properties may differentially moderate genetic risk. For example, some research suggests that the A allele protects only against effects of physical and/or severe abuse (Heim et al., 2009), but amplifies the effects of other kinds of forms of maltreatment (DeYoung et al., 2011). In contrast to earlier studies, our EA composite measure did not explicitly include maltreatment, but instead assessed aspects of the early family environment that contribute to a stressful milieu. Current findings suggest that the A allele exacerbates the effects of these adversities. Furthermore, studies supporting

the protective properties of the A allele have exclusively relied on a single retrospective, self-report measure of maltreatment (CTQ; Bernstein, 1998; e.g., Bradley et al., 2008; Polanczyk et al., 2009 [E-Risk study]; Heim et al., 2009; Hsu et al., 2012). Retrospective self-reports may have been biased by current depressive state, and may not provide precision about timing and duration of occurrence.

Far more research is needed to understand the functional properties of the CRHR1 gene and its association with depression, and we hope that future researchers will bear certain points in mind. First, given the small size of the literature on this gene, it is likely premature to conclusively describe either allele as “risk” or “protective.” Further, as others have noted (Belsky & Pluess, 2009), the concept of risk alleles may be fundamentally oversimplified, as many genotypes may be maladaptive in some circumstances and beneficial in others. Finally, if it is indeed the case that the precise qualitative nature of the environment may influence the $G \times E$ effect, it underscores the importance of using methodologically rigorous measures of environmental stressors, ideally collected longitudinally. As others have noted (Monroe & Reid, 2008), the proliferation of stress assessments with relatively weak psychometric properties has contributed to inconsistent results with the $G \times E$ literature (although even studies with high-quality stress measures can yield inconsistent results; Polanczyk et al., 2009).

In addition, considerable further study of the mechanisms accounting for depressive reactivity to stress is needed to clarify the enormously complex psychobiological processes of development in the first decades of life. We acknowledge the considerable complexity of biological processes linking genetic, neural, and neuroendocrine pathways between EA exposure and internalizing symptomatology. An important next step will be to elucidate specific biological mechanisms driving the $G \times E \times E$ effects revealed in this study. We hypothesize (as have others; e.g., Gillespie et al., 2009) that these genes are associated with susceptibility of HPA axis development to environmental influences during a critical period of development, and that this in turn produces lasting neurological changes that elevate reactivity to further stressors at a later developmental stage. Although our results are consistent with this hypothesis, they do not directly test it. Future research should examine $G \times E \times E$ effects with biological outcomes that more directly correspond to potential biological mechanisms, including markers of HPA axis reactivity, inflammation, and limbic activation in response to stress.

The present study has several implications for the development of empirically supported preventions and interventions. First, it adds to existing support for the premise that early stress contributes to greater vulnerability to later stress. A reasonable corollary of this finding is that interruption of high stress exposure might lead to more adaptive reactions to later stressful experiences; thus, programs aimed at reducing adverse conditions during critical periods of development may have lasting preventative effects. Second, genetic factors may act as biomarkers that identify children at added risk due to early adversity exposure, offering the prospect of targeting limited resources to children who are most vulnerable to dysfunctional outcomes. Empirically-supported treatments for children exposed to adverse conditions are growing in number and appear to improve outcomes in a variety of areas, including indicators of HPA-axis functioning (e.g., Fisher, Gunnar,

Chamberlain, & Reid, 2000; Toth, Rogosch, Manly, & Cicchetti, 2006; for reviews of empirically supported programs see Institute of Medicine and National Research Council, 2013; National Research Council, 2009), although there are a host of challenges associated with implementing such programs with at-risk populations (Toth & Manly, 2011), and further research is needed.

The findings of the present study must be interpreted in the context of some limitations. Studies of gene-environment interaction have many requirements in order to assure credibility, foremost among which are large sample sizes. The present study's moderate size is consistent with those of many $G \times E$ studies in the literature. However, we acknowledge the need for even larger samples, and call for replications of the hypotheses in other and larger samples. The limited sample size also prevented exploration of gender differences; the current study controlled for gender, but it is likely that different patterns of reactivity to stressors may occur and need to be understood. As mentioned above, we also lacked adequate power to evaluate additional joint effects between CRHR1 and 5-HTTLPR. The moderate sample size is balanced by biologically and psychologically plausible hypotheses, and by high quality of measurement of the environmental factors, including a prospective, rather than long-term retrospective, measure of adversity obtained on multiple occasions in the first five years, thus improving on limitations due to memory, and possibly mitigating the biasing effects of current mood on questionnaire reports of adversity. The assessment of adversities limited to the first years of life represents an additional strength, as this may coincide with a sensitive period for the development of stress regulation systems (Heim & Binder, 2012). We also used a reliable and valid interview measure of current and recent chronic stress, and continuous measures of depressive symptoms and broadband internalizing symptoms.

Our measurement of CRHR1 variants was limited, and we based analyses on two highly correlated SNPs that had been prominent in published studies available at the time of genotyping (and we essentially modeled the TAT haplotype with two of three highly correlated SNPs). Nevertheless, it possible and even likely that different functions are served by various CRHR1 polymorphisms, and a fuller understanding of the role of this gene in the stress reactivity process will require many studies of a fuller array of SNPs.

We note that despite the advantages of the prospective assessment of childhood adversity, our measure has shortcomings. It did not include severe maltreatment, and it is possible that our measure did not capture as severe adversity as some other studies have done. However, the most common finding about adversities is that they cluster together, so that use of single focus measures may underestimate more common but perhaps less severe experiences. On the other hand, assessment across diverse content, while more ecologically valid in one sense, may obscure differences between types of adversity, as well as different levels of severity, which could provide important clues about their impact and mechanisms.

In conclusion, framed in terms of interactions among multiple levels of environment and biologically plausible genetic factors, the study supports predictions of increased likelihood of depressive and internalizing symptoms in response to stress among those with EA exposure and A alleles of the CRHR1 rs110402 SNP, and the S alleles of the 5-HTTLPR

gene. The study refines our understanding of which individuals under what conditions are more likely to experience depressive responses to stress, and is consistent with models of genetic susceptibility to the environment and of stress sensitization due to developmentally significant biological and psychosocial processes modified in ways that eventuate in maladaptive outcomes in the face of stressful experiences. Further studies will help to extend these findings with more complete understanding of the mechanisms of stress sensitization and of dysfunctional emotional and coping responses to stress.

Acknowledgments

This research was supported by NIMH R01MH052239 and NIMH T32MH082719 and by a seed grant from the UCLA Cousins Center for Psychoneuroimmunology. The authors thank the MUSP, M900, and M20 Research Teams, and participating families. We thank the principals of the Mater-University Study of Pregnancy study, the Brisbane, Australia staff coordinators, and the Genetic Epidemiology Unit of the Queensland Institute for Medical Research for genotyping of 5-HTTLPR, and for the assistance of the UCLA Inflammatory Biology Core, which is supported by NIH grant P30 AG028748, for the genotyping of CRHR1.

References

- Achenbach, TM. The Achenbach System of Empirically Based Assessment (ASEBA): Development, Findings, Theory, and Applications. Burlington, VT: University of Vermont Research Center for Children, Youth and Families; 2009.
- Andersen SL, Tomada A, Vincow ES, Valente E, Polcari AR, Teicher MH. Preliminary evidence for sensitive periods in the effect of childhood sexual abuse on regional brain development. *The Journal of Neuropsychiatry and Clinical Neurosciences*. 2008; 20:292–301. [PubMed: 18806232]
- Beck, AT., Steer, RA., Brown, GK. The Beck Depression Inventory. 2. San Antonio, TX: The Psychological Corporation; 1996.
- Beck AT, Steer RA, Garbin MG. Psychometric properties of the Beck Depression Inventory: Twenty-five years of evaluation. *Clinical Psychology Review*. 1988; 8:77–100.
- Bedford, A., Foulds, G. Delusions-Symptoms-States Inventory of anxiety and depression. Windsor, England: NFER; 1978.
- Belsky J, Pluess M. Beyond diathesis stress: Differential susceptibility to environmental influences. *Psychological Bulletin*. 2009; 135:885. [PubMed: 19883141]
- Bernstein, D., Fink, LA. Childhood Trauma Questionnaire Manual. San Antonio, TX: The Psychological Corporation; 1998.
- Bowlby, J. Attachment and loss. New York, NY, US: Basic Books; 1980.
- Bradley RG, Binder EB, Epstein MP, Tang Y, Nair HP, Liu W, ... Newport DJ. Influence of child abuse on adult depression: moderation by the corticotropin-releasing hormone receptor gene. *Archives of General Psychiatry*. 2008; 65:190. [PubMed: 18250257]
- Brown GW, Harris TO. Depression and the serotonin transporter 5-HTTLPR polymorphism: A review and a hypothesis concerning gene-environment interaction. *Journal of Affective Disorders*. 2008; 111:1–12. [PubMed: 18534686]
- Carpenter LL, Tyrka AR, McDougle CJ, Malison RT, Owens MJ, Nemeroff CB, Price LH. Cerebrospinal fluid corticotropin-releasing factor and perceived early-life stress in depressed patients and healthy control subjects. *Neuropsychopharmacology*. 2004; 29:777–784. [PubMed: 14702025]
- Carpenter LL, Tyrka AR, Ross NS, Khoury L, Anderson GM, Price LH. Effect of childhood emotional abuse and age on cortisol responsivity in adulthood. *Biological Psychiatry*. 2009; 66:69–75. <http://dx.doi.org/10.1016/j.biopsych.2009.02.030>. [PubMed: 19375070]
- Caspi A, Hariri AR, Holmes A, Uher R, Moffitt TE. Genetic sensitivity to the environment: The case of the serotonin transporter gene and its implications for studying complex diseases and traits. *American Journal of Psychiatry*. 2010; 167:509–527. [PubMed: 20231323]

- Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, et al. Influence of life stress on depression: Moderation by a polymorphism in the 5-HTT gene. *Science*. 2003; 301(5631):386–389. [PubMed: 12869766]
- Cicchetti D. Annual research review: Resilient functioning in maltreated children—past, present, and future perspectives. *Journal of Child Psychology and Psychiatry*. 2013; 54:402–422. [PubMed: 22928717]
- Cicchetti D, Dawson G. Editorial: Multiple levels of analysis. *Development and Psychopathology*. 2002; 14:417–420. [PubMed: 12349866]
- Cicchetti D, Rogosch FA. The impact of child maltreatment and psychopathology on neuroendocrine functioning. *Development and Psychopathology*. 2001; 13(4):783–804. [PubMed: 11771908]
- Cicchetti D, Rogosch FA, Oshri A. Interactive effects of corticotropin releasing hormone receptor 1, serotonin transporter linked polymorphic region, and child maltreatment on diurnal cortisol regulation and internalizing symptomatology. *Development and Psychopathology*. 2011; 23(4): 1125. [PubMed: 22018085]
- Cicchetti D, Toth SL. The past achievements and future promises of developmental psychopathology: The coming of age of a discipline. *Journal of Child Psychology and Psychiatry*. 2009; 50:16–25. DOI: 10.1111/j.1469-7610.2008.01979.x [PubMed: 19175810]
- Comasco E, Aslund C, Orelund L, Nilsson KW. Three-way interaction effect of 5-HTTLPR, BDNF Val66Met, and childhood adversity on depression: A replication study. *European Neuropsychopharmacology*. 2013; Advance online publication. doi: 10.1016/j.euroneuro.2013.01.010
- DeYoung CG, Cicchetti D, Rogosch FA. Moderation of the association between childhood maltreatment and neuroticism by the corticotropin-releasing hormone receptor 1 gene. *Journal of Child Psychology and Psychiatry*. 2011; 52:898–906. [PubMed: 21438878]
- Du L, Bakish D, Hrdina PD. Gender differences in association between serotonin transporter gene polymorphism and personality traits. *Psychiatric Genetics*. 2000; 10:159. [PubMed: 11324940]
- Dunn EC, McLaughlin KA, Slopen N, Rosand J, Smoller JW. Developmental timing of child maltreatment and symptoms of depression and suicidal ideation in young adulthood: Results from the National Longitudinal Study of Adolescent Health. *Depression and Anxiety*. 2013; doi: 10.1002/da.22102]
- First, MB., Spitzer, RL., Gibbon, M., Williams, JBW. *Structured Clinical Interview for DSM-IV Axis I disorders*. Washington, D.C: American Psychiatric Press; 1995.
- Fisher PA, Gunnar MR, Chamberlain P, Reid JB. Preventive intervention for maltreated preschool children: Impact on children's behavior, neuroendocrine activity, and foster parent functioning. *Journal of the American Academy of Child & Adolescent Psychiatry*. 2000; 39:1356–1364. <http://dx.doi.org/10.1097/00004583-200011000-00009>. [PubMed: 11068890]
- Gillespie CF, Phifer J, Bradley B, Ressler KJ. Risk and resilience: Genetic and environmental influences on development of the stress response. *Depression and Anxiety*. 2009; 26:984–992. [PubMed: 19750552]
- Gonzalez A, Jenkins JM, Steiner M, Fleming AS. The relation between early life adversity, cortisol awakening response and diurnal salivary cortisol levels in postpartum women. *Psychoneuroendocrinology*. 2009; 34:76–86. [PubMed: 18835661]
- Gotlib, IH., Colich, NL. Children of depressed parents. In: Gotlib, I., Hammen, C., editors. *Handbook of Depression*. 3. NY: Guilford Press; in press
- Gotlib IH, Joormann J, Minor KL, Hallmayer J. HPA axis reactivity: A mechanism underlying the associations among 5-HTTLPR, stress, and depression. *Biological Psychiatry*. 2008; 63:847–851. [PubMed: 18005940]
- Grabe HJ, Schwahn C, Mahler J, Schulz A, Spitzer C, Fenske K, ... Freyberger HJ. Moderation of adult depression by the serotonin transporter promoter variant (5-HTTLPR), childhood abuse and adult traumatic events in a general population sample. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*. 2012; 159B:298–309.
- Gunnar M, Quevedo K. The neurobiology of stress and development. *Annual Review of Psychology*. 2007; 58:145–173.

- Gunnar MR, Van Dulmen MHM. Behavior problems in postinstitutionalized internationally adopted children. *Development and Psychopathology*. 2007; 19:129–148. [PubMed: 17241487]
- Hammen C, Adrian C, Gordon D, Burge D, Jaenicke C, Hiroto D. Children of depressed mothers: Maternal strain and symptom predictors of dysfunction. *Journal of Abnormal Psychology*. 1987; 96:190–198. [PubMed: 3680756]
- Hammen C, Brennan P. Depressed adolescents of depressed and nondepressed mothers: Tests of an interpersonal impairment hypothesis. *Journal of Consulting and Clinical Psychology*. 2001; 69:284–294. [PubMed: 11393605]
- Hammen C, Brennan P, Keenan-Miller D. Patterns of adolescent depression to age 20: The role of maternal depression and youth interpersonal dysfunction. *Journal of Abnormal Child Psychology*. 2008; 36:1189–1198. [PubMed: 18473162]
- Hammen C, Brennan P, Keenan-Miller D, Hazel N, Najman J. Chronic and acute stress, gender, and serotonin transporter gene-environment interactions predicting depression in youth. *Journal of Child Psychiatry and Psychology*. 2010; 51:180–187.
- Hammen C, Hazel N, Brennan P, Najman J. Intergenerational transmission and continuity of stress and depression: Depressed women and their offspring in 20 years of follow-up. *Psychological Medicine*. 2012; 42:931–942. [PubMed: 22018414]
- Hammen C, Henry R, Daley S. Depression and sensitization to stressors among young women as a function of childhood adversity. *Journal of Consulting and Clinical Psychology*. 2000; 68:782–787. [PubMed: 11068964]
- Hammen, C., Rudolph, KR., Abaied, J. Child and Adolescent Depression. In: Mash, EJ., Barkley, RA., editors. *Child Psychopathology*. 3. New York: Guilford; in press
- Harkness KL, Bruce AE, Lumley MN. The role of childhood abuse and neglect in the sensitization to stressful life events in adolescent depression. *Journal of Abnormal Psychology*. 2006; 115:730–741. [PubMed: 17100530]
- Hayes, AF. *Introduction to Mediation, Moderation, and Conditional Process Analysis: A Regression-Based Approach*. New York: Guilford; 2013.
- Hazel N, Hammen C, Brennan P, Najman J. Early childhood adversity and adolescent depression: Mediating role of continued stress. *Psychological Medicine*. 2008; 38:581–589. [PubMed: 18261247]
- Heim C, Binder EB. Current research trends in early life stress and depression: Review of human studies on sensitive periods, gene–environment interactions, and epigenetics. *Experimental Neurology*. 2012; 233:102–111. <http://dx.doi.org/10.1016/j.expneurol.2011.10.032>. [PubMed: 22101006]
- Heim C, Bradley B, Mletzko TC, Deveau TC, Musselman DL, Nemeroff CB, ... Binder EB. Effect of childhood trauma on adult depression and neuroendocrine function: Sex-specific moderation by CRH Receptor 1 gene. *Frontiers in Behavioral Neuroscience*. 2009; 3:41. [PubMed: 20161813]
- Heim C, Nemeroff CB. The role of childhood trauma in the neurobiology of mood and anxiety disorders: Preclinical and clinical studies. *Biological Psychiatry*. 2001; 49:1023–1039. [PubMed: 11430844]
- Heim C, Newport DJ, Mletzko T, Miller AH, Nemeroff CB. The link between childhood trauma and depression: insights from HPA axis studies in humans. *Psychoneuroendocrinology*. 2008; 33(6): 693–710. [PubMed: 18602762]
- Homberg JR, van den Hove DLA. The serotonin transporter gene and functional and pathological adaptation to environmental variation across the life span. *Progress in Neurobiology*. 2012; 99:117–127. [PubMed: 22954594]
- Hsu DT, Mickey BJ, Langenecker SA, Heitzeg MM, Love TM, Wang H, ... Zubieta JK. Variation in the corticotropin-releasing hormone receptor 1 (CRHR1) gene influences fMRI signal responses during emotional stimulus processing. *Journal of Neuroscience*. 2012; 32:3253–3260. [PubMed: 22378896]
- Institute of Medicine and National Research Council. *New directions in child abuse and neglect research*. Washington, DC: National Academies Press; 2013.
- Ishitobi Y, Nakayama S, Yamaguchi K, Kanehisa M, Higuma H, Maruyama Y, ... Akiyoshi J. Association of CRHR1 and CRHR2 with major depressive disorder and panic disorder in a

- Japanese population. *Am J Med Genet B Neuropsychiatr Genet.* 2012; 159B:429–436. [PubMed: 22467522]
- Karg K, Burmeister M, Shedden K, Sen S. The serotonin transporter promoter variant (5-HTTLPR), stress, and depression meta-analysis revisited: Evidence of genetic moderation. *Archives of General Psychiatry.* 2011; 68:444–454. [PubMed: 21199959]
- Keeping JD, Najman JM, Morrison J, Western JS, Andersen MJ, Williams GM. A prospective longitudinal study of social, psychological, and obstetrical factors in pregnancy: Response rates and demographic characteristics of the 8,556 respondents. *British Journal of Obstetrics and Gynecology.* 1989; 96:289–297.
- Keiley MK, Howe TR, Dodge KA, Bates JE, Pettit GS. The timing of child physical maltreatment: A cross-domain growth analysis of impact on adolescent externalizing and internalizing problems. *Development and Psychopathology.* 2001; 13:891–912. [PubMed: 11771913]
- Kendler K, Baker J. Genetic influences on measures of the environment. *Psychological Medicine.* 2007; 37:615–626. [PubMed: 17176502]
- Kessler RC, Davis CG, Kendler KS. Childhood adversity and adult psychiatric disorder in the U.S. National Comorbidity Survey. *Psychological Medicine.* 1997; 27:1101–1119. [PubMed: 9300515]
- Kessler RC, Magee WJ. Childhood adversities and adult depression: Basic patterns of association in a US national survey. *Psychological Medicine.* 1993; 23(3):679–690. [PubMed: 8234575]
- Kessler RC, McGonagle KA, Swartz M, Blazer DG, Nelson CB. Sex and depression in the National Comorbidity Survey I: Lifetime prevalence, chronicity, and recurrence. *Journal of Affective Disorders.* 1993; 29:85–96. [PubMed: 8300981]
- Kessler R, McLaughlin K, Green J, Gruber M, Sampson N, Zaslavsky A, ... Williams D. Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *British Journal of Psychiatry.* 2010; 197:378–385. [PubMed: 21037215]
- Lasa L, Ayuso-Mateos JL, Vazquez-Barquero JL, Diez-Manrique FJ, Dowrick CF. The use of the Beck Depression Inventory to screen for depression in the general population: A preliminary analysis. *Journal of Affective Disorders.* 2000; 57:261–265. [PubMed: 10708841]
- Lewis G, Collishaw S, Harold G, Rice F, Thapar A. Maternal depression and child and adolescent depression symptoms: An exploratory test for moderation by *CRHR1*, *FKBP5* and *NR3C1* gene variants. *Behavior Genetics.* 2012; 42:121–132. [PubMed: 21789663]
- Mandelli L, Serretti A, Marino E, Pirovano A, Calati R, Colombo C. Interaction between serotonin transporter gene, catechol-O-methyltransferase gene and stressful life events in mood disorders. *The International Journal of Neuropsychopharmacology.* 2007; 10:437–447. [PubMed: 16756688]
- Masten A. Resilience in developing systems: Progress and promise as the fourth waves rises. *Development and Psychopathology.* 2007; 19:921–930. [PubMed: 17705908]
- McLaughlin K, Conron K, Koenen K, Gilman S. Childhood adversity, adult stressful life events, and risk of past year psychiatric disorder: a test of the stress sensitization hypothesis in a population based sample of adults. *Psychological Medicine.* 2010; 40:1647–1658. [PubMed: 20018126]
- McLaughlin K, Green J, Gruber M, Sampson N, Zaslowsky A, Kessler R. Childhood adversities and first onset of psychiatric disorders in a national sample of US adolescents. *Archives of General Psychiatry.* 2012; 69:1151–1160. [PubMed: 23117636]
- Moffitt T, Caspi A, Rutter M. Strategy for investigating interactions between measured genes and measured environments. *Archives of General Psychiatry.* 2005; 62:473–481. [PubMed: 15867100]
- Monroe SM, Reid MW. Gene-environment interactions in depression research: Genetic polymorphisms and life-stress polyprocedures. *Psychological Science.* 2008; 19:947–956. [PubMed: 19000200]
- Munafò MR, Brown SM, Hariri AR. Serotonin transporter (5-HTTLPR) genotype and amygdala activation: A meta-analysis. *Biological Psychiatry.* 2008; 63:852–857. [PubMed: 17949693]
- National Research Council. Preventing mental, emotional, and behavioral disorders among young people: Progress and possibilities. The National Academies Press; 2009.
- Papiol S, Arias B, Gastó C, Gutiérrez B, Catalán R, Fañanás L. Genetic variability at HPA axis in major depression and clinical response to antidepressant treatment. *Journal of Affective Disorders.* 2007; 104:83–90. [PubMed: 17467808]

- Pariante CM, Lightman SL. The HPA axis in major depression: Classical theories and new developments. *Trends in Neurosciences*. 2008; 31:464–468. [PubMed: 18675469]
- Pergamin-Hight L, Bakermans-Kranenburg MJ, van IJzendoorn MH, Bar-Haim Y. Variations in the promoter region of the serotonin transporter gene and biased attention for emotional information: A meta-analysis. *Biological Psychiatry*. 2012; 71:373–379. [PubMed: 22138391]
- Polanczyk G, Caspi A, Williams B, Price TS, Danese A, Sugden K, ... Moffitt TE. Protective effect of CRHR1 gene variants on the development of adult depression following childhood maltreatment: Replication and extension. *Archives of General Psychiatry*. 2009; 66:978–985. [PubMed: 19736354]
- Posener JA, DeBattista C, Williams GH, Chmura Kraemer H, Kalehzan BM, Schatzberg AF. 24-h monitoring of cortisol and corticotropin secretion in psychotic and nonpsychotic major depression. *Archives of General Psychiatry*. 2000; 57:755–760. [PubMed: 10920463]
- Ray L, Sehl M, Bujarski S, Hutchison K, Blaine S, Enoch MA. The CRHR1 gene, trauma exposure, and alcoholism risk: A test of G x E effects. *Genes, Brain, and Behavior*. 2013; 12:361–369.
- Ressler K, Bradley B, Mercer K, Deveau T, Smith A, Gillespie C, ... Binder E. Polymorphisms in CRHR1 and the serotonin transporter loci: Gene x gene x environment interactions on depressive symptoms. *American Journal of Medical Genetics B: Neuropsychiatric Genetics*. 2010; 153B:812–24. [PubMed: 20029939]
- Risch N, Herrell R, Lehner T, Liang KY, Eaves L, Hoh J, ... Merikangas. Interaction between the serotonin transporter gene (5-HTTLPR), stressful life events, and risk of depression. *Journal of the American Medical Association*. 2009; 301:2462–2471. [PubMed: 19531786]
- Rogeness GA, McClure EB. Development and neurotransmitter-environmental interactions. *Development and Psychopathology*. 1996; 8(1):183–199.
- Rutter M. Environmentally mediated risks for psychopathology: Research strategies and findings. *Journal of the American Academy of Child and Adolescent Psychiatry*. 2005; 44:3–18. [PubMed: 15608539]
- Spanier GB. Measuring dyadic adjustment: New scales for assessing the quality of marriage and similar dyads. *Journal of Marriage and the Family*. 1976; 38:15–28.
- Starr LR, Hammen C, Brennan PA, Najman JM. Serotonin transporter gene as a predictor of stress generation in depression. *Journal of Abnormal Psychology*. 2012; 4:810–818.
- Taylor SE, Karlamangla AS, Friedman EM, Seeman TE. Early environment affects neuroendocrine regulation in adulthood. *Social Cognitive & Affective Neuroscience*. 2011; 6(2):244–251. [PubMed: 20400490]
- Toth SL, Manly JT. Bridging research and practice: Challenges and successes in implementing evidence-based preventive intervention strategies for child maltreatment. *Child Abuse & Neglect*. 2011; 35:633–636. <http://dx.doi.org/10.1016/j.chiabu.2011.05.003>. [PubMed: 21855142]
- Toth, SL., Rogosch, FA., Manly, JT., Cicchetti, D. The efficacy of toddler-parent psychotherapy to reorganize attachment in the young offspring of mothers with major depressive disorder: A randomized preventive trial. Vol. 74. US: American Psychological Association; 2006. p. 1006-1016.
- Uher R, McGuffin P. The moderation by the serotonin transporter gene of environmental adversity in the etiology of depression: 2009 update. *Molecular Psychiatry*. 2010; 15:18–22. [PubMed: 20029411]
- van IJzendoorn MH, Schuengel C, Bakermans-Kranenburg MJ. Disorganized attachment in early childhood: Meta-analysis of precursors, concomitants, and sequelae. *Development and Psychopathology*. 1999; 11:225–249. [PubMed: 16506532]
- Wasserman D, Wasserman J, Rozanov V, Sokolowski M. Depression in suicidal males: Genetic risk variants in the CRHR1 gene. *Genes, Brain and Behavior*. 2009; 8:72–79.
- Wasserman D, Wasserman J, Sokolowski M. Genetics of HPA-axis, depression and suicidality. *European Psychiatry*. 2010; 25(5):278–280. [PubMed: 20444578]
- Way B, Taylor SE. Social influences on health: Is serotonin a critical mediator? *Psychosomatic Medicine*. 2010; 72:107–112. [PubMed: 20145277]

- Wendland JR, Martin BJ, Kruse MR, Lesch KP, Murphy DL. Simultaneous genotyping of four functional loci of human *slc6a4*, with a reappraisal of 5-HTTLPR and rs25531. *Molecular Psychiatry*. 2006; 11:224–226. [PubMed: 16402131]
- Wisner Fries AB, Shirtcliff EA, Pollak SD. Neuroendocrine dysregulation following early social deprivation in children. *Developmental Psychobiology*. 2008; 50(6):588–599. [PubMed: 18683181]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Figure 1.

Illustration of three-way interaction between CRHR1 rs110402 genotype, early adversity, and age 20 chronic stress, predicting age 20 depressive symptoms, with differential association between chronic stress and depressive symptoms by genotype presented at low and high levels of early adversity (defined as 10th and 90th percentile, respectively). For ease of visual interpretation, CRHR1 rs110402 genotype has been dichotomized into G/G versus AG and AA.

Figure 2.

Illustration of three-way interaction between 5-HTTLPR genotype, early adversity, and age 20 chronic stress, predicting age 20 depressive symptoms, with differential association between chronic stress and depressive symptoms by genotype presented at low and high levels of early adversity (defined as 10th and 90th percentile, respectively). For ease of visual interpretation, 5-HTTLPR genotype has been dichotomized into L/L versus S/L and S/S.

Table 1
Study Variables Means and Demographic Information for Full Sample and by 5-HTTLPR and CRHR1 Genotype

	Full Sample	CRHR1 rs110402 Genotype			5-HTTLPR Genotype		
	<i>M (SD)</i>	G/G (<i>n</i> =140)	A/G (<i>n</i> =208)	A/A (<i>n</i> =96)	L/L (<i>n</i> = 101)	S/L (<i>n</i> =189)	S/S (<i>n</i> =91)
Early Adversity by age 5	1.77 (1.54)	1.74	1.83	1.83	1.92	1.86	1.55
Chronic stress age 20	22.59 (4.52)	22.90	22.73	21.81	22.30	22.99	21.96
BDI age 20	7.05 (8.40)	7.46	7.94	6.25	7.63	8.43	6.33
Demographic Characteristics							
Gender							
Male	50.6%	12.2%	19.8%	9.0%	11.5%	18.4%	9.2%
Female	49.4%	19.4%	27.0%	12.6%	15.0%	31.2%	14.7%
Race							
White	91.4%	30.6%	42.8%	18.7%	25.2%	46.7%	21.0%
Asian	3.6%	0.5%	1.8%	1.8%	0.0%	1.3%	2.1%
Maori/Islander	1.0%	0.0%	0.9%	0.2%	0.0%	0.8%	0.3%
Australian Aborigine	1.0%	0.2%	0.2%	0.0%	0.3%	0.0%	0.3%

Notes. *N*s for full sample range from 633 to 815. Group differences in study variables by genotype are not significant (*p*s > .05). Race reflects maternally-reported self-identification; additional participants did not report race or identified with another group.

Table 2

Bivariate Correlations between Major Study Variables

	1.	2.	3.	4.	5.	6.
1. Early Adversity before Age 5	---					
2. Age 20 Chronic Stress	.26***	---				
3. Age 20 BDI	.20***	.48***	---			
4. Age 20 Internalizing	.14**	.42***	.71***	---		
5. CRHR1 rs110402 # of A alleles	.02	-.08	-.04	-.04	---	
6. 5-HTTLPR # of S alleles	-.08	-.02	-.05	-.03	.05	---

Notes.

**
 $p < .01$, $p < .001$.