Conclusions

Clinical guidelines provide a useful reference for practising clinicians. However, they are often long, not easily approachable, and therefore underused. The sample case summaries presented here and the treatment flow chart in figure 1 provide a basis for the practical implementation of current guidelines and more recent clinical trial evidence.

Contributors: This review is a combined effort on the part of the two authors. CBC wrote the case examples.

Competing interests: CBC has been a consultant and speaker for Biotechnology General Corp, Boehringer Ingelheim, Glaxo-SmithKline, Pfizer, and VIASYS Healthcare. He is a member of the scientific advisory board of Boehringer Ingelheim/Pfizer for tiotropium related questions. He has also received research grants from Biotechnology General Corp, Boehringer Ingelheim, GlaxoSmithKline, ONO Pharma, and Pfizer. DPT has been a consultant and speaker for AstraZeneca, Boehringer Ingelheim, Forest, GlaxoSmithKline, and Pfizer, and he has received research grants from Altana, AstraZeneca, Boehringer Ingelheim, Forest, GlaxoSmithKline, and Pfizer.

- 1 Ezzati M, Lopez AD. Estimates of global mortality attributable to smoking in 2000. *Lancet* 2003;362:847-52.
- 2 Global Initiative for Chronic Obstructive Lung Disease. *Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease*. Bethesda, MD: National Heart Lung and Blood Institute, World Health Organization, 2003. www.goldcopd.com/2004clean.pdf (accessed 19 Jan 2005).
- 3 Celli BR, MacNee W, and committee members. Standards for the diagnosis and treatment of patients with COPD: a summary of the ATS/ERS position paper. *Eur Respir J* 2004;23:932-46.
- 4 National Institute for Clinical Excellence. *Chronic obstructive pulmonary disease*. *Thorax* 2004;59:1-232.
- 5 O'Donnell DE, Aaron S, Bourbeau J, Hernandez P, Marciniuk D, Balter M, et al. Canadian Thoracic Society recommendations for management of chronic obstructive pulmonary disease, 2003. *Can Respir J* 2003;10(suppl A):11A-65A.
- 6 Ramsey SD. Suboptimal medical therapy in COPD: exploring the causes and consequences. *Chest* 2000;117:33S-7S.
- 7 Brusasco V, Hodder R, Miravittles M, Lee A, Towse LJ, Kesten S. Health outcomes in a 6-month placebo controlled trial of once-daily tiotropium compared with twice-daily salmeterol in patients with COPD. *Thorax* 2003;58:399-404.
- 8 Szafranski W, Cukier A, Ramirez A, Menga G, Sansores R, Nahabedian S, et al. Efficacy and safety of budesonide/formoterol in the management of chronic obstructive disease. *Eur Respir J* 2003;21:74-81.
- 9 Mahler DA, Wire P, Horstman D, Chang CN, Yates J, Fischer T, et al. Effectiveness of fluticasone propionate and salmeterol combination delivered

via the diskus device in the treatment of chronic obstructive pulmonary disease. *Am J Respir Crit Care Med* 2002;166:1084-91.

- 10 Calverley P, Pauwels R, Vestbo J, Jones P, Pride N, Gulsvik A, et al. Combined salmeterol and fluticasone in the treatment of chronic obstructive pulmonary disease: a randomised controlled trial. *Lancet* 2003;361:449-56.
- 11 Rennard SI, Serby CW, Ghafouri M, Johnson PA, Friedman M. Extended therapy with ipratropium is associated with improved lung function in patients with COPD. A retrospective analysis of data from seven clinical trials. *Chest* 1996;110:62-70.
- 12 Liesker JJW, Wijkstra PJ, Ten Hacken NHT, Koeter GH, Postma DS, Kerstjens HAM. A systematic review of the effects of bronchodilators on exercise capacity in patients with COPD. *Chest* 2002;121:597-608.
- 13 Tashkin DP, Cooper CB. The role of long-acting bronchodilators in the management of stable COPD. *Chest* 2004;125:249-59.
- 14 Donohue JF, Menjoge S, Kesten S. Tolerance to bronchodilating effects of salmeterol in COPD. *Respir Med* 2003;97:1014-20.
- 15 Celli B, ZuWallack R, Wang S, Kesten S. Improvement in resting inspiratory capacity and hyperinflation with tiotropium in COPD patients with increased static lung volumes. *Chest* 2003;124:1743-8.
- 16 Hattotuwa KL, Gzycski MJ, Ansari TW, Jeffery PK, Barnes NC. The effects of inhaled fluticasone on airway inflammation in chronic obstructive pulmonary disease: a double-blind, placebo-controlled biopsy study. *Am J Respir Crit Care Med* 2002;165:1592-6.
- 17 Alsaedi A, Sin DD, McAlister FA. The effects of inhaled corticosteroids in chronic obstructive pulmonary disease: a systematic review of randomized placebo-controlled trials. *Am J Med* 2002;2002:59-65.
- 18 Sutherland ER, Allmers H, Ayas NT, Venn AJ, Martin RJ. Inhaled corticosteroids reduce the progression of airflow limitation in chronic obstructive pulmonary disease: a meta-analysis. *Thorax* 2003;58:937-41.
- 19 Highland KB, Strange C, Heffner JE. Long-term effects of inhaled corticosteroids on FEV1 in patients with chronic obstructive pulmonary disease. A meta-analysis. *Ann Intern Med* 2003;138:969-73.
- 20 Lipworth BJ. Systemic adverse effects of inhaled corticosteroid therapy: A systematic review and meta-analysis. *Arch Intern Med* 1999;159:941-55.
- 21 van Noord JA, de Munck DRAJ, Bantje T, Hop WCJ, Akveld MLM, Bommer AM. Long-term treatment of chronic obstructive pulmonary disease with salmeterol and the additive effect of ipratropium. *Eur Respir J* 2000;15:878-85.
- 22 Tennant RC, Erin EM, Barnes PJ, Hansel TT. Long-acting beta 2-adrenoceptor agonists or tiotropium bromide for patients with COPD: is combination therapy justified? *Curr Opin Pharmacol* 2003;3:270-6.
- 23 Cazzola M, Marco FD, Santus P, Boveri B, Verga M, Matera MG, et al. The pharmacodynamic effects of single inhaled doses of formoterol, tiotropium and their combination in patients with COPD. *Pulm Pharmacol Ther* 2004;17:35-9.
- 24 Hanania NA, Darken P, Horstman D, Reisner C, Lee B, Davis S, et al. The efficacy and safety of fluticasone propionate (250 microg)/salmeterol (50 microg) combined in the Diskus inhaler for the treatment of COPD. *Chest* 2003;124:834-43.
- 25 Calverley PM, Boonsawat W, Cseke Z, Zhong N, Peterson S, Olsson H. Maintenance therapy with budesonide and formoterol in chronic obstructive pulmonary disease. *Eur Respir J* 2003;22:912-9.

(Accepted 1 December 2004)

Monitoring in chronic disease: a rational approach

Paul Glasziou, Les Irwig, David Mant

"Know which abnormality you are going to follow during treatment. Pick something you can measure."

Meador C. *A Little Book of Doctors' Rules*.
Lyons: IARC Press, 1999.

The ritual of routine visits for most chronic diseases usually includes monitoring to check on the progress or regress of the disease and the development of complications. Such checks require that we choose what to monitor, when to monitor, and how to adjust treatment. Poor choices in each can lead to poor control, poor use of time, and dangerous adjustments to treatment. For example, an audit of serum digoxin monitoring in a UK teaching hospital more than 20 years ago showed that the logic behind more than 80% of the tests requested could not be established, the timing of tests reflected poor understanding of the clinical pharmacokinetics, and about one result in four was followed by an inappropriate clinical decision.¹

Improvements are possible. For example, a computerised reminder of inappropriate testing reduced the volume of testing for the concentration of antiepileptic drugs by 20%²; a decision support system for anticoagulation with warfarin led to an improvement from 45% to 63% of patients being within target range³; and quality control charts for peak flow measurements for people with asthma could detect exacerbations four days earlier than conventional methods.⁴ Given the extent of monitoring, even modest improvements are likely to improve benefits for patients and may reduce costs.

Monitoring is periodic measurement that guides the management of a chronic or recurrent condition. It can be done by clinicians, patients, or both. In Australia, monitoring comprises between a third and half of all tests ordered in general practice and outpatients (Pirozzo, personal communication, 2002). Despite the considerable staff time and resources

University of
Oxford,
Department of
Primary Health
Care, Oxford
OX3 7LF UK
Paul Glasziou
professor
David Mant
professor
continued over

BMJ 2005;330:644-8

involved, monitoring is a surprisingly understudied area. We review the current literature (based on a Medline search using the terms “monitor*”, “therap*” or “treat*”, “limit*” or “threshold*”, “chronic” and a check of references of relevant papers found) on the clinical uses of monitoring, and develop some principles for good monitoring strategies.

Should we monitor at all?

Although intuitively monitoring should be beneficial, clinicians accept non-monitoring in many areas. Aspirin, for example, is used to prevent stroke without assessing platelet aggregation. Establishing benefit for patients is important, as it must be balanced against the downsides of monitoring, such as the inconvenience and costs, and the impact of false positive and false negative (monitoring) results that can lead to inappropriate or delayed actions. As with other interventions, this ideally requires a randomised controlled trial. The benefit should preferably be measured by outcomes that are relevant to patients rather than time spent in the target range (a surrogate marker). For example, in a trial of drug monitoring in epilepsy, monitored patients were more often within control limits than unmonitored patients (8% *v* 25%), but the proportion remaining free of seizures did not change (38% *v* 41%).⁵ However, few monitoring practices have undergone such trials.

Monitoring is a complex intervention. It can have an impact through several means, including improvement of adherence, better selection of treatments based on individual response, better titration of treatment, and patients' learning about non-treatment factors that alter the condition's control. Because of this complexity, any trial should be preceded by developmental work to understand the optimal strategies.

Monitoring and adjustment may be controlled by clinicians or patients. For patients, monitoring may provide a signal for action or simply provide motivation to adhere to treatment. For example, a recent trial in which some diabetes patients were randomised to self monitoring of risk factors showed a better achievement of target measurements of blood pressure, low density lipoprotein cholesterol, and HbA_{1c}, and a reduction in clinical events.⁶ However, a randomised trial of the motivational effect of cholesterol measurement in general practice in the United Kingdom showed only negligible benefit—a difference of 0.1 mmol/l in total cholesterol measurements.⁷

The optimal process is not straightforward. A recent study compared three modes of peak flow monitoring in childhood asthma: regular daily

measurements, measurement only when symptomatic, and monitoring only at times of symptoms. The second group—children randomised to measure their peak expiratory flow rate only when symptomatic—had lower asthma severity scores, fewer days of symptoms, and fewer healthcare visits than children randomised to either daily peak flow monitoring or monitoring only at times of symptoms.⁸ The study showed that peak flow monitoring is helpful but that the less intensive monitoring regimen is preferable.

The phases of monitoring

The objective and methods of monitoring change over the course of treatment. This course of monitoring can be usefully divided into the five phases listed in table 1.

Figure 1 shows a control chart for these stages. At (a) we first note the abnormal measurement and begin a quick series of measurements before treatment to confirm the abnormal result; (b) then, if appropriate, initiate treatment and monitor at short intervals to check response and achieve control; (c) but once control is achieved, the intervals may be longer (d), although this may be supplemented by patients' self monitoring (small arrows) (e); but when one measurement is more than 3 standard deviations (SD) or two measurements are more than 2 standard deviations from target, we adjust therapy to re-establish control and shorten the re-check interval; and finally (f), if treatment becomes unnecessary, a period of monitoring after its cessation may be required. We now look at these phases in more detail.

Pretreatment monitoring

Monitoring before treatment should establish the need for treatment and then a baseline to judge the response to treatment or changes in the patient's condition. Treatment should not start until sufficient measurements for a firm baseline have been obtained. This firm baseline confirms that the degree of abnormality is beyond the initiation threshold. Serial measurements often “normalise” before treatment for several reasons, such as training effects (for example, with a peak flow meter), accommodation to measurement (for example, with blood pressure measurement), and, perhaps most importantly, regression to the mean (the tendency of repeat tests to be closer to normal⁹). For example, in the first biennial check in the Framingham study, the average blood pressure fell by 3.4 mm Hg systolic and 2.4 mm Hg diastolic.¹⁰ Among 99 Dutch patients with apparently raised blood pressure, re-measurement resulted in average reductions of 9 mm Hg systolic and 4 mm Hg diastolic.¹¹ The same study showed that two measurements were sufficient to

Screening and Test Evaluation Program, School of Public Health, University of Sydney, Camperdown, NSW 2006, Australia

Les Irwig
professor

Correspondence to:
P Glasziou
paul.glasziou@
dphpc.ox.ac.uk

The objectives for the five phases of monitoring

Phase	Monitoring objectives	Optimal interval
Pretreatment	Check need for treatment Establish a baseline for determining response and change	Short—based on variability within patients
Initial titration	Assess individual response to treatment Assess immediate adverse effects Achieve control	Medium—based on pharmacokinetics (such as drug half life) and pharmacodynamics (physiological impact time) (wash-in)
Maintenance	Detect drift from control limits Detect long term harms	Long—based on rate of random and systematic “drift”
Re-establish control	Bring level back within control limits	Medium—as above
Cessation	Check safety of cessation	Medium—as above

establish the need for treatment in patients who were well above the initiation threshold. For borderline patients, even four measurements resulted in substantial misclassification.

Initial titration: response, control, and safety

After establishing the baseline, we should set a target and start titration to achieve that target. However, achieving the target is just one of several objectives of the initial titration phase, which should include checking the individual's response to treatment, detecting unacceptable adverse effects, and achieving the desired target range.

This initial monitoring checks adequate response to treatment—that is, whether it “works” as expected on the basis of clinical trials in other patients. Sometimes individual response to treatment can be predicted by other measurements, such as genetic testing (for example, the dose requirement on initiating warfarin treatment is strongly related to CYP2C9 gene variants¹²). Sometimes it can be predicted by pharmacokinetic studies—for example, in initiating tricyclic antidepressants, short term measurement of the drug concentration can characterise individual metabolism and guide the long term dose used.¹³ However, personalising treatment usually requires some pragmatic trial and error. We can therefore think of this phase as an “n = 1” trial.¹⁴ Ideally, the estimate of effect will be based on both the measurements in that patient and the known effect from trials.¹⁵

The initiation phase should also detect immediate or short term adverse effects.¹⁶ Measures of potential harm should hence be assessed and monitored. Almost half of the medicines in the electronic Medicines Compendium (www.medicines.org.uk) include a suggestion for some monitoring. Many drugs affect renal function, and regular monitoring of creatinine and electrolytes is widely recommended. A more specific example is treatment with clozapine, for which white cell counts are measured on a weekly basis for the first few months to detect agranulocytosis, which occurs in 0.8% of patients. However, the rationale for the timing of measurement is seldom explicit. The criteria for monitoring for adverse effects are similar to those for screening, including that the effect is serious, that a simple test is available, that earlier detection is predictive, and that change in treatment leads to a better outcome.¹⁷

Fig 1 The five phases of treatment monitoring

Monitoring during treatment

Once the target is achieved, the objective of monitoring is to ensure that measurements stay within reasonable limits, called control limits. The control limits ensure that we detect real changes in the level of the target measure while minimising false positives resulting from variable measurements in the short term or errors in the technical measurement. The degree of short term variation can be estimated from population studies, subpopulations, or from the individual's own measurements. Because extreme measurements are unlikely to be due to variability of short term measurements, they may justify action to re-establish control. One approach suggested by statistical control theory is to consider that a shift from control has occurred if a single measurement is outside an upper and lower control limit of 3 standard deviations, or if two or three successive measurements are more than 2 standard deviations from the target.¹⁸ Figure 1 shows these two sets of action thresholds; one for action (SD 3) and one for re-measurement (SD 2), with action if the repeat result is also more than 2 standard deviations from target.

Monitoring during treatment can be less frequent than during the initiation phase. The interval depends on the probability of being outside the control limits, which in turn depends on both random drift and systematic changes (progression or regression of disease).

The measurements intervals may be shorter than the decision interval. For example, monitoring measurements of blood pressure might be done daily by patients, but the decision made at a monthly consultation with the doctor. This is illustrated in figure 1, where multiple measures (3(b)—small arrows) occur between the decision points (3(a)—large arrows). Ideally a graphical presentation, such as a control chart, should be used to aid recording and deciding treatment changes.

Adjustment to re-establish control

When a clear drift beyond the control limits occurs, we should re-establish control. As in the initiation phase, a shorter measurement interval is generally warranted until control is re-established.

We have already cited several examples (for example, monitoring digoxin therapy) in which audits have shown that clinical decisions taken as a result of monitoring are suboptimal—sometimes tending to the conservative. A New Zealand study of digoxin monitoring showed that 53% of ordered measurements were inappropriately timed and that 5% of the ordered measurements led to inappropriate dose adjustments.¹⁹ Patients too find this difficult: a survey of diabetes educators showed that correct adjustment of insulin dosage is the single hardest skill to teach.²⁰ This may be because of the lack of a planned and explicit response to test results or because the tests are seldom considered in context—they usually arrive as a single value, without reference to past values or a clear and useful statement about the variability of measurements. The appropriate degree of adjustment may be helped by nomograms or computer algorithms.

Cessation of treatment

Most therapies are not lifelong. Monitoring should inform stopping decisions. Again, such decisions are

Summary points

Monitoring aims to establish the response to treatment, detect the need to adjust treatment, and detect adverse effects

Monitoring is not always necessary or beneficial and can lead to inappropriate changes

Control charts help distinguish natural variability from true change and reduce unnecessary adjustment

Monitoring for both benefit and harm is important, preferably with a single measurement

The interval between measurements varies between phases and is shorter after changes in treatment

Contributors: PG conceived the study, and all authors contributed to the ideas and writing. PG is guarantor.

Funding: PG and LI were in part funded by the Australian National Health and Medical Research Council for this work, as part of a programme grant 211205 on testing.

Competing interests: None declared.

- 1 Clague HW, Twum-Barima Y, Carruthers SG. An audit of requests for therapeutic drug monitoring of digoxin: problems and pitfalls. *Ther Drug Monit* 1983;5:249-54.
- 2 Chen P, Tanasijevic MJ, Schoenenberger RA, Fiskio J, Kuperman GJ, Bates DW. A computer-based intervention for improving the appropriateness of antiepileptic drug level monitoring. *Am J Clin Pathol* 2003; 119:432-8.
- 3 Ryan PJ, Gilbert M, Rose PE. Computer control of anticoagulant dose for therapeutic management. *BMJ* 1989;299:1207-8.
- 4 Gibson PG, Wlodarczyk J, Hensley MJ, Murree-Allen K, Olson LG, Salto N. Using quality-control analysis of peak expiratory flow recordings to guide therapy for asthma. *Ann Intern Med* 1995;123:488-92.
- 5 Jannuzzi G, Cian P, Fattore C, Gatti G, Bartoli A, Monaco F, Perucca E. A multicenter randomized controlled trial on the clinical impact of therapeutic drug monitoring in patients with newly diagnosed epilepsy. The Italian TDM Study Group in Epilepsy. *Epilepsia* 2000;41:222-30.
- 6 Rachmani R, Levi Z, Slavachevski I, Avin M, Ravid M. Teaching patients to monitor their risk factors retards the progression of vascular

- complications in high-risk patients with Type 2 diabetes mellitus—a randomized prospective study. *Diabet Med* 2002;19:385-92.
- 7 Robertson I, Phillips A, Mant D, Thorogood M, Fowler G, Fuller A, Yudkin P, Woods M. Motivational effect of cholesterol measurement in general practice health checks. *Br J Gen Pract* 1992;42:469-72.
- 8 Yoos HL, Kitzman H, McMullen A, Henderson C, Sidor K. Symptom monitoring in childhood asthma: a randomized clinical trial comparing peak expiratory flow rate with symptom monitoring. *Ann Allerg Asthma Immunol* 2002;88:283-91.
- 9 Bland JM, Altman DG. Regression towards the mean. *BMJ* 1994; 308:1499.
- 10 Dauber TR. *The Framingham study: the epidemiology of atherosclerotic disease*. Cambridge, MA: Harvard University Press, 1980.
- 11 Brueren MM, Petri H, van Weel C, van Ree JW. How many measurements are necessary in diagnosing mild to moderate hypertension? *Fam Pract* 1997;14:130-5.
- 12 Peyvandi F, Spreafico M, Siboni SM, Moia M, Mannucci PM. CYP2C9 genotypes and dose requirements during the induction phase of oral anticoagulation therapy. *Clin Pharmacol Ther* 2004;75:198-203.
- 13 Braithwaite RA, Dawling S, Montgomery SA. Prediction of steady-state plasma concentrations and individual dosage regimens of tricyclic antidepressants from a single test dose. *Ther Drug Monit* 1982;4:27-31.
- 14 Guyatt GH, Keller JL, Jaeschke R, Rosenbloom D, Adachi JD, Newhouse MT. The n-of-1 randomized controlled trial: clinical usefulness. Our three-year experience. *Ann Intern Med* 1990;112:293-9.
- 15 Irwig L, Glasziou P, Wilson A, Macaskill P. Estimating an individual's true cholesterol level and response to intervention. *JAMA* 1991;266:1678-85.
- 16 Aronson JK, Ferner RE. Joining the DoTS: new approach to classifying adverse drug reactions. *BMJ* 2003;327:1222-5.
- 17 Pirmohamed M, Ferner RE. Monitoring drug treatment. *BMJ* 2003; 327:1179-81.
- 18 Boggs PB, Hayati FH, Washburne WF, Wheeler DA. Using statistical process control charts for the continual improvement of asthma care. *J Qual Improve* 1999;25:163-81.
- 19 Sidwell A, Barcaly M, Begg E, Moore G. Digoxin therapeutic monitoring: an audit and review. *N Z Med J* 2003;116:U704.
- 20 Bonnet C, Gagnayre R, d'Ivernois JF. Learning difficulties of diabetic patients: a survey of educators. *Patient Educ Couns* 1998;35:139-47.
- 21 Begg CB, Leung D. On the use of surrogate end points in randomized trials. *J R Stat Soc A* 2000;163:15-28.
- 22 Irwig L. This added to my multiple myopia. *BMJ* 2003;326:1336.
- 23 Scottish Intercollegiate Guidelines Network. Lipids and the primary prevention of coronary heart disease. Edinburgh: SIGN, 1999. www.sign.ac.uk/guidelines/fulltext/40/index.html (accessed 5 Jan 2005).
- 24 Smith DH, Matzek KM, Kempthorne-Rawson J. Dose response and safety of telmisartan in patients with mild to moderate hypertension. *J Clin Pharmacol* 2000;40:1380-90.
- 25 Lowy. A memorable patient: Home glucose monitoring, who started it? *BMJ* 1998;316:1467.
- 26 Cappuccio FP, Kerry SM, Forbes L, Donald A. Blood pressure control by home monitoring: meta-analysis of randomised trials. *BMJ* 2004; 329:145-8.
- 27 Thoonen BP, Schermer TR, van Den Boom G, Molema J, Folgering H, Akkermans RP, et al. Self-management of asthma in general practice, asthma control and quality of life: a randomised controlled trial. *Thorax* 2003;58:30-6.
- 28 Schroeder K, Fahey T, Ebrahim S. How can we improve adherence to blood pressure-lowering medication in ambulatory care? Systematic review of randomized controlled trials. *Arch Intern Med* 2004;164:722-32. (Accepted 20 December 2004)

Health Services
Research, Primary
Care Sciences
Research Centre,
Keele University,
Keele ST5 5BG
Bie Nio Ong
professor

Light Oaks, Stoke
on Trent ST2 7DE
Daphne Evans
patient

continued over

BMJ 2005;330:648-50

A patient's journey with myalgic encephalomyelitis

Bie Nio Ong, Daphne Evans, Andrew Bartlam

Myalgic encephalomyelitis (ME) presents patients and their general practitioners with the challenge of managing a life shaped by chronic debilitating illness, pain, and uncertainty. The notion that body limitations resulting from illness must be recognised is shown in the literature on chronic illness, but at the same time it is necessary to understand the social context of change and diversity within which illness is conceptualised. People give meaning to their illness and altered life course with reference to socioeconomic relations, cultural perceptions and beliefs, and their interactions with health and social care services.

Myalgic encephalomyelitis represents the conundrum of uncertainty in the diagnosis, treatment, and prognosis of the disease. Many of the symptoms that people present with could be indications for other diseases, making it difficult to diagnose. Moreover, there

is still some debate about the existence of the disease as a clinical category. Current guidance focuses on the symptoms of fatigue and malaise, cognitive impairment, and pain. The patient's account in this paper was collected as part of a research study on osteoarthritis of the knee. The patient, Daphne Evans, was interviewed by BNO, but the discussion centred on myalgic encephalomyelitis because she considered this condition of primary importance. Her doctor's account was added to her own account when the paper was drafted.

Diagnosing uncertain disease

Daphne's story illustrates the syndrome of myalgic encephalomyelitis as a chronic and fluctuating illness with a profusion of symptoms that affect many parts of