Published in final edited form as:

J Affect Disord. 2017 September ; 219: 193–200. doi:10.1016/j.jad.2017.04.042.

Severity of Anxiety– but not Depression– is Associated with Oxidative Stress in Major Depressive Disorder

Lisa R. Steenkamp^{1,2}, Christina M. Hough¹, Victor I. Reus¹, Felipe A. Jain¹, Elissa S. Epel¹, S. Jill James³, Alexandra E. Morford¹, Synthia H. Mellon⁴, Owen M. Wolkowitz^{1,*}, and Daniel Lindqvist^{1,5}

¹Department of Psychiatry, University of California San Francisco (UCSF) School of Medicine, San Francisco, California, United States of America ²Institute of Psychology, Leiden University, Leiden, the Netherlands ³Arkansas Children's Research Institute, Department of Pediatrics, University of Arkansas for Medical Sciences, Little Rock, Arkansas, United States of America ⁴Department of OB/GYN and Reproductive Sciences, University of California San Francisco (UCSF) School of Medicine, San Francisco, California, United States of America ⁵Lund University, Faculty of Medicine, Department of Clinical Sciences, Lund, Psychiatry, Sweden

Abstract

Background—Oxidative stress is implicated in both depression and anxiety, but it is currently unclear whether this relates to syndromal diagnoses or trans-diagnostic dimensional symptoms. We examined the relationship between oxidative stress and severity of depression and anxiety symptoms in individuals with Major Depressive Disorder (MDD).

Methods—Plasma oxidative stress markers F2-isoprostanes and oxidized glutathione (GSSG), and the antioxidant reduced glutathione (GSH), were assessed in 69 physically healthy, medication-free MDD subjects. Symptoms of anxiety and depression were assessed using the Hamilton Anxiety (HAM-A) and Hamilton Depression (HAM-D) Rating Scales. Total HAM-A and HAM-D scores, along with “core” anxiety and depression subscales, and individual HAM-D items “psychic anxiety” and “depressed mood,” were related to oxidative stress markers. Analyses controlled for age, sex, BMI, and smoking.

Results—Total HAM-A ratings were positively associated with F2-isoprostanes ($\beta=.26$, $p=0.042$) and GSSG ($\beta=.25$, $p=0.049$), but not GSH ($\beta=.05$, $p=0.711$). Core anxiety severity was positively associated with F2-isoprostanes ($\beta=.34$, $p=0.012$) and GSSG, although this did not reach significance ($\beta=.24$, $p=0.074$). None of the biological markers were significantly associated with total HAM-D or core depression ratings (all $p>0.13$). Subjects scoring high on “psychic anxiety” had elevated F2-isoprostanes ($p=0.030$) and GSSG ($p=0.020$). This was not seen with “depressed mood” scores (all $p>0.12$).

*Corresponding author: Owen M. Wolkowitz, MD, PhD, Department of Psychiatry, University of California, San Francisco (UCSF), School of Medicine, 401 Parnassus Avenue, San Francisco, CA 94143, USA, Phone: 415-476-7433, Owen.Wolkowitz@ucsf.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Limitations—We assessed peripheral oxidative markers, but their relationship to the brain is unclear.

Conclusions—Oxidative stress is more closely related to anxiety than depression symptoms in MDD. This highlights the importance of relating oxidative stress to specific symptoms and could provide new insights into the biological correlates of affective disorders.

Keywords

Major Depressive Disorder; Oxidative Stress; Anxiety; F2-isoprostanes; Oxidized Glutathione; Reduced Glutathione

1. Introduction

Individuals with Major Depressive Disorder (MDD) have an increased risk of developing medical illnesses, including cardiovascular diseases, diabetes, obesity, Alzheimer's disease, and cancer (Penninx et al., 2013), conditions that have also been associated with oxidative stress (Ceriello and Motz, 2004, Emerit et al., 2004, Furukawa et al., 2004, Maritim et al., 2003 and Reuter et al., 2010). Indeed, many studies have found evidence of increased oxidative stress in MDD, as well as increased gene expression regulated by oxidative stress, suggesting that this is an important contributor to depression pathophysiology (Lindqvist et al., 2017, Maes et al., 2011, Mellon et al., 2016 and Wolkowitz et al., 2011).

Oxidative stress occurs when an organism is not able to effectively detoxify or repair the harmful effects of free radicals, for example, when pro-oxidant species (e.g., F2-isoprostanes and oxidized glutathione) exceed the capacity of anti-oxidant defenses (e.g., reduced glutathione; Coyle and Puttfarcken, 1993). Oxidative stress has detrimental effects on cellular membranes, proteins, nucleic acids, and lipids. Although nearly all tissues are sensitive to oxidative damage, the brain is especially sensitive, since it is a major consumer of oxygen and is rich in oxidizable lipids (Halliwell, 2006). Therefore, oxidative stress can result in altered neuronal functioning, neurotransmission, and overall brain activity (Bakunina et al., 2015, Cardozo-Pelaez et al., 1999 and Wang et al., 2010).

Recent meta-analyses have concluded that peripheral oxidative stress markers are elevated in depression, while levels of antioxidants are decreased (Black et al., 2015 and Palta et al., 2014). Although MDD has been the most widely studied psychiatric illness in this regard, other psychiatric illnesses, including anxiety disorders, are also characterized by increased oxidative stress (Atmaca et al., 2008, Bouayed et al., 2009, Kuloglu et al., 2002 and Ozdemir et al., 2009), raising the possibility that oxidative stress may be a non-specific marker of psychiatric illness or that it may relate to certain trans-diagnostic dimensional symptoms, as opposed to specific syndromal diagnoses *per se*.

Depression and anxiety disorders are highly co-morbid, and there is a significant diagnostic overlap (Zbozinek et al., 2012). In fact, the majority of individuals with MDD suffer from a co-morbid anxiety disorder or subsyndromal anxiety symptoms (Zimmerman et al., 2000), often associated with poorer disease outcomes, including higher rates of suicide, more functional impairments, and worse response to antidepressant treatment (Belzer and

Schneier, 2004). Although there are certain symptoms considered to be “core” features of anxiety and depressive disorders, there are also several shared symptoms, such as fatigue or loss of energy, difficulty concentrating, and sleep disturbances, leading to overlapping diagnostic boundaries (Zbozinek et al., 2012).

Most studies examining oxidative stress in MDD have focused on syndromally-defined MDD, and few, if any, have mapped oxidative stress onto specific symptom profiles, including symptoms of anxiety within MDD (Chung et al., 2012 and Irie et al., 2005). Therefore, it is currently unclear whether the observed increase in oxidative stress in MDD is driven by symptoms of depression, anxiety, or both. It is important to better understand the psychiatric correlates of oxidative stress, as this could facilitate the development of novel anti-oxidant treatments in selected subjects with MDD.

F2-isoprostanes and oxidized and reduced glutathione (GSSG and GSH, respectively) are among the most studied oxidative stress markers in MDD. F2-isoprostanes are a highly accurate measure of lipid peroxidation and are considered a robust marker of oxidative stress (Milne et al., 2005 and Patrignani and Tacconelli, 2005). Lipid peroxidation leads to impaired cell functioning, loss of membrane elasticity and fluidity, and potentially cell death (Montuschi et al., 2004). GSH is a nonenzymatic antioxidant and is oxidized by reactive oxygen species to form GSSG (Schulz et al., 2000). GSH levels, therefore, are considered as an indication of antioxidant capacity, whereas GSSG is an indication of oxidative stress. The GSH system, one of the major antioxidant systems of the brain (Dringen, 2000), is essential for cell proliferation (Poot et al., 1995) and is involved in the regulation of apoptosis (Hall, 1999). In fact, accumulation of GSSG levels within cells has been found to directly trigger (neural) cell death (Filomeni et al., 2003 and Park et al., 2009).

In the current study, we investigated the relationship between anxiety symptoms, depression symptoms, and the peripheral oxidative stress markers F2-isoprostanes and GSSG, as well as the antioxidant GSH, in unmedicated individuals with MDD. Based on previous findings of increased oxidative stress and reduced antioxidant capacity in MDD and anxiety disorders (Black et al., 2015, Ng et al., 2008, Sarandol et al., 2007 and Smaga et al., 2015), we hypothesized that more severe anxiety and depression symptoms would be related to increased levels of F2-isoprostanes and GSSG, and decreased levels of GSH.

2. Methods

2.1 Subjects

The study was approved by the University of California, San Francisco (UCSF) Committee on Human Research. All subjects gave written informed consent to participate. Subjects were recruited by clinical referrals, newspaper advertisement, flyers, bulletin board notices, and Craigslist postings.

We enrolled 69 subjects with unipolar MDD who were medication-free for at least six weeks. Oxidative stress data from subsets of this sample have been previously reported in relation to hypotheses not explored in the present study (Lindvist et al., 2017; Rawdin et al., 2013). MDD diagnoses were determined using the Structured Clinical Interview for the

DSM-IV (First et al., 1997). Subjects were required to have a minimum score of 17 on the 17-item Hamilton Depression Rating Scale (HAM-D; Hamilton, 1960).

Subjects with current psychosis or a history of bipolar disorder or psychotic symptoms outside of a major depressive episode were excluded. In addition, subjects were excluded if they had an eating disorder or post-traumatic stress disorder (PTSD) within one month of entering the study, or alcohol or substance dependence or abuse within six months of entering the study. Depressed subjects with co-morbid anxiety disorders were allowed (except PTSD) if the primary diagnosis was MDD.

The presence of medical conditions was assessed by medical history, physical examinations, and routine blood screening. Subjects were excluded if they had acute illnesses or infections, neurological disorders, chronic inflammatory disorders, or other major medical conditions that were considered potentially confounding.

All subjects were free of any psychotropic medication, steroid-containing contraceptives, hormone supplements, and any other potentially confounding medications for at least six weeks prior to participation, and none were taking vitamin supplements above the United States Recommended Daily Allowances. Subjects were allowed to use short-acting sedative hypnotics for sleep up to three times a week, but no sedative hypnotics were allowed within one week prior to entering the study.

2.2 Procedures

After having fasted (except water) since 2200 hours the night before, subjects were admitted as outpatients to the UCSF Clinical and Translational Science Institute between 0800 and 1100 hours. First, subjects were required to test negative on a urine toxicology screening and, if applicable, on a urine pregnancy test. Blood was drawn after 25 – 45 minutes of relaxation. Plasma for oxidative stress markers was collected in lavender-top EDTA tubes and was placed on ice immediately after being drawn. Plasma was separated by centrifugation at 14000 rpm at 4°C for 15 minutes in a refrigerated centrifuge within 15-20 minutes after the blood draw.

2.3 Assay of oxidative stress markers

Due to differing recruitment periods (2006 – 2008 and 2011 – 2014), samples were assayed in separate batches. As noted below (Section 2.5), this factor was statistically corrected for in all analyses.

F2-isoprostanes were quantified at Vanderbilt University, Eicosanoid Laboratory. The F2-isoprostanes were extracted and purified using solid-phase extraction and thin layer liquid chromatography. Subsequently, they were converted into trimethyl-ether derivatives and analyzed using chromatography– mass spectrometry (GC–MS), which is described elsewhere (Milne et al., 2007 and Morrow and Roberts, 2002).

For GSSG and GSH assessment, the use of different assay batches necessitated a change in assay laboratories and methods over the course of the study.

The first batch consisted of samples from 16 MDD subjects. Samples were processed by adding 400 μ L ice cold 5% meta-phosphoric acid and 100 μ L of whole blood into a 1.5mL microcentrifuge tube. Immediately, the microcentrifuge tube was capped and vortexed for 5 seconds and then centrifuged at 4°C for 2 minutes at 13000 RPM in a microcentrifuge. The supernatant was aliquoted to a vial and was stored at -80°C until assay at the Kronos Science Laboratory. The measurements were carried out using LC-MS/MS on a system that consisted of two Shimadzu LC-10AD pumps, a Shimadzu degasser (Shimadzu Scientific Instruments, Columbia, MD), and a Perkin Elmer autosampler (Perkin Elmer LLC, Norwalk, CT) directly interfaced with a triple-stage quadrupole mass spectrometer (API2000, Applied Biosystems, Foster City, CA) equipped with a TurboIonSpray ionization source. Nitrogen was used as the collision gas. Glutathione under MRM mode was measured using Positive ionization methods. GSSG and GSH were monitored with ion pairs (m/z) of 613/355 and 308/179, respectively. A seven-point linear calibration curve was established for each analyte using internal and external standards. Three quality-control samples were injected at the beginning, the end, and after every 10 samples to monitor intra- and inter-day assay accuracy and precision. Recoveries were consistently in excess of 90% and coefficients of variance ranged from 3% to 10%.

The second and third batches of GSSG and GSH consisted of samples from 48 MDD subjects. Samples were processed by adding 200 μ L ice cold 10% meta-phosphoric acid to 200 μ L of plasma, after which the sample was incubated for 30 minutes in order to precipitate proteins. The sample was centrifuged at 14,000 rpm for 15 min at 4°C, after which the supernatant was filtered through a 0.2 mm filter (PGC Scientific, Frederic, MD). A 20 μ L aliquot was injected in the HPLC-electrochemical detection system and the ESA autosampler (model 542). HPLC with an ESA solvent delivery system (model 580) and a reverse phase Capcell pak C18 column (3 mm; 4.6 $\times$ 150 mm, Phenomenex, Inc., Torrance, CA, USA) were used for the analyses. A 16 channel CouloArray electrochemical detector (ESA, Inc., Chelmsford, MA) with a high sensitive four-sensor cell (model 6210) was used to determine the plasma metabolites. Using HPLC software, the concentrations of the plasma metabolites were calculated from peak areas and standard calibration curves included with each run. The coefficients of variation were 5% and 8% for GSH and GSSG, respectively.

2.4 Anxiety and depression measures

The 17-item HAM-D was used to determine the severity of depression (Hamilton, 1960). To assess severity of anxiety, we used the Hamilton Anxiety Rating Scale (HAM-A; Hamilton, 1959). The HAM-A is a widely used clinician-rated scale consisting of 14 items that are each scored on a scale of 0 (not present) to 4 (severe). The HAM-D and HAM-A have good reliability (Bagby et al., 2004) and validity (Cusin et al., 2009 and Maier et al., 1988).

It is commonly acknowledged that the HAM-A and HAM-D assess certain overlapping symptoms of anxiety and depression, making it difficult to parse out the differences between anxiety and depression (Zbozinek, et al., 2012). To address this, we used subscales of the HAM-A and HAM-D to investigate the relationship between oxidative stress and “core” symptoms of anxiety and depression. To measure core anxiety, we used the Hamilton Core

Anxiety Scale (HAM-A₆; Bech, 2006, Bech et al., 2005 and Kent et al., 2016). The HAM-A₆ includes the items anxious mood, tension, fears, intellectual, somatic (muscular), and anxious behavior at interview. To measure core depression, we used the Core Depression Factor Subscale of the HAM-D (Bech, 2006, Detke et al., 2002 and Perahia et al., 2006). We specifically chose this subscale because it does not include the item “psychic anxiety” that is frequently present in other subscales of core depression. The core depression factor subscale includes the HAM-D items depressed mood, feelings of guilt, suicide, work and activities, and psychomotor retardation (Bech, 2006).

Another approach to capture specific symptomatology is to select a single item that best reflects the primary state of the pathology. We selected the HAM-D item “psychic anxiety” to capture the emotion of “pure” anxiety (Russel et al., 2007 and Thase et al., 2014). We classified subjects into one of two subgroups, according to their scores on the single item psychic anxiety: absent to mild anxiety (score 0 or 1) or moderate to severe anxiety (score 2, 3, or 4; Russel et al., 2007). To capture the emotion of “pure” depression, we selected the HAM-D item “depressed mood” (Hieronimus et al., 2016). Subjects were classified into one of two subgroups according to their score on the single item depressed mood: low to mild depressed mood (score 1 or 2) or moderate to severe depressed mood (score 3 or 4).

2.5 Data analysis

Statistical analyses were performed with IBM SPSS Statistics, version 23. All tests were 2-tailed with an alpha=0.05. Since oxidative stress markers were analyzed in separate assay batches (two for F2-isoprostanes and three for GSH and GSSG), we transformed biomarker levels into z-scores before combining data across batches. We examined the distributions for normality, and non-normal distributions were either natural log transformed (HAM-A and HAM-D) or Blom transformed in cases where log transformation was not feasible (F2-isoprostanes and GSH; Blom, 1958).

There were missing values for both HAM-A (16% missing data) and oxidative stress markers (6-7% missing data), due to clinical constraints and difficulties while obtaining blood samples, respectively. These randomly missing data were handled in two ways: First, multiple imputation was used, employing the fully conditional approach (FCS) and predictive mean matching (PMM), with 100 maximum iterations and 20 imputed data sets (as described by Rubin, 1986, 1987). Second, we performed a sensitivity analysis using the original data, in which subjects with missing data were excluded.

To investigate the relationship between symptom scores (independent variable) and oxidative stress markers (dependent variable), we performed a series of linear regression analyses, one model for each pair of symptom scores and oxidative stress markers. We used residual values of the oxidative stress markers that were adjusted for age, sex, BMI, and tobacco use (Black et al., 2016a and Block et al., 2002).

We performed independent t-tests to investigate whether the subgroups low vs. high anxiety and low vs. high depression (single-item measures) differed in levels of oxidative stress markers (using residual values corrected for age, sex, BMI, and tobacco use).

3. Results

3.1 Demographics

Table 1 displays the characteristics of the study sample. HAM-D scores ranged from 17 (moderate depression) to 34 (very severe depression), with an average score of 20, indicating that overall, our sample was moderately to severely depressed. HAM-A scores ranged from 8 (mild anxiety) to 33 (severe anxiety), with an average score of 17, indicating a mild to moderate overall anxiety severity in our sample.

3.2 Hamilton Anxiety and Depression Rating Scales

Figure 1 shows the relationship between HAM-A ratings (panel A and C), HAM-D ratings (panel B and D), and the oxidative stress markers F2-isoprostanes and GSSG. As hypothesized, HAM-A ratings positively predicted levels of F2-isoprostanes ($\beta = .26$, 95% CI [.010, .506], $p = .042$) and levels of GSSG ($\beta = .25$, 95% CI [.003, .487], $p = .049$). HAM-A ratings did not, however, significantly predict GSH levels ($\beta = .05$, 95% CI [-.205, .300], $p = .711$).

HAM-D ratings did not significantly predict levels of F2-isoprostanes ($\beta = .13$, 95% CI [-.123, .374], $p = .323$), GSSG ($\beta = .13$, 95% CI [-.118, .369], $p = .314$), or GSH ($\beta = -.12$, 95% CI [-.357, .126], $p = .347$).

3.3 Core anxiety and core depression

We next sought to determine if higher “core” anxiety severity was related to increased oxidative stress. Core anxiety severity positively predicted levels of F2-isoprostanes ($\beta = .34$, 95% CI [.081, .606], $p = .012$). Core anxiety severity also positively predicted levels of GSSG, but this did not reach significance ($\beta = .24$, 95% CI [-.024, .506], $p = .074$). Further, we found no significant correlation between core anxiety severity and levels of GSH ($\beta = .04$, 95% CI [-.257, .327], $p = .803$).

Core depression severity did not significantly predict any of the oxidative stress markers; F2-isoprostanes ($\beta = .19$, 95% CI [-.054, .425], $p = .130$), GSSG ($\beta = .18$, 95% CI [-.053, .422], $p = .128$), and GSH ($\beta = .08$, 95% CI [-.165, .315], $p = .540$).

3.4 Low vs. high anxiety and low vs. high depression (single-item analysis)

Oxidative stress markers in individuals with high vs. low “psychic anxiety” and high vs. low “depressed mood” items are summarized in Table 2. We found significantly higher levels of F2-isoprostanes and GSSG in the high anxiety subgroup compared to the low anxiety subgroup, with medium effect sizes (Cohen's $d = 0.54$ and 0.62 , respectively; Cohen, 1988). There were no significant differences in GSH levels between subjects with high anxiety compared to subjects with low anxiety. With regard to depression severity, we found no significant differences in F2-isoprostanes, GSSG, and GSH between the low and high depression subgroups.

The sensitivity analyses (in which cases with missing data were excluded) resulted in nearly identical findings, with only a small difference in significance values for the relationship

between GSSG and HAM-A ratings ($p = .049$ for multiply imputed data, and $p = .059$ for original, non-imputed data), and the relationship between GSSG and core anxiety ($p = .074$ for multiply imputed data, and $p = .030$ for original, non-imputed data).

4. Discussion

In the present study, we examined the relationship between oxidative stress, anxiety, and depression symptoms in somatically healthy, medication-free individuals with MDD. Our main finding was that MDD subjects with more severe anxiety, but not subjects with more severe depression, showed increased oxidative stress, after controlling for age, sex, BMI, and smoking.

Specifically, we found that higher plasma levels of F2-isoprostanes and GSSG were associated with higher total HAM-A scores, higher core anxiety scores (HAM-A₆), and higher scores on the single HAM-D item “psychic anxiety.” The group differences in oxidative stress markers between those with low vs. high anxiety were of medium effect size (Cohen's $d = 0.54$ and 0.62). Similarly, the continuous relationships between severity of anxiety symptoms and oxidative stress marker levels showed r values ranging from .24 to .34, indicating small to medium effect sizes (Cohen, 1988). Surprisingly, F2-isoprostanes and GSSG levels showed no significant associations with depression severity, as quantified by total HAM-D scores, core depression scores, and scores on the single HAM-D item “depressed mood”. As opposed to the oxidative stress markers, the antioxidant GSH was not significantly associated with either anxiety or depression severity.

To the best of our knowledge, this is the first study to show a relationship between increased oxidative stress and more severe symptoms of anxiety, but not depression, in medication-free individuals with MDD. This conclusion is supported by the concordance of our findings across three different characterizations of anxiety and depression symptoms. Our findings raise the possibility that the increased oxidative stress reported in many studies of MDD may be more attributable to anxiety rather than to depression *per se*, although this needs to be confirmed in future studies. Our findings may also point to the possibility that increased oxidative stress contributes to, or is a consequence of, the worse illness course seen in MDD patients with co-morbid anxiety compared to those with more “pure” depressive symptoms (Belzer and Schneier, 2004).

Peripheral markers of increased oxidative stress, as well as decreased antioxidants, have been reported in individuals with MDD (Black et al., 2015, Lindqvist et al., 2017 and Palta et al., 2014) and in individuals with anxiety disorders, including panic disorder, social phobia, and obsessive compulsive disorder (Atmaca et al., 2008, Kuloglu et al., 2002 and Ozdemir et al., 2009). To our knowledge, only two prior studies examined the relationship between oxidative stress and specific symptom dimensions of MDD. Chung et al. (2013) examined the relationship between oxidative stress and the Profiles of Mood State (POMS) questionnaire in MDD subjects, but found no significant associations with any of the subscales. Irie et al. (2005) found similar results as in the present study, namely, a positive relationship between oxidative stress and core anxiety severity (“tension-anxiety” subscale of the POMS), but not core depression severity (“depression-rejection” subscale of POMS)

in female MDD subjects. The effect sizes of the relationship between anxiety and oxidative stress reported by Irie et al. (2005) were medium to large in size, which are larger in magnitude than the effect sizes found in this study. However, the results of the present study were based on a substantially larger sample of depressed individuals, all of whom were medication-free, and analyses controlled for potential confounders, in contrast to the two prior studies.

A dimensional approach in psychiatric research has been recently advocated for, exemplified by the introduction of the RDoC criteria by the NIMH (Cuthbert and Insel, 2013). In this paradigm, anxiety symptoms are part of the “Threat” construct in the Negative Valence Systems domain, and depression symptoms are part of the “Loss” construct and “Reward” constructs in the Negative and Positive Valence Systems domains, respectively (Woody and Gibb, 2015). Although the present study did not employ an RDoC design, it did attempt to parse out anxiety vs. depression components in relation to oxidative stress.

Both preclinical (Bouayed et al., 2007, Rammal et al., 2008 and Vollert et al., 2011) and clinical studies (Atmaca et al., 2008 and Kologlu et al., 2002) have found a positive relationship between oxidative stress and severity of anxiety symptoms. Interestingly, several animal studies have shown that *induction* of oxidative stress results in anxiety-like behavior (Masood et al., 2008 and Salim et al., 2010). Moreover, a genomics study showed that overexpression of genes involved in oxidative stress metabolism (glutathione reductase 1 and glyoxalase 1) in the mouse brain increased anxiety-like behavior, whereas inhibition of glyoxalase 1 expression had the opposite effect (Hovatta et al., 2005). In contrast, we are not aware of animal studies that have examined causality of oxidative stress in depression-like behaviors (Salim, 2014).

Whether oxidative stress is the cause or consequence of anxiety, or if the two are jointly related to a third factor (e.g., genetics or lifestyle factors; Hovatta et al., 2010), is still unknown. Nonetheless, the hypothesis that oxidative stress is involved in the genesis of anxiety has been widely proposed (Bouayed et al., 2009 and Hassan et al., 2014). The neurobiological mechanisms may involve brain regions such as the amygdala and the hippocampus, which are especially sensitive to oxidative damage (Masood et al., 2008, Salim, 2016 and Salim et al., 2010) and which are implicated in anxiety-related disorders (Shin and Liberzon, 2010). Oxidative stress may lead to neuroendocrine alterations in the amygdala, including dendritic shrinkage and amygdala hyperactivity (Salim, 2016). Moreover, oxidative damage to the hippocampus might disrupt the neural circuitry involved in anxiety and fear conditioning (Hassan et al., 2014). Additional mechanisms proposed to mediate the oxidative stress/anxiety relationship include impaired secretion of brain-derived neurotrophic factor (BDNF; Salim et al., 2011) and pathologic alterations in NMDA receptor activity, mitochondrial functioning, and neurogenesis (Haxaire et al., 2012, Hovatta et al., 2010 and Rai et al., 2013). However, any inferences about central nervous system effects, based on our data, are speculative since we only studied peripheral oxidative stress markers. Further, none of these explanations would predict a relationship exclusively with anxiety without also predicting a relationship with other psychiatric symptoms such as depression.

Our findings are in contrast to some previous studies showing a positive relationship between oxidative stress and overall depression severity (Bilici et al., 2001, Dimopoulos et al., 2008, Forlenza and Miller, 2006, Irie et al., 2005, Jorgensen et al., 2013, Sarandol et al., 2007 and Yanik et al., 2004). The discrepant findings may be explained by the large heterogeneity of oxidative stress measures across studies. In fact, two out of three previous MDD studies that examined the relationship between F2-isoprostanes and depression severity were, as the present study, negative (Chung et al., 2012 and Yager et al., 2010). Further, any differences between our study and others could be explained by our stringent inclusion and exclusion criteria, resulting in a sample of somatically healthy and medication-free individuals. This, and our decision to adjust for the potentially confounding influences of age, sex, smoking, and BMI, are in contrast to most prior studies examining the relationship between oxidative stress and depression symptom severity (Bilici et al., 2001, Dimopoulos et al., 2008, Chung et al., 2012, Irie et al., 2005, Jorgensen et al., 2013, Sarandol et al., 2007 and Yanik et al., 2004).

The finding that GSH was not related to either anxiety or depression severity is in contrast to our hypothesis that decreased antioxidant activity would be related to more severe symptoms. However, our finding is in line with a previous report from our group that GSH levels were not decreased in MDD subjects compared to healthy controls, despite the fact that F2-isoprostanes were elevated (Lindqvist et al., 2017). Our findings may indicate that chronic oxidative stress results in a compensatory but inadequate increase in GSH synthesis in attempt to reduce further oxidative damage (Andreazza et al., 2009), but this is purely hypothetical.

Our study has several limitations. Firstly, we assessed oxidative stress markers at only one time point on a single day. F2-isoprostanes are chemically stable and day-to-day variability, as well as diurnal variability, are relatively limited (Montuschi et al., 2004). GSSG levels also have a minimal diurnal variation, but, on the other hand, GSH levels show fluctuations over the day, especially related to food intake (Blanco et al., 2007). This might be another possible reason for not finding correlations between symptom severity and GSH levels. However, we obtained blood samples between 8 and 11 AM in all subjects to minimize diurnal variability, and subjects had fasted since the night before to diminish the effect of food intake on plasma oxidative stress levels. Secondly, we only assessed the oxidative markers in peripheral blood samples, therefore extrapolations to brain oxidative stress would be purely conjectural. However, peripheral markers of oxidative stress have recently been correlated with decreased hippocampal CA3/dentate gyrus volume in unmedicated individuals with MDD (Lindqvist et al., 2014). Thirdly, although we controlled for the potential confounding effects of age, sex, BMI, and tobacco use, we cannot rule out that other variables, not accounted for in our analyses, may influence oxidative stress in MDD. Dietary patterns have recently been identified as potential confounders in the association between oxidative stress and depression (Black et al., 2016c). Future studies should include dietary measures to further examine the role of this factor in the relationship between MDD and oxidative stress. Fourthly, the relationship between oxidative stress and non-pathological levels of anxiety and depression, such as in healthy controls, was not addressed in this study. Lastly, due to our strict inclusion and exclusion criteria, the findings may not be generalizable to real-world MDD patients seen in clinical practice.

Strengths of the study are that we used somatically healthy depressed patients who had all been free of antidepressants and other possibly confounding drugs for a minimum of six weeks. The use of antidepressants may influence oxidative stress markers and may thus be a serious confounder (Black et al., 2016b and Mellon et al., 2016). Another strength is that we used several strategies to parse out anxiety and depression symptomatology in relation to oxidative stress. The congruence of findings among these approaches strengthens the overall conclusions and provides face validity to our results.

To conclude, the present study showed a relationship between increased peripheral oxidative stress and more severe anxiety, but not depression symptoms, in physically healthy, medication-free individuals with MDD. Our findings highlight the importance of mapping oxidative stress, and perhaps other biomarkers, onto specific dimensional symptoms in addition to syndromal diagnoses. Future studies should expand the range of symptom dimensions surveyed (i.e., using the RDoC) and the number and type of oxidative stress measures studied. Moreover, future studies should test whether successful treatment of underlying symptoms diminishes oxidative stress. Although not addressed in the current study, it might be possible that antioxidant treatments could play a role in ameliorating anxiety symptoms in individuals with MDD (Balmus et al., 2016 and Deepmala et al., 2015). Also, to the extent anxiety symptoms are directly correlated with oxidative stress, it will be important to determine whether a causal relationship exists, and, if so, its directionality. Distinguishing these possibilities could lead to a better recognition of the biological underpinnings of affective disorders.

Acknowledgments

The authors gratefully acknowledge the assistance of Kevin Delucchi, PhD, Phuong Hoang, and Alanie Lazaro (all at UCSF), the nursing and other staff of UCSF CTSI Clinical Research Center, and all past and present UCSF PNE Lab volunteer research assistants, as well as all of the participants in this research.

Funding: This study was funded by grants from the National Institute of Mental Health (NIMH; grant No. R01-MH083784), the Tinberg family, the O'Shaughnessy Foundation, the UCSF Academic Senate and the UCSF Research Evaluation and Allocation Committee (REAC). This project was also supported by the National Institutes of Health/National Center for Research Resources (NIH/NCRR) and the National Center for Advancing Translational Sciences, NIH (through UCSF-CTSI grant No. UL1 RR024131). D.L. was supported by the Swedish Research Council (registration No. 2015-00387), Marie Skłodowska-Curie Actions, COFUND, (Project INCA 600398), the Söderström-Königska Foundation, the Sjöbring Foundation, and the province of Scania (Sweden) state grants (ALF).

References

- Andreazza AC, Kapczinski F, Kauer-Sant'Anna M, Walz JC, Bond DJ, Gonçalves CA, Young LT, Yatham LN. 3-Nitrotyrosine and glutathione antioxidant system in patients in the early and late stages of bipolar disorder. *J Psychiatry Neurosci*. 2009; 34:263–271. [PubMed: 19568477]
- Atmaca M, Kuloglu M, Tezca E, Ustundag B. Antioxidant enzyme and malondialdehyde levels in patients with social phobia. *Psychiatry Research*. 2008; 159:95–100. [PubMed: 18339429]
- Bagby RM, Ryder AG, Schuller DR, Marshall MB. The Hamilton Depression Rating Scale: has the gold standard become a lead weight? *Am J Psychiatry*. 2004; 161:2163–2177. [PubMed: 15569884]
- Bakunina N, Pariante CM, Zunszain PA. Immune mechanisms linked to depression via oxidative stress and neuroprogression. *Immunology*. 2015; 144:365–373.
- Bech P. Rating scales in depression: limitations and pitfalls. *Dialogues Clin Neurosci*. 2006; 8:207–215. [PubMed: 16889106]

- Bech P, Lunde M, Uden M. An inventory for the measurement of generalised anxiety distress symptoms, the GAD-10 Inventory. *Acta Psychiatrica Belgica*. 2005; 105:111–118.
- Belzer K, Schneier FR. Comorbidity of anxiety and depressive disorders: issues in conceptualization, assessment, and treatment. *J Psychiatr Pract*. 2004; 10:296–306. [PubMed: 15361744]
- Bilici M, Efe H, Köro lu MA, Uydu HA, Bekaro lu M, De er O. Antioxidative enzyme activities and lipid peroxidation in major depression: alterations by antidepressant treatments. *J Affect Disord*. 2001; 64:43–51. [PubMed: 11292519]
- Black CN, Bot M, Scheffer PG, Cuijpers P, Penninx BW. Is depression associated with increased oxidative stress? A systematic review and meta-analysis. *Psychoneuroendocrinology*. 2015; 51:164–175. [PubMed: 25462890]
- Black CN, Bot M, Scheffer PG, Penninx BW. Oxidative stress in major depressive and anxiety disorders, and the association with antidepressant use; results from a large adult cohort. *Psychol Med*. 2016a:1–13.
- Black CN, Bot M, Scheffer PG, Penninx BW. Sociodemographic and Lifestyle Determinants of Plasma Oxidative Stress Markers 8-OHdG and F2-Isoprostanes and Associations with Metabolic Syndrome. *Oxid Med Cell Longev*. 2016b; 2016:7530820. [PubMed: 27006748]
- Black CN, Penninx BW, Bot M, Odegaard AO, Gross MD, Matthews KA, Jacobs DR. Oxidative stress, anti-oxidants and the cross-sectional and longitudinal association with depressive symptoms: results from the CARDIA study. *Transl Psychiatry*. 2016c; 6:e743. [PubMed: 26905415]
- Blanco RA, Ziegler TR, Carlson BA, Cheng PY, Park Y, Cotsonis GA, Accardi CJ, Jones DP. Diurnal variation in glutathione and cysteine redox states in human plasma. *Am J Clin Nutr*. 2007; 86:1016–1023. [PubMed: 17921379]
- Blom, G. Statistical estimates and transformed beta-variables. J. Wiley & Sons; Almqvist & Wiksell; New York; Stockholm: 1958.
- Bouayed J, Rammal H, Soulimani R. Oxidative stress and anxiety: relationship and cellular pathways. *Oxid Med Cell Longev*. 2009; 2:63–67. [PubMed: 20357926]
- Cardozo-Pelaez F, Song S, Parthasarathy A, Hazzi C, Naidu K, Sanchez-Ramos J. Oxidative DNA damage in the aging mouse brain. *Mov Disord*. 1999; 14:972–980. [PubMed: 10584672]
- Ceriello A, Motz E. Is oxidative stress the pathogenic mechanism underlying insulin resistance, diabetes, and cardiovascular disease? The common soil hypothesis revisited. *Arterioscler Thromb Vasc Biol*. 2004; 24:816–823. [PubMed: 14976002]
- Chung CP, Schmidt D, Stein CM, Morrow JD, Salomon RM. Increased oxidative stress in patients with depression and its relationship to treatment. *Psychiatry Res*. 2013; 206:213–216. [PubMed: 23245537]
- Coyle JT, Puttfarcken P. Oxidative stress, glutamate, and neurodegenerative disorders. *Science*. 1993; 262:689–695. [PubMed: 7901908]
- Cusin, C., Yang, H., Yeung, A., Fava, M. Rating scales for depression. In: Baer, L., Blais, MA., editors. *Handbook of clinical rating scales and assessment in psychiatry and mental health*. Humana Press; New York: 2009. p. 7-35.
- Cuthbert BN, Insel TR. Toward the future of psychiatric diagnosis: the seven pillars of RDoC. *BMC Med*. 2013; 11:126. [PubMed: 23672542]
- Detke MJ, Lu Y, Goldstein DJ, McNamara RK, Demitrack MA. Duloxetine 60 mg once daily dosing versus placebo in the acute treatment of major depression. *J Psychiatr Res*. 2002; 36:383–390. [PubMed: 12393307]
- Dimopoulos N, Piperi C, Psarra V, Lea RW, Kalofoutis A. Increased plasma levels of 8-iso-PGF 2 α and IL-6 in an elderly population with depression. *Psychiatry Research*. 2008; 16:59–66.
- Emerit J, Edeas M, Bricaire F. Neurodegenerative diseases and oxidative stress. *Biomed Pharmacother*. 2004; 58:39–46. [PubMed: 14739060]
- Filomeni G, Rotilio G, Ciriolo MR. Glutathione disulfide induces apoptosis in U937 cells by a redox-mediated p38 MAP kinase pathway. *FASEB J*. 2003; 17:64–66. [PubMed: 12424221]
- First, MB., Spitzer, RL., Gibbon, M., Williams, JB. User's guide for the Structured clinical interview for DSM-IV axis I disorders SCID-I: clinician version. American Psychiatric Pub; 1997.

- Forlenza MJ, Miller GE. Increased serum levels of 8-hydroxy-2'-deoxyguanosine in clinical depression. *Psychosom Med*. 2006; 68:1–7. [PubMed: 16449405]
- Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, Nakajima Y, Nakayama O, Makishima M, Matsuda M, Shimomura I. Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest*. 2004; 114:1752–1761. [PubMed: 15599400]
- Halliwell B. Reactive species and antioxidants. Redox biology is a fundamental theme of aerobic life. *Plant Physiol*. 2006; 141:312–322. [PubMed: 16760481]
- Hamilton M. The assessment of anxiety states by rating. *Br J Med Psychol*. 1959; 32:50–55. [PubMed: 13638508]
- Hamilton M. A rating scale for depression. *J Neurol Neurosurg Psychiatry*. 1960; 23:56–62. [PubMed: 14399272]
- Hassan W, Silva CE, Mohammadzai IU, da Rocha JB, J LF. Association of oxidative stress to the genesis of anxiety: implications for possible therapeutic interventions. *Curr Neuropharmacol*. 2014; 12:120–139. [PubMed: 24669207]
- Haxaire C, Turpin FR, Potier B, Kervern M, Sinet PM, Barbanel G, Mothet JP, Dutar P, Billard JM. Reversal of age-related oxidative stress prevents hippocampal synaptic plasticity deficits by protecting D-serine-dependent NMDA receptor activation. *Aging Cell*. 2012; 11:336–344. [PubMed: 22230264]
- Hieronymus F, Emilsson JF, Nilsson S, Eriksson E. Consistent superiority of selective serotonin reuptake inhibitors over placebo in reducing depressed mood in patients with major depression. *Mol Psychiatry*. 2016; 21:523–530. [PubMed: 25917369]
- Hovatta I, Juhila J, Donner J. Oxidative stress in anxiety and comorbid disorders. *Neurosci Res*. 2010; 68:261–275. [PubMed: 20804792]
- Hovatta I, Tennant RS, Helton R, Marr RA, Singer O, Redwine JM, Ellison JA, Schadt EE, Verma IM, Lockhart DJ, Barlow C. Glyoxalase 1 and glutathione reductase 1 regulate anxiety in mice. *Nature*. 2005; 438:662–666. [PubMed: 16244648]
- Irie M, Miyata M, Kasai H. Depression and possible cancer risk due to oxidative DNA damage. *J Psychiatr Res*. 2005; 39:553–560. [PubMed: 16005897]
- Jones DP. Redox potential of GSH/GSSG couple: assay and biological significance. *Methods Enzymol*. 2002; 348:93–112. [PubMed: 11885298]
- Jorgensen A, Krogh J, Miskowiak K, Bolwig TG, Kessing LV, Fink-Jensen A, Nordentoft M, Henriksen T, Weimann A, Poulsen HE, Jorgensen MB. Systemic oxidatively generated DNA/RNA damage in clinical depression: associations to symptom severity and response to electroconvulsive therapy. *J Affect Disord*. 2013; 149:355–362. [PubMed: 23497793]
- Kent JM, Daly E, Kezic I, Lane R, Lim P, De Smedt H, De Boer P, Van Nueten L, Drevets WC, Ceusters M. Efficacy and safety of an adjunctive mGlu2 receptor positive allosteric modulator to a SSRI/SNRI in anxious depression. *Prog Neuropsychopharmacol Biol Psychiatry*. 2016; 67:66–73. [PubMed: 26804646]
- Kuloglu M, Atmaca M, Tezcan E, Ustundag B, Bulut S. Antioxidant enzyme and malondialdehyde levels in patients with panic disorder. *Neuropsychobiology*. 2002a; 46:186–189. [PubMed: 12566935]
- Kuloglu M, Atmaca M, Tezcan E, Ustundag B, Bulut S. Antioxidant enzyme and malondialdehyde levels in patients with panic disorder. *Neuropsychobiology*. 2002b; 46:186–189. [PubMed: 12566935]
- Lindqvist D, Dhabhar FS, James SJ, Hough CM, Jain FA, Bersani FS, Reus VI, Verhoeven JE, Epel ES, Mahan L, Rosser R, Wolkowitz OM, Mellon SH. Oxidative stress, inflammation and treatment response in major depression. *Psychoneuroendocrinology*. 2017; 76:197–205. [PubMed: 27960139]
- Lindqvist D, Mueller S, Mellon SH, Su Y, Epel ES, Reus VI, Rosser R, Mahan L, Mackin RS, Yang TT, Wolkowitz OM. Peripheral antioxidant markers are associated with total hippocampal and CA3/dentate gyrus volume in MDD and healthy controls-preliminary findings. *Psychiatry Res*. 2014; 224:168–174. [PubMed: 25266915]
- Maes M, Galecki P, Chang YS, Berk M. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the (neuro) degenerative processes

- in that illness. *Prog Neuropsychopharmacol Biol Psychiatry*. 2011; 35:676–692. [PubMed: 20471444]
- Maier W, Buller R, Philipp M, Heuser I. The Hamilton Anxiety Scale: reliability, validity and sensitivity to change in anxiety and depressive disorders. *J Affect Disord*. 1988; 14:61–68. [PubMed: 2963053]
- Maritim AC, Sanders RA, Watkins JB. Diabetes, oxidative stress, and antioxidants: a review. *J Biochem Mol Toxicol*. 2003; 17:24–38. [PubMed: 12616644]
- Masood A, Nadeem A, Mustafa SJ, O'Donnell JM. Reversal of oxidative stress-induced anxiety by inhibition of phosphodiesterase-2 in mice. *J Pharmacol Exp Ther*. 2008; 326:369–379. [PubMed: 18456873]
- Mellon SH, Wolkowitz OM, Schonemann MD, Epel ES, Rosser R, Burke HB, Mahan L, Reus VI, Stamatiou D, Liew CC, Cole SW. Alterations in leukocyte transcriptional control pathway activity associated with major depressive disorder and antidepressant treatment. *Transl Psychiatry*. 2016; 6:e821. [PubMed: 27219347]
- Milne GL, Musiek ES, Morrow JD. F2-isoprostanes as markers of oxidative stress in vivo: an overview. *Biomarkers*. 2005; 10(Suppl 1):S10–23. [PubMed: 16298907]
- Ng F, Berk M, Dean O, Bush AI. Oxidative stress in psychiatric disorders: evidence base and therapeutic implications. *Int J Neuropsychopharmacol*. 2008; 11:851–876. [PubMed: 18205981]
- Ozdemir E, Cetinkaya S, Ersan S, Kucukosman S, Ersan EE. Serum selenium and plasma malondialdehyde levels and antioxidant enzyme activities in patients with obsessive-compulsive disorder. *Prog Neuropsychopharmacol Biol Psychiatry*. 2009; 33:62–65. [PubMed: 18957313]
- Palta P, Samuel LJ, Miller ER, Szanton SL. Depression and oxidative stress: results from a meta-analysis of observational studies. *Psychosom Med*. 2014; 76:12–19. [PubMed: 24336428]
- Park HA, Khanna S, Rink C, Gnyawali S, Roy S, Sen CK. Glutathione disulfide induces neural cell death via a 12-lipoxygenase pathway. *Cell Death Differ*. 2009; 16:1167–1179. [PubMed: 19373248]
- Patrignani P, Tacconelli S. Isoprostanes and other markers of peroxidation in atherosclerosis. *Biomarkers*. 2005; 10(Suppl 1):S24–29. [PubMed: 16298908]
- Penninx BW, Milaneschi Y, Lamers F, Vogelzangs N. Understanding the somatic consequences of depression: biological mechanisms and the role of depression symptom profile. *BMC Med*. 2013; 11:129. [PubMed: 23672628]
- Perahia DG, Wang F, Mallinckrodt CH, Walker DJ, Detke MJ. Duloxetine in the treatment of major depressive disorder: a placebo- and paroxetine-controlled trial. *Eur Psychiatry*. 2006; 21:367–378. [PubMed: 16697153]
- Rai S, Kamat PK, Nath C, Shukla R. A study on neuroinflammation and NMDA receptor function in STZ (ICV) induced memory impaired rats. *J Neuroimmunol*. 2013; 254:1–9. [PubMed: 23021418]
- Rammal H, Bouayed J, Younos C, Soulimani R. Evidence that oxidative stress is linked to anxiety-related behaviour in mice. *Brain Behav Immun*. 2008; 22:1156–1159. [PubMed: 18620042]
- Rawdin BJ, Mellon SH, Dhabhar FS, Epel ES, Puterman E, Su Y, Burke HM, Reus VI, Rosser R, Hamilton SP, Nelson JC, Wolkowitz OM. Dysregulated relationship of inflammation and oxidative stress in major depression. *Brain Behav Immun*. 2013; 31:143–152. [PubMed: 23201587]
- Reuter S, Gupta SC, Chaturvedi MM, Aggarwal BB. Oxidative stress, inflammation, and cancer: how are they linked? *Free Radic Biol Med*. 2010; 49:1603–1616. [PubMed: 20840865]
- Rubin, DB. Multiple imputation for nonresponse in surveys. Wiley; New York: 1987.
- Rubin DB. Statistical matching using file concatenation with adjusted weights and multiple imputations. *J Bus Econ Stat*. 1986; 4:87–94.
- Russell J, Raskin J, Wiltse C, Walker D, Brawman-Mintzer O. Efficacy and tolerability of duloxetine treatment in elderly patients with major depressive disorder and concurrent anxiety symptoms. *Psychiatry (Edmont)*. 2007; 4:33–45. [PubMed: 20711334]
- Salim S. Oxidative stress and psychological disorders. *Curr Neuropharmacol*. 2014; 12:140–147. [PubMed: 24669208]
- Salim S. Oxidative Stress and the Central Nervous System. *J Pharmacol Exp Ther*. 2016

- Salim S, Asghar M, Chugh G, Taneja M, Xia Z, Saha K. Oxidative stress: a potential recipe for anxiety, hypertension and insulin resistance. *Brain Res.* 2010; 1359:178–185. [PubMed: 20816762]
- Sarandol A, Sarandol E, Eker SS, Erdinc S, Vatansever E, Kirli S. Major depressive disorder is accompanied with oxidative stress: short-term antidepressant treatment does not alter oxidative-antioxidative systems. *Hum Psychopharmacol.* 2007; 22:67–73. [PubMed: 17299810]
- Schulz JB, Lindenau J, Seyfried J, Dichgans J. Glutathione, oxidative stress and neurodegeneration. *Eur J Biochem.* 2000; 267:4904–4911. [PubMed: 10931172]
- Shin LM, Liberzon I. The neurocircuitry of fear, stress, and anxiety disorders. *Neuropsychopharmacology.* 2010; 35:169–191. [PubMed: 19625997]
- Smaga I, Niedzielska E, Gawlik M, Moniczewski A, Krzek J, Przeglasiński E, Pera J, Filip M. Oxidative stress as an etiological factor and a potential treatment target of psychiatric disorders. Part 2. Depression, anxiety, schizophrenia and autism. *Pharmacol Rep.* 2015; 67:569–580. [PubMed: 25933971]
- Thase ME, Chen D, Edwards J, Ruth A. Efficacy of vilazodone on anxiety symptoms in patients with major depressive disorder. *Int Clin Psychopharmacol.* 2014; 29:351–356. [PubMed: 24978955]
- Vollert C, Zagaar M, Hovatta I, Taneja M, Vu A, Dao A, Levine A, Alkadhi K, Salim S. Exercise prevents sleep deprivation-associated anxiety-like behavior in rats: potential role of oxidative stress mechanisms. *Behav Brain Res.* 2011; 224:233–240. [PubMed: 21621560]
- Wang X, Michaelis EK. Selective neuronal vulnerability to oxidative stress in the brain. *Front Aging Neurosci.* 2010; 2:12. [PubMed: 20552050]
- Wolkowitz OM, Reus VI, Mellon SH. Of sound mind and body: depression, disease, and accelerated aging. *Dialogues Clin Neurosci.* 2011; 13:25–39. [PubMed: 21485744]
- Woody ML, Gibb BE. Integrating NIMH Research Domain Criteria (RDoC) into Depression Research. *Curr Opin Psychol.* 2015; 4:6–12. [PubMed: 25642446]
- Yager S, Forlenza MJ, Miller GE. Depression and oxidative damage to lipids. *Psychoneuroendocrinology.* 2010; 35:1356–1362. [PubMed: 20417039]
- Yanik M, Erel O, Kati M. The relationship between potency of oxidative stress and severity of depression. *Acta Neuropsychiatr.* 2004; 16:200–203. [PubMed: 26984307]
- Zbozinek TD, Rose RD, Wolitzky-Taylor KB, Sherbourne C, Sullivan G, Stein MB, Roy-Byrne PP, Craske MG. Diagnostic overlap of generalized anxiety disorder and major depressive disorder in a primary care sample. *Depress Anxiety.* 2012; 29:1065–1071. [PubMed: 23184657]
- Zimmerman M, McDermut W, Mattia JI. Frequency of anxiety disorders in psychiatric outpatients with major depressive disorder. *Am J Psychiatry.* 2000; 157:1337–1340. [PubMed: 10910803]

Highlights

- Oxidative stress is poorly studied in relation to specific psychiatric symptoms.
- The sample comprised 69 physically healthy, unmedicated individuals with MDD.
- Oxidative stress is related to symptoms of anxiety, but not depression, in MDD.
- Anxiety may be an important contributor to, or a result of, oxidative stress in MDD.

Figure 1.

Scatter plots of the relationship between HAM-A scores (x-axis) and levels of (a) F2-isoprostanes and (c) GSSG (y-axis), and the relationship between HAM-D scores (x-axis) and levels of (b) F2-isoprostanes and (d) GSSG (y-axis). HAM-A and HAM-D scores are log transformed and F2-isoprostane and GSSG levels are (Blom) normalized and regressed on confounders age, sex, BMI, and smoking. Scatter plots display the data without multiply-imputed data, whereas statistical analyses include multiply-imputed data for missing values, as described in the text.

Table 1

Characteristics of the study sample.

	MDD	<i>n</i>^I
Age, years, mean (SD)	38.8 (13.9)	69
Sex, <i>n</i> men (%)	28 (40.6 %)	69
Body mass index, mean (SD)	25.7 (4.5)	69
Smoking, <i>n</i> (%)		69
Not at all	52 (75.4 %)	
Some days	9 (13.0 %)	
Every day	8 (11.6 %)	
HAM-D score, mean (SD)	20.0 (3.2)	69
Core Depression Factor Subscale score, mean (SD)	8.7 (1.8)	69
HAM-A score, mean (SD)	16.9 (4.9)	58
Hamilton Core Anxiety Scale score, mean (SD)	8.0 (2.8)	58
Psychic anxiety, <i>n</i> (%)		
Low	47 (68.1 %)	69
High	22 (31.9 %)	
Depressed mood, <i>n</i> (%)		69
Low	48 (69.6 %)	
High	21 (30.4 %)	
F2-isoprostanes, ng/ml, mean (SD)	.030 (.015)	65
GSSG, uM, mean (SD)	.198 (.090)	64
GSH, uM, mean (SD)	1.64 (.488)	64

^I Sample sizes are not all equivalent due to some subjects missing HAM-A or biochemical data. These missing data were handled using multiple imputation (discussed in section 2.5).

Table 2

Levels of oxidative stress markers for low vs. high anxiety and low vs. high depression (single item) subgroups.

	Low anxiety (<i>n</i> = 47)		High anxiety (<i>n</i> = 22)		Low depression (<i>n</i> = 48)		High depression (<i>n</i> = 21)	
F2-isoprostanes, ng/ml (mean, SD)	-.16 ± .12	.35 ± .23	-2.17	.030	-.03 ± .12	.07 ± .13	-0.41	.685
GSSG, uM (mean, SD)	-.18 ± .14	.39 ± .18	-2.32	.020	-.12 ± .13	.28 ± .23	-1.58	.115
GSH, uM (mean, SD)	-.05 ± .14	.11 ± .24	-0.58	.560	-.06 ± .15	.15 ± .22	-0.78	.436

Abbreviations: SD = standard deviation. Levels of oxidative stress markers are normalized residuals corrected for age, sex, BMI, and smoking. Accordingly, values can be positive or negative.