CORRESPONDENCE

The Cardiovascular Effect of Musical Genres—A Randomized Controlled Study on the Effect of Compositions by W. A. Mozart, J. Strauss, and ABBA

by Prof. Dr. med. Hans-Joachim Trappe, and Dr. med. Gabriele Voit
in issue 20/2016

Music and Neurophysiology

Trappe and Voit's interesting article on the cardiovascular effect of listening to music is the first to approach an interesting topic in a systematic fashion (1). In our opinion, the next step to be investigated should be the question of an association with neurophysiological parameters. This could be done relatively simply by using non-invasive, non-ionizing methods (for example, multichannel EEG surface mapping, functional near-infrared spectroscopy). In this way, it would be possible to verify whether the delinking between individually preferred musical styles and the measured effects also happens at the cerebral level. Subsequently, further controlled studies would be desirable, to compare the cardiovascular and neurophysiological measurements in healthy subjects and patients; patients with (chronic) pain or insomnia could be of multifarious interest in this setting (2, 3).

DOI: 10.3238/arztebl.2017.0043a

REFERENCES

1. Trappe H-J, Voit G: The cardiovascular effect of musical genres—a randomized controlled study on the effect of compositions by W. A. Mozart, J. Strauss, and ABBA. *Dtsch Arztebl Int* 2016; 113: 347–52.
2. Riemann D, Nissen C, Palagini L, Otte A, Perlis ML, Spiegelhalter K: The neurobiology, investigation, and treatment of chronic insomnia. *Lancet Neurol* 2015; 14: 547–58.
3. Otte A: Johann Sebastian Bach's „Goldberg variations“ to treat insomnia from renal lithiasis pain. *Sleep research in nuclear medicine. Hell J Nucl Med* 2016; 19: 13–4.

Prof. Dr. med. Andreas Otte
Offenburg University, andreas.otte@hs-offenburg.de

Methodological Limitations

Trappe and Voit's article picks up an interesting old topic but has methodological limitations (1). The slight changes in blood pressure are of questionable relevance even though they reached significance, especially as the method for measuring blood pressure was not described. In spite of being described as Riva-Rocci (auscultatory) measurement, the values—some of which were reported to two decimals—imply that the actual measuring technique was oscillometry, whose precision is suspected to be only ± 3 mm Hg. The effect of the measuring procedure itself was not considered.

The discussion attempts to explain the effects primarily by means of the characteristics of the composition, although hardly any published data exist to support this and, furthermore, this was not the subject of the study. We conducted an earlier study of the cardiovascular, hormonal, and psychological effects of the music of J Strauss (2), and we found fewer objective stress-reducing effects by comparison.

In another study using the same pieces of music, we found stress-reducing effects in patients with hypertension especially for the meditative music of Ravi Shankar, although this music was not con-

sistent with the kind of music the participants habitually consumed (3). We agree that elements of the composition are important, but we would also consider the factor of distraction (for example, from worries) as a further important component of the effect.

On the basis of these insights, we tested specifically composed music with a high degree of periodicity as receptive music therapy in a coronary sports group. We found that the patients benefited from the music intervention in terms of a lower degree of worry and anxiety, lower β -endorphin concentrations, and lower systolic blood pressure, in spite of physical exercise/activity (4).

It is regrettable that Trappe et al. did not make reference to these and further studies that investigated the effect of music on (psycho)physiological parameters. DOI: 10.3238/arztebl.2017.0043b

REFERENCES

1. Trappe H-J, Voit G: The cardiovascular effect of musical genres—a randomized controlled study on the effect of compositions by W. A. Mozart, J. Strauss, and ABBA. *Dtsch Arztebl Int* 2016; 113: 347–52.
2. Möckel M, Rucker L, Stork T, et al.: Immediate physiological responses of healthy volunteers to different types of music: cardiovascular, hormonal and mental changes. *Eur J Appl Physiol Occup Physiol* 1994; 68: 451–9.
3. Möckel M, Stork T, Vollert J, et al.: [Stress reduction through listening to music: effects on stress hormones, hemodynamics and mental state in patients with arterial hypertension and in healthy persons]. *Dtsch Med Wochenschr* 1995; 120: 745–52.
4. Vollert JO, Stork T, Rose M, Möckel M: [Music as adjuvant therapy for coronary heart disease. Therapeutic music lowers anxiety, stress and beta-endorphin concentrations in patients from a coronary sport group]. *Dtsch Med Wochenschr* 2003; 128: 2712–6.

Prof. Dr. med. Martin Möckel
Dr. med. Jörn Ole Vollert
Charité – Universitätsmedizin Berlin, martin.moeckel@charite.de

Effect Depends on Subjective Perception

Trappe and Voit showed in a methodologically sound study that merely listening to music goes hand in hand with a reduction in the concentrations of the stress hormone cortisol (1). This effect reached significance and was observed independently of the musical genre of the music on offer; furthermore, it was also not merely the expression of a physiological effect of rest.

This finding is consistent with music therapeutic routine experience and with the results of musical preference research—namely, that individually different musical genres are used to achieve a subjective state of relaxation. By contrast, the effects of different musical styles on heart rate and blood pressure were obviously less pronounced than in the control setting without music (1). Standardized serial measurements of heart rate variability (HRV) during rest might be suitable for confirming such changes (2). In a study of the effects of “calming/relaxing” or “stimulating” music on the HRV, we also found a drop over time, independently of the musical genre (Bertling B, Harenbrock S, Agelink MW: *Musik und Vegetativum* [music and the autonomic nervous system] . [conference abstract] Jahrestagung der Deutschen Gesellschaft für Musikpsychologie [AGM of the German Society for Music Psychology]; Karlsruhe, 17–19 September 1999). Neither musical style resulted in significant changes to the HRV. What was interesting, however, was that subjects who sub-

jectively liked the “calming/relaxing” music had a tendentially greater increase in the vagal modulation of the heart frequency than subjects who did not like the same piece of music. By contrast, Trappe and Voit could not demonstrate any significant effects resulting from subjective ratings, but then 90% of their subjects described all three musical styles as a pleasant sensation (1). The idea of compiling a “musical pharmacy” from certain musical genres that has a defined therapeutic effect in an existing indication is exciting—but it should not be allowed to blank out the subjective perception of the individual. DOI: 10.3238/arztebl.2017.0043c

REFERENCES

1. Trappe H-J, Voit G: The cardiovascular effect of musical genres—a randomized controlled study on the effect of compositions by W. A. Mozart, J. Strauss, and ABBA. *Dtsch Arztebl Int* 2016; 113: 347–52.
2. Agelink MW, Malessa R, Baumann B, et al.: Standardized tests of heart rate variability: normal ranges obtained from 309 healthy humans, and effects of age, gender and heart rate, *Clin Auton Res* 2001; 11: 99–108.

Stefan Harenbrock

Evangelische Kliniken Gelsenkirchen
Akademisches Lehrkrankenhaus der Universität Duisburg Essen

PD Dr. med. Marcus W. Agelink

Klinik für Psychiatrie, Psychotherapie & Psychosomatik
Klinikum Herford,
Sek.Psychiatrie@klinikum-herford.de

Improved Learning Performance

With regard to the article by Trappe and Voit on the cardiovascular effects of different musical genres, I would like to point out that the effect of different musical genres in cardiovascular parameters was confirmed previously. What is interesting, however, is the additionally observed drop in serum cortisol concentrations. This is a parallel trend to that observed in studies that have investigated the effect of different musical genres on psychological parameters (learning performance, attention, reading performance, memory performance) (1). In this particular study, music in general improves learning performance in terms of meaningless syllables, as these keep distracting stimuli in the background. This study showed—as did the one reported by Trappe and Voit—that rock music has the smallest positive effect (1, 2). A positive correlation between volume and improved learning performance of meaningless syllables was also seen. Our own study (3) using music by Mozart and Deep Purple yielded the same result. In view of the fact that the central importance of the stress hormone cortisol on psychological and biological parameters is being described to an increasingly more precise degree, and on the basis of the assumption that music has a crucial effect on this very cortisol level, the effect of music as “add-on” therapy should also be investigated for other parameters that are affected by cortisol. DOI: 10.3238/arztebl.2017.0044a

REFERENCES

1. Carlson JA, Hergenbahn BB: Effects of rock—roll and classical music on the learning of nonsense syllables. *Psychological Reports* 1967; 20: 2021–2.
2. Trappe H-J, Voit G: The cardiovascular effect of musical genres—a randomized controlled study on the effect of compositions by W. A. Mozart, J. Strauss, and ABBA. *Dtsch Arztebl Int* 2016; 113: 347–52.
3. Niederhofer H: Der Einfluß unterschiedlicher Musiktypen auf die Lernleistungen von Kindern. *Heilpädagogik* 2002; 45: 1–6.

CA DP Dr. phil. Dr. med. Helmut Niederhofer

Bezirkskrankenhaus Bayreuth
helmutniederhofer@yahoo.de

In Reply:

Although the association of music and health has been discussed for centuries, the interest in scientific studies of this topic is still quite low (1, 2). We therefore welcome the many positive responses and reactions to our study of the cardiovascular effect of different musical genres (3). Although we confirmed an effect on blood pressure and heart rate for the music of Strauss and Mozart, many unanswered questions remain that should be the subject of future studies. Especially neurophysiological studies are of interest—for example, using multichannel EEG—and especially in patients with (chronic) pain or insomnia (4). Further studies of the heart rate variability (HRV) at rest and of the effect of music and subjective perceptions are exciting and crucially essential in terms of assessing musical styles as a “musical pharmacy” (5). The fact that music affects psychological (learning performance, attention, memory), biological, and cortisol-affected parameters is well known and should certainly be the subject of further studies (6). Although our prospective randomized studies showed unequivocally positive effects on blood pressure and heart rate, it is too early to draw conclusions along the lines of “music as medication” for the purpose of blood pressure lowering. The phenomena of blood pressure lowering by means of music have also been observed by other authors in clinical studies and animal experiments. However, one publication cannot possible reference all other studies on the same subject, especially as *Deutsches Ärzteblatt* places great importance on up-to-the-minute reference lists.

The subject of music and medicine will certainly also be of great interest in future and is likely to gain in importance in conservative and operative medicine. In this setting, music needs to be understood as a multifactorial phenomenon, which achieves its effects by means of very diverse mechanisms. It remains to be seen whether there will be “music as medication”—for example, in pain therapy/management or arterial hypertension.

DOI: 10.3238/arztebl.2017.0044b

REFERENCES

1. Trappe HJ: The effects of music on the cardiovascular system and cardiovascular health. *Heart* 2010; 96: 1868–71.
2. Trappe HJ, Breker IM: Effects of different styles of music on human cardiovascular response: a prospective controlled trial. *Music & Medicine* 2016; 8: 8–16.
3. Trappe H-J, Voit G: The cardiovascular effect of musical genres—a randomized controlled study on the effect of compositions by W. A. Mozart, J. Strauss, and ABBA. *Dtsch Arztebl Int* 2016; 113: 347–52.
4. Riemann D, Nissen C, Palagini L, Otte A, Perlis ML, Spiegelhalder K: The neurobiology, investigation, and treatment of chronic insomnia. *Lancet Neurol* 2015; 14: 547–58.
5. Agelink MW, Malessa R, Baumann B, et al.: Standardized tests of heart rate variability: normal ranges obtained from 309 healthy humans, and effects of age, gender and heart rate. *Clin Auton Res* 2001; 11: 99–108.
6. Niederhofer H: Der Einfluß unterschiedlicher Musiktypen auf die Lernleistungen von Kindern. *Heilpädagogik* 2001; 45: 1–6.

On behalf of the authors

Prof. Dr. med. Hans-Joachim Trappe
Medizinische Universitätsklinik-Klinik II,
Marien Hospital Herne
Ruhr-Universität Bochum
hans-joachim.trappe@marienhospital-herne.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists.