Summary points

Organisational differences between the United States and United Kingdom present many possibilities for joint learning

The NHS primary care system offers many learning opportunities

The United States is ahead of the United Kingdom in research to understand the multiple determinants of disease but lags behind on equity in health

The United Kingdom could make better use of national patient data

marketing techniques to ascertain satisfaction. Too great a focus on patient satisfaction will detract from a more concerted effort to gain evidence from clinical practices, thus depriving both practitioners as well as consumers of a rational rather than a preference basis for decision making. It could be argued that need, rather than demand, should be the primary criterion for providing services, in the interest of better outcomes and more equitable distribution of resources.

Potential of teamwork

Both UK and US organisations employ teams of practitioners to provide primary care. In US managed care, it is common for nurse practitioners to act as frontline providers of care, with primary care doctors as back up. Because of the proprietary nature of managed care data, little is written about how decisions are made about who does what, and its impact in terms of quality and outcomes of care. This issue could well serve as a basis for a bi-national collaborative evaluation, with benefit to both countries.

Contributors and sources: This article reflects BS's experience in health services research and health policy, which derives from personal and policy contacts in many industrialised and developing nations.

Funding: This work was supported in part by Grant No 6 U30 CS 00189-05 S1 R1 of the Bureau of Primary Health Care, Health Resources and Services Administration, Department of Health and Human Services, to the Primary Care Policy Center for the Underserved at Johns Hopkins University.

Competing interests: None declared.

- 1 Starfield B, Shi L, Grover A, Macinko J. The role of evidence in physician workforce policy [web exclusive]. *Health Aff (Millwood)* 2005;Mar.
- 2 Forrest CB, Majeed A, Weiner JP, Carroll K, Bindman AB. Comparison of specialty referral rates in the United Kingdom and the United States: retrospective cohort analysis. *BMJ* 2002;325:370-1.
- 3 Forrest CB, Majeed A, Weiner JP, Carroll K, Bindman AB. Referral of children to specialists in the United States and the United Kingdom. *Arch Pediatr Adolesc Med* 2003;157:279-85.
- 4 Starfield B, Forrest CB, Nutting PA, von Schrader S. Variability in physician referral decisions. *J Am Board Fam Pract* 2002;15:473-80.
- 5 Fireman B, Bartlett J, Selby J. Can disease management reduce health care costs by improving quality? *Health Aff (Millwood)* 2004;23:63-75.
- 6 Ham C, York N, Sutch S, Shaw R. Hospital bed utilisation in the NHS, Kaiser Permanente, and the US Medicare programme: analysis of routine data. *BMJ* 2003;327:1257.
- 7 National Committee for Vital and Health Statistics. *Information for health: a strategy for building the national health information infrastructure*. Washington, DC: US Department of Health and Human Services, 2001.
- 8 Nolte E, McKee M. Measuring the health of nations: analysis of mortality amenable to health care. *BMJ* 2003;327:1129.
- 9 Bunker J. *Medicine matters after all: measuring the benefits of medical care, a healthy lifestyle and a just social environment*. London: Nuffield Trust, 2001.
- 10 Van den Akker M, Buntinx F, Roos S, Kottnerus JA. Problems in determining occurrence rates of multimorbidity. *J Clin Epidemiol* 2001;54:675-9.
- 11 Starfield B. New paradigms for quality in primary care. *Br J Gen Pract* 2001;51:303-9.
- 12 Salem-Schatz S, Moore G, Rucker M, Pearson SD. The case for case-mix adjustment in practice profiling. When good apples look bad. *JAMA* 1994;272:871-4.
- 13 Fisher ES, Wennberg DE, Stukel TA, Gottlieb DJ, Lucas FL, Pinder EL. The implications of regional variations in Medicare spending. Part 1: the content, quality, and accessibility of care. *Ann Intern Med* 2003;138:273-87.
- 14 Baicker K, Chandra A. Medicare spending, the physician workforce, and beneficiaries' quality of care [web exclusive]. *Health Aff (Millwood)* 2004; Apr 7. <http://content.healthaffairs.org/cgi/reprint/hlthaff.w4.184v1.pdf> (accessed 24 Feb 2004).
- 15 Wennberg JE. Perspective: Practice variations and health care reform: connecting the dots. *Health Aff* 2004; Suppl:VAR-140-VAR-144.
- 16 Majeed A, Evans N, Head P. What can PACT tell us about prescribing in general practice? *BMJ* 1997;315:1515-9.
- 17 US Department of Health and Human Services. *Healthy People 2010*. 2nd ed. Washington, DC: US Government Printing Office, 2000.
- 18 Black D. *Inequalities in health: report of a research working group*. London: Department of Health and Social Security, 1980.
- 19 Acheson D. *Independent inquiry into inequalities in health: report*. London: Stationery Office, 1998.
- 20 Senn S. Individual response to treatment: is it a valid assumption? *BMJ* 2004;329:966-8.
- 21 Mushlin AI, Appel FA. Testing an outcome-based quality assurance strategy in primary care. *Med Care* 1980;18:1-100.

A fatal error on the day of the new moon

In India the day of the new moon (Amavasya) in the month of July is considered to be highly auspicious. Various ancient rituals are practised on that day, one of which is drinking a herbal concoction prepared from the bark of the tree *Alstonia scholaris* (dita bark, devil tree, pale mara) early in the morning on an empty stomach. The bark of this tree is collected fresh, well before the sunrise, and is ground up to make a herbal preparation called kashaya. This preparation is believed to cure and prevent bowel related disorders. Clinical uses of this tree have been well described in homoeopathic and ayurvedic literature.

As they usually did, six members of a family prepared kashaya, but this year making it from the bark of a relatively young tree. Within a few minutes, all of them had severe muscle spasms and started having convulsive movements. They were rushed to hospital, but the two children, aged between 6 and 8 years, were dead on arrival. The four adult members of the family were conscious but had muscle stiffness with periodic convulsive movements of the limbs and opisthotonos. The clinical picture

was consistent with strychnine poisoning, and it was later confirmed that, in the dark, the family had taken the bark from *Strychnos nux-vomica* (strychnine tree) instead of *Alstonia*.

Ancient traditions and rituals tend to abound with precepts and injunctions. To modern eyes, these can seem meaningless superstition, but much ancient wisdom can be hidden amidst such rituals. This family had blindly followed the ritual of taking kashaya but had neglected to follow the golden rule, to "Select only those trees with old scars on their trunk." This practice of selecting trees with scars on the trunk is the only sure and safe way to identify the correct tree in darkness; old scars on the trunk affirm the tree's use in the previous year. Had our patients followed this tip, identification of the right tree would not have been difficult, and the tragedy could have been averted.

Sadananda Naik *cardiologist, Akva's Health Centre, Moodabidri, India* (sadananda@consultant.com)