Published in final edited form as:

Addiction. 2017 October ; 112(10): 1709–1715. doi:10.1111/add.13763.

How can we conceptualize behavioral addiction without pathologizing common behaviors?

Daniel Kardefelt-Winther^{a,*}, Alexandre Heeren^b, Adriano Schimmenti^c, Antonius van Rooij^d, Pierre Maurage^e, Michelle Carras^f, Johan Edman^g, Alexander Blaszczynski^h, Yasser Khazaalⁱ, and Joël Billieux^{e,j,k,*}

^aDepartment of Clinical Neuroscience, Karolinska Institutet, Sweden ^bDepartment of Psychology, Harvard University, USA ^cFaculty of Human and Social Sciences, Kore University of Enna, Italy ^dDepartment of Communication Sciences, imec-MICT-Ghent University, Belgium ^eLaboratory for Experimental Psychopathology, Université Catholique de Louvain, Belgium ^fDepartment of Mental Health, Johns Hopkins Bloomberg School of Public Health, USA ^gCentre for Social Research on Alcohol and Drugs, Stockholm University, Sweden ^hGambling Treatment Clinic and Research, School of Psychology, University of Sydney ⁱGeneva University Hospital, University of Geneva, Switzerland ^jInstitute for Health and Behavior. Integrative Research Unit on Social and Individual Development (INSIDE), University of Luxembourg, Luxembourg ^kInternet and gambling disorders Clinic, Department of Adult Psychiatry, Cliniques Universitaires Saint-Luc, Belgium

Abstract

Following the recent changes to the diagnostic category for addictive disorders in DSM-5, it is urgent to clarify what constitutes behavioral addiction to have a clear direction for future research and classification. However, in the years following the release of DSM-5, an expanding body of research has increasingly classified engagement in a wide range of common behaviors and leisure activities as possible behavioral addiction. If this expansion does not end, both the relevance and the credibility of the field of addictive disorders might be questioned, which may prompt a dismissive appraisal of the new DSM-5 sub-category for behavioral addiction. We propose an operational definition of behavioral addiction together with a number of exclusion criteria, to avoid pathologizing common behaviors and provide a common ground for further research. The definition and its exclusion criteria are clarified and justified by illustrating how these address a number of theoretical and methodological shortcomings that result from existing conceptualizations. We invite other researchers to extend our definition under an Open Science Foundation framework.

*Corresponding authors: Daniel Kardefelt-Winther, PhD, Department of Clinical Neuroscience, Karolinska Institutet, Tomtebodavägen 18A, 17176 Stockholm, Sweden., Phone: +44 79 46567850, Daniel.kardefelt.winther@ki.se. Joël Billieux, PhD, Research Unit INSIDE, University of Luxembourg, Maison des Sciences Humaines, 11, Porte des Sciences – L-4366 Esch-sur-Alzette, Luxembourg, T.: ++352 46 66 44 9000.; Joël.Billieux@uni.lu.

No competing financial interests exist.

1. Introduction

The fifth edition of the Diagnostic and Statistical Manual of Mental Disorders (1) includes a major change to the diagnostic category for addictions. The Substance-related Abuse and Dependency category has been relabeled Substance-Related and Addictive Disorders and modified to include two subdivisions, substance-related disorders and non-substance-related disorders, where the latter is defined as addictive disorders that do not involve ingestion of a psychoactive substance. This category is commonly referred to as behavioral addiction, although we note that there is no consensus yet as to how such a disorder should be defined¹.

The present debate paper has been written by a group of researchers from a number of different countries and academic fields who have a shared interest in how research on behavioral addiction is currently developing. In this paper, we wish to focus on the ongoing expansion of the behavioral addiction research area and its consequences for future research and clinical practice. We argue that considerable resources are increasingly being diverted to conduct research on excessive behaviors that lack indications of functional impairment, psychological distress or a clear separation from normative behavior in context, and therefore do not seem to constitute behavioral addiction. We are concerned that if this expansion does not end, both the relevance and credibility of this field might be questioned, which may prompt a dismissive appraisal of the new DSM-5 sub-category.

The sole condition that is currently included in the category for behavioral addiction is gambling disorder, with internet gaming disorder included in Appendix III as a potential addition to the category that first requires further study. That this category contains only one disorder is partly due to a lack of consensus on what precisely constitutes behavioral addiction, which is a consequence of insufficient peer-reviewed evidence for its etiology, onset and course (1). In addition to scarce evidence, we contend that research on behavioral addiction currently lacks a sound theoretical framework that can guide research in this area, help produce quality evidence and ensure a common ground for theoretical development. Although theoretical models exist that could guide research on addictions, whether substance-related or not, these are rarely used in behavioral addiction research, which tends to be largely atheoretical (2, 3).

For example, the syndrome model of addiction (4–6) usefully distinguishes the distal antecedents of an addiction (etiological processes) from its multiple expressions and manifestations (the behaviors and their symptoms). However, in the behavioral addiction research field, only the presence or absence of symptoms tend to be studied, most often guided by the symptom-based components model of addiction (7). Core elements of theory such as etiological processes and unique features of behavioral addiction are rarely investigated to the extent they are for substance addiction (8). Furthermore, the components model relies on symptoms of substance addiction in its definition of behavioral addiction. Several of these symptoms, such as tolerance and withdrawal, are difficult to convincingly

¹For the remainder of this paper we will use the term behavioral addiction to signify the DSM-5 category for non-substance-related disorders. Substance addiction will be used to signify the DSM-5 category for substance-related disorders.

apply and measure in relation to behaviors, which questions whether these symptoms form a useful and valid part of a behavioral addiction definition (2, 9, 10, 11).

Although West (8, 12) has provided an overview of many different theoretical approaches that can be used to understand substance addiction, ranging from neurological to sociological, none of these theories see much use in recent behavioral addiction research, and evaluations of the applicability of different approaches are rare. This has led to a situation where atheoretical and confirmatory approaches are far more common than research that is exploratory and theory-driven, which is unsuitable and ineffective for an emerging research area. The consequence has been a considerable expansion of the conceptualization of behavioral addiction to the point where we risk pathologizing common behaviors due to the lack of a clear theoretical framework.

To rectify this issue and in an attempt to provide a common ground for continued research, we propose an operational definition of behavioral addiction based on our collective understanding of harmful and persistent problem behaviors:

“A repeated behavior leading to significant harm or distress. The behavior is not reduced by the person and persists over a significant period of time. The harm or distress is of a functionally impairing nature.”

Additionally, diagnostic approaches often qualify the listing of inclusion signs with additional exclusion criteria. This is rarely considered when new expressions of behavioral addiction are proposed, but if we allow other potential explanations for a problem behavior to become part of the definition of behavioral addiction, we may cloud the treatment target, prolong potential suffering, and fail to identify problems that are actually due to another cause. Therefore, we include crucial exclusion criteria together with the definition and contend that a behavior should not be conceptualized as behavioral addiction if:

1. The behavior is better explained by an underlying disorder (e.g., a depressive disorder or impulse-control disorder).
2. The functional impairment results from an activity that, although potentially harmful, is the consequence of a willful choice (e.g. high-level sports).
3. The behavior can be characterized as a period of prolonged intensive involvement that detracts time and focus from other aspects of life, but does not lead to significant functional impairment or distress for the individual.
4. The behavior is the result of a coping strategy.

We recognize that this definition is not exhaustive and likely to require further input by experts in this field of research. It is meant to provide a starting point and guide researchers when conducting research on behavioral addiction, to avoid confirmatory research practices and an overreliance on substance addiction symptoms. In the interest of further conceptual development we have established an Open Science Foundation webpage to facilitate extension of the proposed operational definition in a transparent manner and invite interested researchers to comment and provide input (see 13 for a link).

Moving forward with this paper, we will highlight what we consider to be the most problematic theoretical and methodological practices and assumptions in the behavioral addiction research area and show how our proposed definition addresses each of these issues.

2. Which symptoms are useful and valid in behavioral addiction research?

Studies of behavioral addiction have often contended that individuals who engage excessively in certain behaviors experience a common set of symptoms frequently associated with substance addiction, such as salience, mood modification, tolerance, withdrawal, conflict and relapse, and that these symptoms can be used to identify behavioral addiction in people (7). The implication is that there is no limit as to what might be conceptualized as behavioral addiction as long as some of the common addiction symptoms are observed in relation to the behavior.

Problematically, these symptoms are likely to manifest in relation to most activities that people find interesting or engaging, without reflecting clinically significant functional impairment or distress for the individual or a burden to public health in populations. Some examples found in recent literature include (but are not limited to) “study addiction” (14), “work addiction” (15), “dancing addiction” (16), “mobile phone addiction” (17), “social network site addiction” (18), “fortune telling addiction” (19) and “body image addiction” (20). A positive addiction diagnosis or classification for these behaviors is particularly likely to be made when responses are captured through survey research using DSM-style polythetic cut-off scoring (e.g., meet 5/9 criteria for positive diagnosis), even though the symptoms in and of themselves are not indicative of functional impairment in community samples (9, 21). This diagnostic system also implicitly assumes that symptoms have equal weight and therefore contribute equally to a total score, which might not be the case (22). Additionally, an issue with the use of substance addiction symptoms when studying behavioral addiction is the failure to recognize that what constitutes a problematic symptom in relation to one activity (e.g., substance use) is not necessarily problematic in a different context (e.g., video gaming). For example, preoccupations with video games are still considered harmful in a similar way as preoccupations related to drugs, even though the former is a common everyday activity related to far fewer problematic consequences than the latter (21, 23, 24). This shows why using substance addiction symptoms in the conceptualization and assessment of behavioral addiction is inadequate and can lead to pathologizing of common behaviors or leisure activities. We propose instead a definition of behavioral addiction that focuses only on two components: a) significant functional impairment or distress as a direct consequence of the behavior, and b) persistence over time. While some studies consider functional impairment to be indicated by the presence of symptoms such as withdrawal, tolerance and conflict, the absence of in-depth studies with clinical cases means that it remains unclear whether these symptoms truly manifest in relation to behavioral addiction and, if they do, whether the symptoms are actually functionally impairing (23, 24). We therefore suggest not using these symptoms when operationalizing the definition unless more evidence is provided with regards to their relevance. Furthermore, while we include persistence as a key component, we have not yet proposed a period of time after which the behavior is classified as persistent because the lack

of evidence makes any such recommendation arbitrary. Persistence over time is an area where future research could usefully be conducted to improve the definition (e.g., 25).

For the purpose of research into new expressions of behavioral addiction we recommend directly assessing functional impairment in clinical settings, supported by a health professional, rather than via surveys implemented in a healthy population. A serious inquiry into excessive dancing as an expression of behavioral addiction would need to account for multiple explanations for any psychological distress or functional impairment that result. Acute stress reactions in pursuit of an unattainable ideal performance might be seen as a consequence of dysfunctional beliefs or elevated and rigid standards, such as those characterizing clinical perfectionism (26). The skeletal or muscle injuries that can result from excessive dancing (e.g., professional ballet) can be seen as a natural but unhealthy consequence of a career that puts a lot of pressure on the body. The exclusion criteria proposed together with our definition reflect both points raised. The first exclusion criterion is that *“the behavior is better explained by an underlying disorder”*, and the second exclusion criterion is that *“the functional impairment results from an activity that, although potentially harmful, is the consequence of a willful choice (e.g. high-level sports).”* These exclusion criteria are crucial to consider when conceptualizing behavioral addiction.

3. Avoid creating new disorders with old recipes

Billieux and colleagues (2) have argued that most studies identifying new expressions of behavioral addiction are confirmatory in nature. While this is a known issue for academic research that originates in epistemological discussions of hypothesis testing and falsification, it constitutes a considerable issue for research on behavioral addiction. Generally, research that identifies new expressions of behavioral addiction tends to follow the same three-step methodological pattern:

1. The research process seeks to confirm what the researchers believe to be true, rather than aiming to generate new findings, test hypotheses and contribute to theory-building (2, 27, 28). For example, based on initial observations of seemingly excessive involvement in a specific activity (e.g., some dancers can spend a whole day or night pursuing their hobby and will be tired at work the next day), the behavior is *a priori* conceptualized as behavioral addiction despite no indications of functional impairment and investigated as such because of its assumed similarity to substance addiction.
2. In the second step, diagnostic criteria and related screening tools are constructed by paraphrasing items initially developed to study substance addiction, in an attempt to match the substance addiction symptoms to the presumed behavioral addiction.
3. In the third step, these tools are used to collect cross-sectional survey data in order to investigate whether individuals who score highly on the items intended to capture core symptoms of this proposed behavioral addiction also report known risk factors for substance addiction (e.g., impulsivity traits, abnormal reward processing, cue reactivity). Any subsequent correlations are used to

establish that the disorder is similar to substance addiction in terms of correlates with known risk factors and therefore deserves clinical attention.

This process is inadequate in distinct ways for each step. At the first step, no alternative conceptualization is considered even though many seemingly excessive behaviors could also be considered an impulse control disorder, an obsessive-compulsive disorder, a maladaptive coping strategy, an engaging leisure activity or career. In other words, the proposed behavioral addiction is not clearly delineated from other disorders or from normative behavior. This is crucial because it is incorrect to assume that initial observation of excessive involvement is sufficient to warrant the label of behavioral addiction, or even that excessive involvement leads to functional impairment or distress. Most professional athletes and musicians could be considered as addicted if this approach was applied. This illustrates why substance addiction symptoms cannot uncritically be applied to behaviors, as they cannot adequately distinguish addiction from high engagement or passion (3, 29). For many behaviors and activities, without evidence of serious functional impairment, a substance addiction dimension such as craving when assessed by a self-reported questionnaire is, at the phenomenological level, similar to common desire (30). Furthermore, some behaviors are likely to be part of everyday life in many societies today even when engaged in very frequently, such as studying, playing video games and using mobile phones (31). These arguments justify the third exclusion criterion, which is that *“the behavior is characterized as a period of prolonged intensive involvement that detracts time and focus from other aspects of life, but does not lead to significant functional impairment or distress for the individual”*. While this criterion is implicitly accounted for by including functional impairment as part of the definition, the tendency in behavioral addiction research to conflate high engagement with addiction compels us to make this point explicit as an exclusion criterion. The risk of making a disorder out of a normative behavior or passion demands extra caution because it is likely to also affect and stigmatize individuals outside of the disordered population, for whom the behavior might bring many benefits.

For the second step, we reinforce our message that developing assessment instruments, and implicitly a definition, for behavioral addiction by substituting one behavior for another is inadequate (2, 21, 23). As highlighted with our example of athletes and musicians, assessment instruments aiming to measure a new disorder cannot have validity without an in-depth understanding of the problem domain, its unique features and natural boundaries (27, 32), even if some common features are shared with existing disorders at the phenomenological level. A revealing example of why this is problematic is the largely unsuccessful effort to apply the criteria for tolerance and withdrawal in behavioral addiction research, which has been criticized for involving either metaphorical use of these terms or the use of fairly coarse behavioral criteria such as patient's complaints of feeling irritable (10, 33, 34). Additionally, the physiological component involved in withdrawal and tolerance symptoms do not manifest through behavior alone which makes direct translation of the criteria unreliable (9, 10). The practice of using substance addiction symptoms in the study of behavioral addiction has resulted in a lack of theoretical specificity for its different expressions since new and more accurate criteria are rarely – if ever – developed (3, 27). From an epistemological perspective, our understanding of excessive and repetitive behaviors is currently restricted to the theoretical boundaries of substance addiction (27).

This is a concern when approaching new problem domains with unknown problem manifestations.

As a consequence, we note that in the third step, to investigate patterns of correlation with known risk factors of substance addiction is at this point almost meaningless, as the validity of a measurement constructed in a confirmatory fashion is highly questionable. Establishing similarities between substance addiction and behavioral addiction is useful only in so far as we can trust our understanding of the two disorders; the above paragraphs have illustrated why current understandings of behavioral addiction are likely to be flawed in many respects. Putting too much faith in comparisons with substance addiction at this point could lead to a number of negative consequences for research and clinical practice, such as an inflation of prevalence figures, misdiagnosis or ineffective prevention and treatment. Instead of relying on substance addiction symptoms, researchers should work to identify the etiological and psychological processes that underlie a specific excessive behavior, which demands focus on the unique manifestations of the behavior as well as the resulting impairment, rather than on symptoms of excessive involvement *per se*. Long-term impairment seems particularly relevant to assess for behavioral addiction, as longitudinal studies have shown that a number of excessive behaviors seem to be fairly transient for most people and are episodic rather than steady in nature (25). This justifies our focus on the persistence of the behavior in the proposed definition and explains why we deliberately chose not to include other substance addiction symptoms.

Finally, an excessive behavior (whether it leads to treatment or not) may also constitute a helpful or maladaptive coping strategy, which on the surface may be similar to behavioral addiction as it can be repetitive and frequently recurring (35, 36). If the possibility of a coping strategy is not considered when the behavior is evaluated, the patient may be treated for behavioral addiction instead. The behavior may or may not change as a result, but the underlying problem is unlikely to be affected (37). Thus, we include coping strategies as the fourth exclusion criterion in the proposed definition, which needs to also be considered in relation to a targeted behavior.

4. Considerations for improving research on behavioral addiction

To avoid the theoretical and methodological issues presented in this paper, studies should seek to generate evidence that identifies and promotes a further description of the factors that distinguish healthy repeated behavior from harmful disorder. A clear deviation from normative behavior should be observed and reported functional impairment and psychological distress should be required for a behavior to warrant further study as an expression of behavioral addiction. To observe excessive involvement alone, which might only reflect engagement, passion, or coping, is inadequate. We believe this is captured in our proposed definition.

We suggest that one way to yield useful evidence that has not yet been given enough attention is by taking a person-centered approach to research, focusing on qualitative studies with individuals reporting significant functional impairment and distress as a consequence of a targeted behavior. A key aim should be to explore the phenomenology of a proposed

condition and identify its etiology and course. Only in a second step should items be developed and assessment instruments evaluated in terms of psychometric properties (27). This approach has been used with much success in the study of gambling disorder (38–41), but is rather rare in the wider study of behavioral addiction. The phenomenology and symptomatology of an excessive behavior needs to be theorized and empirically evaluated through hypothesis-driven theory testing to ensure that we are not restricting the scientific venture to inquire only about symptoms of known relevance for substance addiction, and to ensure that the behavior is not misinterpreted as behavioral addiction where other explanations are more appropriate. Such detailed theoretical knowledge would provide the greater confidence in classification required to conduct more advanced research, such as treatment, neuro-imaging or genetics studies. Following this line of thought – where appropriately developed theory precedes survey research for a specific excessive behavior – we maintain that excessive problem behaviors should not be *a priori* theorized as behavioral addiction without sound empirical evidence to support this conceptualization. We suggest that future research conceptualizes new expressions of behavioral addiction according to our proposed definition, after serious consideration of the exclusion criteria. As argued by Clark and Watson (42), *a good theory articulates not only what a construct is, but also what it is not* (p.5).

We realize that the exclusion criteria proposed in this paper raises the bar significantly in terms of what may or may not be conceptualized as an expression of behavioral addiction. This is also our intention, as we believe no one stands to gain if behaviors are conceptualized as expressions of behavioral addiction when, in fact, they are not. By introducing a more stringent definition we implicitly ask researchers to explore each problem behavior in-depth before conceptualizing it as behavioral addiction, which means that our overall understanding of behavioral addiction and its boundaries to other problem behaviors should improve.

To conclude, we hope that our proposed definition will be helpful for researchers working on behavioral addiction and we look forward to further developing this definition together with others. We also hope that the methodological issues and problematic research practices highlighted in this paper can be avoided as we move forward, partly through the use of a more appropriate definition of behavioral addiction and the proposed exclusion criteria. We believe future research on new expressions of behavioral addiction should consider the public and/or mental health implications of a behavior (i.e., impact on subjective distress, nature and severity of harms, functional impairment), as well as clearly establish the boundary between healthy engagement and disorder, taking into account normative aspects of a given behavior. The proposed definition should initially be used to guide qualitative research with patients who report significant functional impairment as a consequence of a repeated behavior with the aim of improving the definition, rather than for population-based studies.

Acknowledgments

Disclosed funding sources: Michelle Carras: National Institute of Mental Health (NIMH) Training Grant 5T32MH014592-39; Alexandre Heeren: Helaers Foundation for Medical Research (Grant : Complexity in psychopathology); Belgian Foundation for Vocation (Scientific Vocation), and the WBI World Excellence Grant in

Life Sciences - BLOWIN (grant number: sub/2015/228106243177); Pierre Maurage: Belgian Fund for Scientific Research (FRS-FNRS)

References

1. American Psychiatric Association. Diagnostic and Statistical Manual of Mental Disorders. 5. Arlington, VA: Washington, DC: 2013.
2. Billieux J, Schimmenti A, Khazaal Y, Maurage P, Heeren A. Are we overpathologizing everyday life? A tenable blueprint for behavioral addiction research. *Journal of Behavioral Addictions*. 2015a; 4:119–23. [PubMed: 26014667]
3. Kardefelt-Winther D. Problems with atheoretical and confirmatory research approaches in the study of behavioral addictions. *Journal of Behavioral Addictions*. 2015; 4:126–29. [PubMed: 26551896]
4. Shaffer HJ, LaPlante DA, LaBrie RA, Kidman RC, Donato AN, Stanton MV. Toward a syndrome model of addiction: multiple expressions, common etiology. *Harvard Review of Psychiatry*. 2004; 12:367–74. [PubMed: 15764471]
5. Shaffer, HJ., LaPlante, DA., Nelson, SE. The APA Addiction Syndrome Handbook (Vol. 1. Foundations, Influences, and Expressions of Addiction). Washington, DC: American Psychological Association Press; 2014.
6. Shaffer, HJ., LaPlante, DA., Nelson, SE., editors. The APA Addiction Syndrome Handbook (Vol. 2. Recovery, Prevention, and Other Issues). Washington, DC: American Psychological Association Press; 2012.
7. Griffiths MDA. “components” model of addiction within a biopsychosocial framework. *Journal of Substance Use*. 2005; 10:191–97.
8. West, R., Hardy, A. Theory of addiction. Oxford: Wiley-Blackwell; 2006.
9. Van Rooij AJ, Prause N. A critical review of “Internet addiction” criteria with suggestions for the future. *Journal of Behavioral Addictions*. 2014; 3:203–13. [PubMed: 25592305]
10. Starcevic V. Tolerance and withdrawal symptoms may not be helpful to enhance understanding of behavioural addictions. *Addiction*. 2016; 111:1307–08. [PubMed: 27095413]
11. Kaptis D, King DL, Delfabbro PH, Gradisar M. Withdrawal symptoms in internet gaming disorder: A systematic review. *Clinical Psychology Review*. 2015; 43:58–66. [PubMed: 26704173]
12. West, R. EMCDDA Insights Series No 14. Lisbon, Portugal: EMCDDA; 2013. EMCDDA insights: Models of addiction.
13. Billieux, J., Blaszczynski, A., Colder Carras, M., Edman, J., Heeren, A., Kardefelt-Winther, D., Khazaal, Y., Maurage, P., Schimmenti, A., van Rooij, AJ. Behavioral Addiction: open definition development. 2016. <http://doi.org/10.17605/OSF.IO/Q2VVA>
14. Atroszko PA, Andreassen CS, Griffiths MD, Pallesen S. Study Addiction-A new area of psychological study: Conceptualization, assessment, and preliminary empirical findings. *Journal of Behavioral Addictions*. 2015; 4:75–84. [PubMed: 26014668]
15. Griffiths MD. Workaholism is still a useful construct. *Addiction Research and Theory*. 2005; 13:97–100.
16. Maraz A, Urbán R, Griffiths MD, Demetrovics Z. An empirical investigation of dance addiction. *PloS ONE*. 2015; 10:e0125988. [PubMed: 25951077]
17. Chóliz M. Mobile phone addiction: A point of issue. *Addiction*. 2010; 105:373–74.
18. Andreassen CS, Pallesen S. Social network site addiction: An overview. *Current Pharmaceutical Design*. 2014; 20:4053–61. [PubMed: 24001298]
19. Grall-Bronnec M, Bulteau S, Victorri-Vigneau C, Bouju G, Sauvaget A. Fortune telling addiction: unfortunately a serious topic. *Journal of Behavioral Addictions*. 2015; 4:27–31. [PubMed: 25786497]
20. Foster AC, Shorter GW, Griffiths MD. Muscle dysmorphia: could it be classified as an addiction to body image? *Journal of Behavioral Addictions*. 2015; 4:1–5.
21. Griffiths MD, van Rooij AJ, Kardefelt-Winther D, Starcevic V, Király O, Pallesen S, et al. Working towards an international consensus on criteria for assessing Internet Gaming Disorder: A critical commentary on Petry et al (2014). *Addiction*. 2016; 111:167–78. [PubMed: 26669530]

22. Van Rooij, AJ., Van Looy, J., Billieux, J. Internet Gaming Disorder as a formative construct: Implications for conceptualization and measurement; *Psychiatry and Clinical Neurosciences*. 2016. p. 1-14.<http://doi.org/10.1111/pcn.12404>
23. Kardefelt-Winther D. A conceptual and methodological critique of internet addiction research: towards a model of compensatory internet use. *Computers in Human Behavior*. 2014; 31:351–54.
24. King D, Delfabbro P. Is preoccupation an oversimplification? A call to examine cognitive factors underlying internet gaming disorder. *Addiction*. 2014; 109:1566–70. [PubMed: 24735375]
25. Thege BK, Woodin EM, Hodgins DC, Williams RJ. Natural course of behavioral addictions: a 5-year longitudinal study. *BMC Psychiatry*. 2015; :15.doi: 10.1186/s12888-015-0383-3 [PubMed: 25652365]
26. Shafran R, Cooper Z, Fairburn CG. Clinical perfectionism: a cognitive-behavioural analysis. *Behaviour Research and Therapy*. 2002; 40:773–91. [PubMed: 12074372]
27. Kardefelt-Winther D. Making the case for hypothesis-driven theory testing in the study of Internet Gaming Disorder. *Addictive Behaviors*. 2017; 64:234–237. DOI: 10.1016/j.addbeh.2015.09.012 [PubMed: 26455330]
28. Shaffer HJ, Hall M, Vander Bilt J. “Computer addiction”: a critical consideration. *American Journal of Orthopsychiatry*. 2000; 70:162–68. [PubMed: 10826028]
29. Charlton JP, Danforth IW. Distinguishing addiction and high engagement in the context of online game playing. *Computers in Human Behavior*. 2007; 23:1531–48.
30. Kavanagh DJ, Andrade A, May J. Imagery relish and exquisite torture: The elaborated intrusion theory of desire. *Psychological Review*. 2005; 112:446–67. [PubMed: 15783293]
31. Sussman S, Lisha N, Griffiths MD. Prevalence of the addictions: a problem of the majority or the minority? *Eval Health Prof*. 2011; 34:3–56. [PubMed: 20876085]
32. Kendell R, Jablensky A. Distinguishing between the validity and utility of psychiatric diagnoses. *Am J Psychiatry*. 2003; 160:4–12. [PubMed: 12505793]
33. Pies R. Should DSM-V Designate “Internet Addiction” a Mental Disorder? *Psychiatry*. 2009; 6:31–37. [PubMed: 19724746]
34. Billieux J, Maurage P, Fernandez-Lopez O, Kuss DJ, Griffiths MD. Can disordered mobile phone use be considered a behavioral addiction? An update on current evidence and a comprehensive model for future research. *Current Addiction Reports*. 2015; 2:156–162.
35. Lazarus, RS., Folkman, S. Stress, appraisal, and coping. New York: Springer; 1984.
36. Kardefelt-Winther D. Conceptualizing Internet use disorders: Addiction or coping process? *Psychiatry and Clinical Neurosciences*. 2016; doi: 10.1111/pcn.12413
37. Schimmenti A, Caretti V. Psychic retreats or psychic pits? Unbearable states of mind and technological addiction. *Psychoanalytic Psychology*. 2010; 27:115–32.
38. Lesieur H. The Compulsive Gambler’s Spiral of Options and Involvement. *Psychiatry*. 1979; 42:79–87. [PubMed: 760136]
39. Ricketts T, Macaskill A. Gambling as emotion management: Developing a grounded theory of problem gambling. *Addiction Research and Theory*. 2003; 11:383–400.
40. Ricketts T, Macaskill A. Differentiating normal and problem gambling: A grounded theory approach. *Addiction Research and Theory*. 2004; 12:77–88.
41. Wood R, Griffiths M. A qualitative investigation of problem gambling as an escape-based coping strategy. *Psychology and Psychotherapy: Theory, Research and Practice*. 2007; 80:107–25.
42. Clark LA, Watson D. Constructing validity: Basic issues in objective scale development. *Psychological Assessment*. 1995; 7:309–19.