RESEARCH

A quasi-experimental evaluation of dried blood spot testing through community pharmacies in the Tayside region of Scotland

Andrew Radley,¹ Karen Melville,² Jan Tait,³ Brian Stephens,³ Josie M M Evans,⁴ John F Dillon⁵

¹NHS Tayside, Directorate of Public Health, Kings Cross Hospital, Dundee, UK

²NHS Tayside, Tayside Substance Misuse Services, Dundee, UK

³NHS Tayside, Gastroenterology Department, Ninewells Hospital and Medical School, Dundee, UK

⁴Division of Health Sciences and Sport, University of Stirling, Stirling, UK

⁵Division of Cardiovascular Medicines and Diabetes, Ninewells Hospital and Medical School, University of Dundee, Dundee, UK

Correspondence to

Andrew Radley, NHS Tayside, Directorate of Public Health, Kings Cross Hospital, Clepington Road, Dundee DD3 8EA; 425681, UK; Andrew.Radley@nhs.net

Received 9 November 2016

Revised 11 December 2016

Accepted 13 December 2016

Published Online First

6 January 2017

ABSTRACT

Objective Comparison of uptake of dried blood spot testing (DBST) for hepatitis C virus (HCV) infection between community pharmacies and established services.

Design Quantitative evaluation of a service development with qualitative process evaluation undertaken in parallel.

Setting Six pharmacies from 36 community pharmacies within Dundee City, a large urban settlement with high levels of socioeconomic deprivation.

Participants Patients in receipt of opioid substitution therapy (OST) not tested for HCV within 12 months. The 6 pharmacies provided OST for approximately 363 patients from a cohort of 1385 patients within Dundee City.

Intervention Provision of DBST by pharmacists.

Main outcome measure Receipt of DBST between January and December 2014.

Results 43 of 143 service users with no record of testing from the 6 community pharmacies accepted DBST. Of 561 from the remaining 1022 service users with no record of testing, 75 were tested for HCV (30% vs 13%). The OR for increased uptake of testing within the 6 pharmacies was 2.25 (95% CI 1.48 to 3.41, Z statistic=3.81, $p<0.0001$) compared with other services. The DBST taken by the pharmacies provided 12 patients with a reactive test. The process evaluation identified key themes important to staff and recipients of the service. A logic model was constructed.

Limitations Non-experimental service evaluation performed in community pharmacies records service activity in one location across a single time period.

Interpretation Some evidence that DBST from community pharmacies may be feasible. Service

users received the service positively. Staff reported that DBST was straightforward and achievable.

BACKGROUND

Hepatitis C virus (HCV) infection is a bloodborne infection causing liver disease. Around 1% of the population may be infected with HCV and around 0.8% is chronically infected.¹ The greatest risk of acquiring the virus in the UK is through injecting drug use. Patient outcomes from HCV infection vary, with 25% clearing the infection spontaneously and the remainder becoming chronically infected, risking development of cirrhosis and hepatocellular carcinoma. People infected with HCV may show no symptoms, presenting with incurable, end-stage disease. A recent Public Health England report highlighted that less than 3% of those known to be infected with HCV are being treated and less than half of those infected are known.² The largest single infected group is that on opioid substitution therapy (OST).³ Research suggests around 40% of people receiving OST has HCV.^{4 5}

The conventional NHS pathway of care is that patients with a history of intravenous drug use or those currently prescribed OST should be offered HCV testing.¹ Testing is commonly available from a range of primary care and third sector providers. Once diagnosed, patients may be referred to nurse led treatment pathways, based around hepatology or infectious disease teams in secondary care.

CrossMark

To cite: Radley A, Melville K, Tait J, et al. *Frontline Gastroenterology* 2017;**8**:221–228.

In current pathways, less than 10% of the OST population is tested for HCV. Of those tested, at very best 25% start treatment in one of the dedicated centres, with 70–80% successfully completing. This means that only 2–3% of this vulnerable population receives adequate treatment.² The new highly effective directly acting antiviral drugs achieve cure rates in excess of 90%, with once or twice daily tablets for 8–12 weeks and few side effects.⁶

The use of dried blood spot testing (DBST) in non-traditional environments has dramatically increased detection of HCV.^{7–8} DBST has been shown to be a reliable alternative to taking venous blood samples and determining HCV status in drug injectors. With appropriate training it can be carried out by all staff. Several practice-based projects have sought to implement DBST into pharmacy practice.⁹

Pharmacists have daily interactions with patients receiving methadone and we hypothesised that this relationship could be wielded to deliver increases in rates of testing using a DBST approach. To test this we employed a quasi-experimental design¹⁰ to compare the uptake of DBST for HCV in a small group of community pharmacies, with uptake in established services (those in substance misuse services, general practices and in third sector organisations).

METHODS

A quasi-experimental design was chosen as a pragmatic route to evaluate the feasibility and scalability of a service development in community pharmacy, because of uncertainties about the effect size and nature of the intervention.¹¹

This study was carried out in the city of Dundee, within the Tayside region of Scotland, a large urban settlement with significant socioeconomic deprivation.¹² There are 36 community pharmacies that provide OST for approximately 1385 patients, within a total population of 148 000.¹³ Six pharmacies were trained to offer DBST. Pharmacies were selected if they provided OST supervision for at least 30 patients and staff was willing to participate. Patients in the comparator group were prescribed OST and attended a community pharmacy that did not offer DBST.

Design of the intervention

Focus group interviews

A focus group series was undertaken before implementation to identify service user responses to the offer of testing, using a co-production approach.¹⁴ A purposive sampling strategy recruited 41 participants in seven focus groups, to gain a diversity of views (table 1). All interviews and focus group discussions were recorded as digital audio files and transcribed in full for thematic analysis.¹⁵ Analysis drew on the constant comparison method, operationalised within a general thematic approach.¹⁶

Table 1 Focus group participant details

	Participants (n)
Age group at participation, years	
<35	17
35–44	5
45–54	6
Over 55	4
Did not disclose	9
Sex	
Male	31
Female	10
Participant category	
Service user	38
Carer	3
Focus group venue	
Large urban settlement	10
Other urban settlement	10
Accessible rural town	6
Prison educational centre	7
Women's group	4
Peer mentor's group	4

Development of programme theory

Figure 1 describes a logic model that was constructed to explicitly identify targets for evaluation and data collection. The process evaluation took cognisance of the normalisation process theory¹⁷ using the core constructs of coherence, cognitive participation, collective action and reflexive monitoring to conceptualise the processes involved with effectiveness of implementation.

Implementation of DBST

Staff from the six community pharmacies were trained to perform DBST to consenting patients.¹⁸ DBST employs a finger prick of blood, spotted onto protein saving paper. The sample was tested by medical microbiology for HCV, HIV and hepatitis B antibodies.⁷ Fourteen staff members from the pharmacies attended a 2-hour interactive teaching session. The six pharmacies were notified which of their OST patients had no history of HCV testing, as identified by the central virology laboratory. Testing for HCV was offered to each of these individuals. All individuals with a reactive antibody HCV test were offered referral to the specialist hepatitis service.

Participants' comments during the training were noted and each pharmacy was visited several times to discuss progress. Notes of these interactions and emails from participants were retained for analysis. Testing took place during the last quarter of 2014.

Post-testing interviews

Semistructured interviews were conducted with (1) 8 service users and (2) 10 professionals taking part in the study, with all six pharmacies represented. The interviews were conducted using two topic guides

Figure 1 Logic model for the dried blood spot testing (DBST) implementation process.

developed in line with the research aims and programme theory. All interviews were recorded as digital audio files and transcribed in full for thematic analysis.^{15 16} These data contributed to assessment of feasibility and acceptability (including barriers and facilitators), identifying any unintended consequences of participation. Transcripts were inductively analysed to identify themes emergent from the interviews. A deductive analysis was also undertaken to compare findings with programme theory.

Quantitative data collection and analysis

Data on demographic information, risk factors, laboratory tests, referral, follow-up and treatment were collected. The age structures of the intervention and comparator groups were compared by t-test, as were data on DBST taken by the pharmacies. An OR was calculated for the numbers of service users undertaking testing at pharmacies compared with non-pharmacy services.

Ethics approval

Study documents were submitted to the East of Scotland Research Ethics Service, who confirmed that this work did not require ethical approval under its terms of governance. Caldicott permission was gained to enable access to and analysis of patient information.

RESULTS

The age structure of the 143 service users (26% of all OST users) with no record of testing from the six

pharmacies, together with the 1022 service users of the comparator group are presented in [table 2](#).

Analysis of the age structure of intervention and comparator groups was undertaken. A statistically significant difference between the two groups was detected using the two-sided t-test ($p < 0.05$). The intervention group was younger by 1.75 years (mean age of 35.6 years) compared with the comparison group (mean age of 37.4 years). The study design meant that intervention and comparison groups were not randomised, and therefore equal mean ages of the two groups were not expected.

During the test period, 43 (30%) of the 143 service users from the six community pharmacies accepted a DBST. Within the comparator group 561 from 1022 service users had no history of DBST and a total of 75 (13%) patients was tested for HCV by other providers of DBST (30% vs 13%). The OR for increased uptake of testing within the six pharmacies was 2.25 (95% CI 1.48 to 3.41, Z statistic=3.81, $p \leq 0.0001$) in comparison to the other services ([table 2](#)). The six pharmacies identified 12 patients with a reactive test. The uptake of DBST by each of the six pharmacies is presented in [table 3](#). No statistically significant difference in uptake between the sites could be detected at the $p < 0.05$ level using two-sided t-testing.

Understanding of context for delivery

Themes identified from transcript analysis of the focus groups provided information about the context for delivery.¹⁹ Focus groups discussions demonstrated an understanding of HCV including latency and

Table 2 Age structure of service users prescribed opioid substitution therapy (OST) in Dundee

Age (years)	Intervention cohort	Intervention cohort (%)	Comparison cohort	Comparison cohort (%)	Total population	Total population (%)
<19	0	~	0	~	0	~
20–24	14	4	23	2	37	3
25–29	51	14	133	13	184	13
30–34	112	31	256	25	368	27
35–39	86	24	239	23	325	23
40–44	61	17	185	18	246	18
45–49	26	7	120	12	146	11
50–54	10	3	48	5	58	4
55–59	3	1	16	2	19	1
60–64	0	0	2	0	2	0
65–69	0	0	1	0	1	0
	363	100	1023	100	1386	100

Table 3 Uptake of testing by pharmacy site

Pharmacy site	Eligible patients (n)	Tests taken, n (% of eligible patients)	Positive tests (n)
A	23	13 (57)	3
B	22	11 (50)	4
C	30	5 (17)	3
D	26	10 (38)	1
E	26	3 (12)	1
F	16	1 (6)	0
Total	143	43 (30)	12

effects on general health (figure 2). Participants could talk about cirrhosis and cancer. Participants knew about the burden of treatment of interferon-based regimes. Several clients knew about the cost of the new treatments.

Hepatitis C was viewed as a ‘dirty disease’ caught from sharing needles. It was unacceptable to admit to having HCV and participants feared social exclusion if found out. Participants described the shame they expected to feel if their family discovered they had HCV and spoke about ‘letting them down’. The fear of being found out contributed to reticence about testing.

What factors influenced client recruitment and participation?

The transcripts from the post-test interviews demonstrated that service users had both positive and negative perceptions of testing. Interviewees clearly thought that pharmacies were a good place to be tested and valued this service and the positive relationships built with pharmacy staff. Lack of money meant travelling to a local hospital was a barrier to clinic attendance. Service users talked vividly about their diminished horizons: they felt this was due to their dependence on drugs and lack of money. Pharmacies however were viewed as part of the local community.

Some service users’ previous experiences of stigma and discrimination when attending a pharmacy led to caution and suspicion when they were offered testing. One client complained that they were offered DBST standing at the dispensary counter, rather than in the private consultation room. Clients with negative experiences suggested staff would need intensive training before offering the service. Some clients expressed the wish for a better explanation about how the test worked and what the results might mean. Clients with positive experiences raised no issues for service improvement.

Several interviewees distinguished themselves from other ‘junkies’. These interviewees would describe their plans for recovery and would describe their relationship with a partner or young family. These individuals believed that service users with more chaotic lives were less likely to be tested.

What feedback on staff training and implementation was received?

Staff interviewees had clear views about successful implementation. Staff considered that strong leadership and involving all the team were necessary. Comments such as ‘a great opportunity’, ‘an obvious thing to do’ and ‘a no brainer’ were made. The degree of enthusiasm for new roles and positive relationships with patients were important. Only one individual viewed the development ‘as a step beyond what pharmacies do’. There were some initial anxieties expressed about potential contact with infected blood. Some staff viewed DBST as a pharmacist’s role, whereas others thought it was a team role.

The training was evaluated positively, with comments received about the simplicity of DBST. Several teams took lancets and protein-saving cards with them to practice the technique. Implementation was viewed as most successful where all the teams were involved.

Some staff expressed concern about offering DBST when prescription dispensing workload was high. An identified solution was to use time slots when

What do service users understand by hepatitis C infection and how they obtain testing and treatment?

Perceptions of disease

"my daughter and partner have it. It can lie dormant. it makes you feel very tired and you have no appetite" (Participant 5, Carer, December 11)

"You could have hepatitis even though, you could be stable on methadone for years then find out you've got hepatitis" (Participant 35, 22 April)

Perceptions of testing and treatment

I think more people have got it done since they started doing that 'cause it's easier (DBST) (Participant 37, 22 April)

"What they do is they have a look at your white blood cells count and things like that and how your system is reacting to it and whether or not. See, when you get the finger prick test for Hepatitis it'll, if you have been exposed to the Hepatitis virus at all, it will come back positive. That doesn't necessarily mean to say that you have got Hepatitis, so what they then is they will do a blood test and start talking about counts and things like that as its really confusing" (Participant 19, 25 February)

You see my partner is really skinny as it is and I think he will loose, everybody;s different are they, I think he will loose a lot of weight and I think it affects them mentally as well (Participant 33, 7 March)

Preferences for obtaining treatment

It would have tae be someplace where dinnae have tae travel everywhere where its gonnae cost them money that they have nae always got that's, that's what I feel is a big thing having tae travel tae Ninewells all the time (Participant 27, 5 March)

I think that's one of the things that's gonnae be down tae the individual, it depends on how chaotic their lifestyle is otherwise there's gonnae be a lot of money wasted there. And I take it this new stuff isnae cheap coz it never is when its first introduced. (Participant 22, 5 March)

"Well I say your pharmacy because like I say you're there every day, you're only, some people dinnae even go to the DPC once a month. So and then you forget your appointment, some people forget their appointment at the DPC but you're guaranteed to go for your methadone every day so" (Participant 36, 22 April)

Aye but you could be going tae the pharmacy for anything you see, so this again de-stigmatising. Nobody knows what they're in there for, it could be a drug problem it could be... anything (Participant 19, 25 February)

Figure 2 Examples of focus group quotations.

workload was less. Time barriers were most important where DBST was seen as a pharmacist role.

DISCUSSION

Key findings

This study provides some evidence that pharmacies may be a feasible site from which to offer DBST. People receiving OST were more likely to accept a DBST from a pharmacy than from other local providers. The evaluation of the implementation provides some evidence of

the context for delivery and the mechanisms that may lie behind the outcomes observed.¹⁹ Contextual factors included: expectations and experiences of stigma and discrimination; fears about confidentiality; the limited horizons of people receiving OST and the poverty they experience. Identified mechanisms that may influence uptake included the presence of established relationships with pharmacy staff; a pre-existing reason for attending the pharmacy for OST and the proximity of the pharmacy within the local community.

Review of programme theory

The programme theory provided a useful structure from which to consider the process evaluation and enabled a series of insights into the barriers and facilitators to effective implementation. In particular, we valued insights into the context that testing in pharmacies created for service users and the meaning that such testing may have. The evaluation also provided an indication about how the attitudes and behaviours of staff towards the intervention contributed to success or otherwise.

A number of barriers to uptake of testing and treatment including fear of blood tests,²⁰ adverse socio-economic and family circumstances, as well as fear of treatment side effects were identified in other studies as well as being identified in this study.²¹ Evidence suggests that offering DBST may increase the uptake of HCV testing, when compared with venepuncture, although this finding may also reflect increased availability and access when a DBST technique is used.^{7 22 23} Community-based settings for HCV services may also increase acceptance and uptake of testing.^{24–27} Provision of on-site testing may also have positive effects on uptake, due to the proximity of the testing offer.²²

Different settings have been used to increase the uptake of testing and treatment including testing from methadone maintenance services²⁸ in city homeless shelters²² and mental health sites.²⁵ Little work has used the daily interaction with community pharmacists to increase testing, follow-up and treatment adherence. A pilot project introducing DBST in a community pharmacy was undertaken in England in 2009.^{29 30} The pilot concluded that community pharmacies provide a useful route to diagnosing patients with HCV and could successfully reach at-risk groups. The pilot service was most successful when pharmacists worked closely with local drug services and where pharmacists were proactive in discussing risks with clients. Several limitations were noted, including the availability of consultation rooms in pharmacies, the provision of adequate staffing to manage the normal workload of a pharmacy and also the motivation and commitment of the pharmacist to support this new activity. Similar factors were identified in this current study. The provision of clinical standard consultation rooms is now very common in Scotland.

The stigma associated with both OST prescription and HCV infection was strongly recognised in this study. A systematic review of qualitative research into hepatitis testing, recommended framing the positive outcomes of testing in terms of responsibility for individual health and the health of family and community, building positive relationships and targeting stigmatising attitudes.²²

Limitations

The primary limitation of the current feasibility study is the small group of pilot pharmacies. Further work

is now required to establish stronger evidence that pharmacies can positively influence uptake in a greater variety of locations.

The intervention group had a slightly lower average age than the comparator group. Since patients on OST undertake a pathway to recovery, a younger patient group may be expected to have a less stable situation than an older group and be less motivated to engage in healthcare interventions.

Some variance in service delivery is commonplace in pharmacies and in health services generally. The variance may be explained by a range of factors including staffing levels, building configuration, profile of the client group, as well as staff attitudes and the quality of relationships with patients. In this particular study, one pharmacist who was trained to participate in the study, left their post and their work covered by locum staff for the remainder of the study. The rate of testing in this pharmacy was much lower than might have been achieved with consistent staffing. With more experience of delivering training for and services, as well as role

Significant of this study

What is already known on this topic?

- ▶ Less than half of those infected with hepatitis C virus are known to services and rates of uptake of testing and treatment for hepatitis C are low.
- ▶ Dried blood spot testing may increase the uptake of testing for hepatitis C and may be easier to perform than testing by venepuncture.
- ▶ Populations in receipt of opioid substitution therapy (OST) may have a high prevalence of hepatitis C infection.
- ▶ People prescribed OST may visit a community pharmacy regularly to access medication.

What this study adds?

- ▶ Pharmacy staff can be successfully trained to offer dried blood spot testing (DBST) to patients who receive supervised administration of opioid substitution therapy from a pharmacy.
- ▶ The uptake of DBST from community pharmacies may make a useful addition to the identification of the population who are currently infected with hepatitis C.

How might it impact on clinical practice in the foreseeable future?

- ▶ The provision of dried spot testing in community pharmacies may enable greater and more local access to hepatitis C testing.
- ▶ Closer liaison between community pharmacies and hepatology teams may provide additional clinical capacity that can be used to encourage more people infected with hepatitis C to undertake treatment.

acceptance, it is expected that some of the observed variance would diminish.

Interpretation

The study design provided a rapid and simple method of demonstrating that the systems required for DBST testing can be established successfully in community pharmacies. The daily attendance at the pharmacy provides an unexploited opportunity to deliver health interventions. OST recipients usually lived a short walk from their pharmacy.

An increased proportion of OST service users accessed DBST from a pharmacy, compared with other providers. Knowledge of potential issues with implementation should assist with creation of effective service delivery. Further work to evaluate the outcomes associated with this service configuration is required.³¹ The identified barriers to the uptake of testing in this study were overcome by local availability of the pharmacies and positive relationships with pharmacy staff.

Twitter Follow Andrew Radley at @AndrRadl

Reporting Guideline Trend Statement: <https://www.cdc.gov/trendstatement/>

Acknowledgements The authors thank Professor Brian Williams for his contribution towards reviewing this paper. The authors also thank Dr Emma Fletcher and Professor Weiije Wang for providing statistical advice.

Contributors AR: study conception, planning and design, protocol preparation, analysis and interpretation of data, manuscript preparation. KM: acquisition of prescribing data, acquisition of qualitative data, manuscript preparation. JT: acquisition of testing data, analysis of testing data, manuscript preparation. BS: provision of study site staff training, manuscript preparation. JMME: methodological advice, advice on analysis, manuscript preparation. JFD: study conception and protocol preparation, manuscript preparation.

Competing interests AR: Honorarium from Gilead and Research Grants from Gilead and Roche. JT: Honorariums and sponsorship from Abbvie, Bristol Myers Squibb, Gilead, Janssen, Merck Roche Sharp & Dohme and Roche. BS: Honorariums and sponsorship, Gilead and Janssen. JFD: Research Grants, Honorariums and sponsorship from Abbvie, Bristol Myers Squibb, Gilead, Janssen, Merck Roche Sharp & Dohme and Roche.

Funding Gilead Sciences Inc.

Competing interests None declared.

Provenance and peer review Not commissioned; externally peer reviewed.

REFERENCES

- Scottish Intercollegiate Guidelines Network (SIGN). Management of hepatitis C. Edinburgh: SIGN; 2013. (SIGN publication no. 133). [July 2013]. <http://www.sign.ac.uk>
- Hepatitis C annual report: progress made, but much more to do. Public Health England. July 2014. <http://www.gov.uk/government/news/hepatitis-c-annual-report-progress-made-but-much-more-to-do>
- Arain A, Robaey G. Eligibility of persons who inject drugs for treatment of hepatitis C virus infection. *World J Gastroenterol* 2014;20:12722–33.
- Aspinall EJ, Doyle JS, Corson S, *et al.* Targeted hepatitis C antibody testing interventions: a systematic review and meta-analysis. *Eur J Epidemiol* 2015;30:115–29.
- Edlin BR, Kresina TF, Raymond DB, *et al.* Overcoming barriers to prevention, care, and treatment of hepatitis C in illicit drug users. *Clin Infect Dis* 2005; 40(Suppl 5):S276–85.
- Hoofnagle JH, Sherker AH. Therapy for Hepatitis C — the costs of success. *N Engl J Med* 2014;370:1552–3.
- Tait JM, Stephens BJ, McIntyre PG, *et al.* Dry blood spot testing for hepatitis C in people who injected drugs: reaching the populations other tests cannot reach. *Frontline Gastroenterol* 2013;4:255–62.
- McAllister G, Innes H, Mcleod A, *et al.* Uptake of hepatitis C specialist services and treatment following diagnosis by dried blood spot in Scotland. *J Clin Virol* 2014;61:359–64.
- Taheri L. Testing for hepatitis in pharmacies. *Pharm J* 2010;284:51–53.
- Petticrew M, Cummins S, Ferrell C, *et al.* Natural experiments: an underused tool for public health. *Public Health* 2005;119:751–7.
- Craig P, Cooper C, Gunnell D, *et al.* Using natural experiments to evaluate population health interventions: new MRC guidance. *J Epidemiol Community Health* 2012;66:1182–6.
- Jenkins A. Dundee City Council –Scottish Index of Multiple Deprivation 2012 Analysis. http://www.dundeepartnership.co.uk/sites/default/files/SIMD12_Analysis.pdf (accessed 2/2016).
- Midyear population estimate 2014. National Records of Scotland. <http://www.nrscotland.gov.uk/> (accessed Nov 2016).
- Radley A, Melville K, Easton P, *et al.* “Standing outside the Junkie Door”—service users’ experiences of using community pharmacies to access treatment for opioid dependency. *J Public Health* 2016. doi: 10.1093/pubmed/fdw138
- Pope C, Mays N. Reaching the parts other methods cannot reach: an introduction to qualitative methods in health and health services research. *Br Med J* 1995;311:42–5.
- Richie J, Spencer J. Qualitative data analysis for applied policy research. In: Bryman A, Burgess RG, eds. *Analysing qualitative data*. London: Routledge, 1994:173–94.
- Murray E, Treweek S, Pope C, *et al.* Normalisation process theory: a framework for developing, evaluating and implementing complex interventions. *BMC Med* 2010;8:63.
- Dried Blood Spot Testing Guidelines for Hepatitis C virus, Hepatitis B virus and HIV NHS Tayside Sexual Health And BBV Managed Care Network. Dundee 2012. http://staffnet.tayside.scot.nhs.uk/NHSTaysideDocs/idcplg?IdcService=GET_FILE&dDocName=D_OCS_045536&Rendition=web&RevisionSelectionMethod=LatestReleased&noSaveAs=1
- Pawson R, Tilley N. *Realistic evaluation*. London: Sage Publications Ltd, 1997.
- Sahajian F, Bailly P, Vanhems B, *et al.* A randomized trial of viral hepatitis prevention among underprivileged people in the Lyon area of France. *J Public Health* (Oxf) 2011;33:182–92.
- Jones L, Atkinson A, Porcellato L, *et al.* A systematic review of qualitative research on the views, perspectives and experiences of hepatitis B and C testing among practitioners and people at greatest risk of infection. Liverpool Centre for Public Health, 2011.
- Hickman M, McDonald T, Judd A, *et al.* Increasing the uptake of hepatitis C virus testing among injecting drug users in specialist drug treatment and prison settings by using dried blood spots for diagnostic testing: a cluster randomized controlled trial. *J Viral Hepat* 2008;15:250–4.

- 23 Craine N, Parry J, O'Toole J, *et al.* Improving blood-borne viral diagnosis; clinical audit of the uptake of dried blood spot testing offered by a substance misuse service. *J Viral Hepat* 2009;16:219–22.
- 24 Rosenberg S, Goldberg R, Dixon L, *et al.* Assessing the STIRR model of best practices for blood-borne infections in clients with severe mental illness. *Psychiatr Serv* 2010;61:885–91.
- 25 Lindenburg C, Femke A, Lambers F, *et al.* Hepatitis C testing and treatment among active drug users in Amsterdam: results from the DUTCH-C project. *Eur J Gastroenterol Hepatol* 2011;23:23–9.
- 26 Jack K, Willott S, Mannners J, *et al.* Clinical trial: a primary-care-based model for the delivery of anti-viral treatment to injecting drug users infected with hepatitis. *Aliment Pharmacol Ther* 2009;29:38–45.
- 27 Cullen BL, Hutchinson SJ, Cameron SO, *et al.* Identifying former injecting drug users infected with hepatitis C: an evaluation of a general practice-based case-finding intervention. *J Public Health* 2012;34:14–23.
- 28 Harris KA, Van Essen G, Litwin A. Successful integration of hepatitis C evaluation and treatment services with methadone maintenance. *J Addict Med* 2010;4:20–6.
- 29 The Hepatitis C Trust. Diagnosing viral hepatitis in the Community: A 3-month pharmacy testing pilot. London, 2009. <http://www.hepctrust.org.uk/pharmacy-testing-0> (accessed Apr 2016).
- 30 The Hepatitis C Trust. Pharmacy-based hepatitis testing—project review. 2013. London. <http://www.hepctrust.org.uk/pharmacy-testing-0> (accessed Apr 2016).
- 31 NHS Research Authority. Research Summary—SuperDOT-C. <http://clinicaltrials.gov/ct2/show/NCT02706223> (accessed Nov 2016).