Preventable infections are out of control in Canadian hospitals

David Spurgeon *Quebec*

Canadian hospitals are failing to control infections resistant to antibiotics. These kill 8000 patients and cost healthcare systems at least \$C100m (£40m; \$80m; €63m) a year. The Canadian Broadcasting Corporation, in a news investigation, has said that the incidence of methicillin resistant *Staphylococcus aureus* (MRSA) has increased 10-fold in less than a decade (www.cbc.ca/news, 21 March). It also said that since 2003 *Clostridium difficile* has killed more than 600 people in Quebec province alone.

If cases of necrotising fasciitis, the third most common hospital infection, are added, more than 250 000 Canadians are getting ill from preventable infections every year, says the corporation.

Canadian hospitals have fewer regulations for controlling infection than restaurants, says the broadcaster. Many hospitals do not even have the required minimum number of

staff needed to combat infections. The government does not monitor hospitals' infection controls. Restaurants, though, must pass regular inspections or face closure.

Quebec's government has recently announced an increase of \$C20m in funds for hospital infection control. Preliminary findings showed that *C difficile* infections in the province's hospitals had dropped substantially from the previous year (*BMJ* 2005;330:275, 5 Feb).

Andrew Simor, head of the microbiology department at Toronto's Sunnybrook Hospital, says that one study shows that each patient infected with MRSA costs a hospital on average about \$C14 000 (*Infection Control and Hospital Epidemiology* 2001;22:99-104). Projecting this bill across the whole country indicates that the cost of MRSA in Canadian hospitals alone would approach \$C100m a year. Infection control should

be a priority, with a minimum set of standards all hospitals must follow, said Dr Simor.

Donald Low, an infectious disease specialist at Toronto's Mount Sinai Hospital, says that Canada is on the verge of a crisis in controlling preventable infections. He agrees with Dr Simor that hospitals should be forced to meet national standards and report outbreaks to health officials.

In the United Kingdom, an inquiry was launched at Ipswich Hospital, Suffolk, after tests showed that Luke Day, a 2 day old boy who died in February, had septicaemia caused by MRSA. Recent UK government figures showed that there were 3519 MRSA infections between April and September 2004, down from 3574 in the same period in 2003 (www.dh.gov.uk). □

Luke Day, who died from methicillin resistant *Staphylococcus aureus* at Ipswich Hospital, Suffolk, England, aged 36 hours

FDA seeks to ease burden on trial review boards

Bob Roehr *Washington, DC*

The growth in the number and complexity of clinical trials has put great strain on institutional review boards in the United States, say experts in evidence submitted to the US Food and Drug Administration.

The boards were set up to approve and review biomedical research using humans. But the pressures on them, due in part to the large number of sites in trials, are growing to such an extent that a new approach is needed, in particular to recording adverse events, the FDA was told at a public hearing at its headquarters on 21 March.

The meeting was held to find out how the process might be improved to best meet the primary purpose of the review

boards: to protect the rights and welfare of participants in clinical trials.

Greg Koski, a doctor affiliated with Massachusetts General Hospital and a former director of the US Office of Human Research Protections, presented the most detailed and far reaching proposal, in testifying on behalf of the Association of Clinical Research Professionals.

"What we need to do now goes beyond tweaking the existing system," he said. "There is an urgent need and a compelling justification to create and implement a comprehensive national adverse event reporting system that takes advantage of currently available information technologies."

"The best outcome will come from mining a comprehensive database of timely, reliable information. It can and must serve many goals and masters, recognising that not all parties need the same things."

Dr Koski said it was important to keep the collection of data simple, to ensure that the data were collated in the same way and that they were made

available immediately. The system should be based on the internet and be capable of interacting with all types of data processing technologies, he added.

He said the system also should embrace the FDA's Med-Watch programme of recording adverse events associated with drugs once they go on the market. Patients and their families and carers should be encouraged to add entries, he said: "Sometimes the most sensitive observers are the ones who are taking care of mom."

Acknowledging the costs of creating and operating such a system, Dr Koski proposed "a user based national trust fund model analogous to that which supports the national air traffic safety system."

Owen Reese, executive director of the independent Western Institutional Review Board, told the hearing: "The process is plagued with inefficiency and duplication."

The sentiment was shared by all the 19 people who testified at the meeting. Although most boards were willing to shoulder

the burden of examining adverse events that occur at their own site, many complained of being inundated with adverse event reports from other sites in the same trial and sometimes from trials that merely use a similar compound. Boards seldom have either the skills or the resources to evaluate data from external sites.

"Data should not be confused with information. Information is data bestowed with meaning and utility," said Sandra Alfano, vice chairwoman of the independent review board at the Yale University School of Medicine. A strong consensus existed for creating independent data safety monitoring boards for all multicentre trials and for those structures to forward relevant reports in a meaningful context to the relevant independent review boards.

Meanwhile Howard Dickler, speaking for the Association of American Medical Colleges, recommended that the sponsor of the drug or device be made responsible for the preparation of the summary adverse event report. □