tal disorders, such as depression and anxiety, rather than complex psychoses, that contribute most to this rising sickness absence. These are managed almost entirely in primary care, where the focus is on patients with apparently greater clinical needs. Effective evidence based treatments are available for these disorders, including antidepressant medication, problem solving, cognitive behaviour therapy, counselling, and collaborative management.¹⁰ Patients tend to prefer psychological therapies,¹¹ but with a limited capacity to provide them the waiting times are commonly long. Novel approaches to delivery, such as computer based cognitive behaviour therapy, are still at an early stage of development. Both employers and patients require a speedier response than is currently delivered, as the longer an individual remains off work, the more difficult a return to work becomes.

Not uncommonly, a position develops where an individual has recovered sufficiently to consider returning to work but perceives that exposure to his employers, colleagues, or other aspects of work will lead to a relapse. General practitioners can have difficulty linking with employers to effect vocational rehabilitation and, as the patient's advocate, may feel uncomfortable recommending returning to work in this situation. Occupational physicians are best equipped to manage these cases, yet the United Kingdom has very poor provision of occupational health (one specialist for every 43 000 workers) compared with the rest of Europe.¹ A cluster randomised controlled trial in Holland has shown how early psychological interventions for common mental disorders, delivered through the workplace, can enhance health and reduce absence.¹² The intervention consisted of 4-5 sessions of cognitive behaviour therapy to increase activity and coping skills for those off sick for only two weeks. It reduced total sick leave, time taken to return to work, and recurrence at 12 months. If the government is serious about tackling the consequences of common mental disorders then innovative policies, possibly requiring major expansion in occupational health and provision of psychological

therapy service in primary care, will be required alongside research into the most effective and cost effective methods of delivering service. This would be a wise investment given the substantial economic and social costs engendered by the current service framework.

Max Henderson *clinical research fellow in liaison psychiatry*

Institute of Psychiatry, Department of Psychological Medicine, Weston Education Centre, London SE5 9RJ
(m.henderson@iop.kcl.ac.uk)

Nicholas Glozier *consultant occupational psychiatrist*

Department of Occupational Health and Safety, King's College Hospital NHS Trust, London SE5 9RS

Kevin Holland Elliott *professor of occupational health and health risk management*

Brunel University, Uxbridge, Middlesex UB8 3PH

Competing interests: NG has received speaking and consulting fees from private and public employers who may benefit from a greater return to work of trained employees. KHE is chair of Corporate Health and Performance, a charity established to research the link between health, productivity and performance at work.

- 1 Confederation for British Industry. *Room for improvement: Absence and labour turnover (in association with AXA)*. London: CBI, 2004.
- 2 Unum Limited, Institute for Employment Studies. *Towards a better understanding of sickness absence costs*. Dorking: Unum, 2001.
- 3 Department for Work and Pensions. *Pathways to work: Helping people into employment*. London: Stationery Office, 2002.
- 4 Department of Health. *Choosing health: making healthier choices easier*. London: DH, 2004.
- 5 Health and Safety Executive. *Securing health together*. London: HSE, 2000.
- 6 Waddell G, Aylward M, Sawney P. *Back pain, incapacity for work and social security benefits: an international literature review and analysis*. London: Royal Society of Medicine Press, 2002.
- 7 Jones J, Huxtable C, Hodgson J, Price M. Self-reported work-related illness in 2001/2002. London: HSE, 2003.
- 8 Singleton N, Bumpstead R, O'Brien M, Lee A, Meltzer H. *Psychiatric morbidity among adults living in private households*. London: Office for National Statistics, 2000.
- 9 Health and Safety Executive. *Tackling work-related stress. A manager's guide to improving and maintaining employee health and well-being*. London: HSE, 2001.
- 10 Dowrick C. Advances in psychiatric treatment in primary care. *Adv Psychiatric Treat* 2001;7:1-8.
- 11 Bedi N, Lee A, Harrison G, Chilvers C, Dewey M, Fielding K et al. Assessing effectiveness of treatment of depression in primary care: partially randomised preference trial. *Br J Psychiatry* 2000;177:318.
- 12 Van der Klink J, Blonk R, Schene A, van Dijk F. Reducing long term sickness absence by an activating intervention in adjustment disorders: a cluster randomised controlled design. *Occup Environ Med* 2003;60:437.

New professional roles in surgery

Would be effective in selected surgical settings and can offer benefits

New professional roles in surgery are a controversial issue. Recent publicity surrounding surgical care practitioners has illustrated the extent of hostility in parts of the surgical community.¹⁻³ Yet the landscape of the NHS is changing radically. External forces such as the European Working Time Directive are having a profound effect on the United Kingdom's healthcare workforce, and maintaining the current situation is not an option.^{4 5}

The unavoidable reality is that we do not have enough doctors to sustain traditional working patterns. Therefore, developing new professional roles seems a logical response. Moreover, role redesign fits with the government's commitment to widen career opportunities in health care and to develop a flexible training

structure based on individual competences rather than traditional pigeonholes such as doctor or nurse.^{6 7}

Increasing numbers of medically unqualified practitioners are now being trained in surgery related practice, and this is a good time to examine the pros and cons. We write from the perspective of a large university teaching hospital in central London, with a track record of pioneering new roles. Early projects included establishing the United Kingdom's first nurse consultant in coloproctology and a nurse led minor surgery service in west London. Although now widely accepted, these roles aroused great opposition when first introduced.

More recently we have led two national pilot programmes, funded by the Department of Health and

drawing participants from nursing, the operating department, and allied health practitioners. The perioperative specialist practitioner programme provides preoperative and postoperative care, working alongside junior medical staff and as a part of the surgical team.⁸ Participants in the Imperial surgical care practitioner pilots carry out surgical procedures, working under supervision as part of a clinical team. Other surgical care practitioner pilots have taken place elsewhere in the United Kingdom.

Our pilots show that intensive focused training can lead to high degrees of expertise in a relatively short time (one to two years), albeit within clearly defined limits. In our unpublished study, detailed evaluation, using extensive interviews by independent qualitative researchers, showed the high perceived value of these roles in many of the 22 NHS trusts that took part. Equally striking, however, were the high levels of initial anxiety and mistrust that emerged, especially among junior doctors, who felt threatened by changes to traditional working patterns. Interestingly, support for new roles was greatest outside the metropolitan teaching centres, with their traditional reliance on trainees.

At first sight, the advantages of practitioners in new roles in surgery seem obvious. Provided they are suitably trained and supervised they can provide a much needed addition to the workforce. Practitioners in these new roles will not rotate, unlike junior doctors, so continuity will improve. Direct referral pathways from primary care can be developed. By carrying out surgical and perioperative tasks such as excision of skin lesions and preanaesthetic assessment according to clearly defined protocols, such practitioners can allow surgeons to focus on managing more complex clinical problems. In time they could act as a resource for junior surgeons as they learn straightforward procedures. This would free consultants to teach more complex tasks.

However, there are caveats. These new practitioners should not be embraced uncritically or introduced solely as a response to political imperatives. Redesign of surgical roles is relatively new in the United Kingdom, and most information comes from pilot projects whose conditions may be unrepresentative. Moreover, most trainee practitioners for new roles have been experienced healthcare professionals, drawn from other parts of the existing workforce. If role design is to do more than simply transfer staffing shortages from one part of the system to another, the new practitioners must also be recruited from outside the NHS. But little is known about training direct entrants to carry out specialised surgical tasks.

A subtler issue relates to professional expertise. Much routine operative and perioperative practice is

repetitive and underpinned by clinical protocols. Practitioners in new roles have much to offer in this area. But beyond the boundaries of the routine, when clinical presentations are atypical and confusing, the skills of experienced consultants are essential. Such wide ranging expertise takes years to acquire and, although easily recognised, is hard to define.⁹ Yet if surgical practice becomes dominated by narrowly defined roles, future generations of surgeons may lose the mature expertise that allows them to recognise and manage difficult clinical challenges. If that happened we would lose something very valuable.

We believe that practitioners in new roles can be effective in selected surgical settings and can offer noteworthy benefits. Opponents, however, fear a diminution in training opportunities for doctors and see a threat to established lines of clinical responsibility. In our view, the solution is not to oppose the development of practitioners in new roles, for they offer great potential in a rapidly changing health service. Rather we should support what such roles can offer by ensuring that they meet the highest clinical standards. For this to succeed, practitioners in new roles need to be part of a wider national framework, where high quality care delivered by medically unqualified practitioners is combined with the best elements of traditional consultant led surgical practice.

Roger Kneebone *senior lecturer in surgical education*
Ara Darzi *head*

Department of Surgical Oncology and Technology, Imperial College London, St Mary's Campus, London W2 1NY
(r.kneebone@imperial.ac.uk)

Competing interests: None declared.

- 1 Laurence J. NHS revolution: nurses to train as surgeons. *Independent* 6 Dec 2004. http://news.independent.co.uk/uk/health_medical/story.jsp?story=590105 (accessed 17 Feb 2005).
- 2 A closed shop for surgery does patients no favours. *Independent* 6 Dec 2004. http://comment.independent.co.uk/leading_articles/story.jsp?story=590065 (accessed 17 Feb 2005).
- 3 Mulholland H. Nurses "to carry out routine operations." *Guardian* 6 Dec 2004. <http://society.guardian.co.uk/NHSstaff/story/0,7991,1367616,00.html> (accessed 17 Feb 2005).
- 4 Phillips H, Cripps J. EWTD developments. *Bull R Coll Surgeons Engl* 2004;6:190-2.
- 5 The Royal College of Surgeons of England and the European Working Time Directive. http://www.rcseng.ac.uk/service_provision/ewtd/ (accessed 22 Feb 2005).
- 6 Buchan J, Calman L. Skill mix and policy change in the health workforce: nurses in advanced roles. OECD health working papers number 17. Paris: OECD Publications Service, 2004. www.oecd.org/dataoecd/30/28/33857785.pdf (accessed 17 Feb 2005).
- 7 Mahon A, Harris C, Walshe K, Higgins J. The European Working Time Directive and doctors in training: an early evaluation of 19 NHS pilot. Report by Manchester Centre for Healthcare Management, University of Manchester, 2003.
- 8 NHS Modernisation Agency. Working time directive pilots programme. www.modern.nhs.uk/scripts/default.asp?site_id=50 (accessed 16 Feb 2005).
- 9 Talbot M. Good wine may need to mature: a critique of accelerated higher specialist training. Evidence from cognitive neuroscience. *Med Educ* 2004;38:399-408.