was commissioned by the Department of Health from UnitedHealth Europe, the company that piloted the programme together with nine NHS primary care trusts in England over 17 months. The pilots covered 1333 patients.

The Evercare programme identifies vulnerable elderly patients, who are then allocated to an advanced practice nurse who manages their cases and coordinates all aspects of the patients' care, working with GPs, geriatricians, each patient's family, and social services. The nurse also advises patients in managing their own care.

The report, published last week, looks at the initial outcome of the pilot scheme, which was introduced into the trusts from April 2003.

Among the 793 patients who were identified as having had previous unplanned admissions, unplanned admissions fell by 38% (from an annualised rate of 174 per 100 before the scheme was introduced to an annualised rate of 108 per 100).

In a smaller group of 183 patients, who were predicted by general practitioners to be at high risk of admission, admissions rose from an annualised rate of 21 per 100 to 62 per 100. Adrian O'Dowd *London*

Assessment of the Evercare Programme in England in 2003-2004: Final Report is available at www.united-healthcare.co.uk/ouractivities/evercare.html

Radiation for prostate cancer increases risk of rectal cancer

Men with prostate cancer treated with external beam radiation have a 70% higher risk of developing rectal cancer than men who have surgery, a US study has found. Radiation is now used for about 17% of patients with prostate cancer.

The retrospective population based study used data from the US National Cancer Institute's surveillance, epidemiology, and end results (SEER) register, from 1973 to 1994, and

included more than 85 000 men aged 18 to 80. The participants had had invasive microscopically confirmed prostate cancer but no evidence of metastases or previous history of colorectal cancer (*Gastroenterology* 2005;128:819-24).

After controlling for other factors, the results showed that radiation was a significant risk factor for rectal cancer ($P<0.0001$), with a hazard ratio of 1.7 (95% confidence interval 1.4 to 2.2). This is equivalent to the same level of risk as having a first degree relative who has had colon cancer.

"Overall, the risk is still quite low, about 5 in 1000 for those treated with surgery and 10 in 1000 for those treated with radiation," she said.

In all, 55 263 men had had surgery and 30 552 had been treated with radiation. The average age was 67.6, and the men given radiation were on average about two years older than those who had surgery.

Janice Hopkins Tanne *New York*

Young people who binge drink are likely to continue into adulthood

Binge drinking is widespread among British men and women throughout their 20s, 30s, and into their 40s, new research shows (*Addiction* 2005;100:543-9). And those who binge drink in their early 20s are more likely to be binge drinking 26 years later: the risk is twofold for men and 1.5-fold for women.

Men who drank more than seven units a week at the age of 16 were one and a half times more likely than light drinkers to binge drink in their 30s or 40s. Women who rarely or never drank at this age were less likely than light drinkers to binge drink later on.

Binge drinkers were defined as men who consumed 10 or more units of alcohol on each occasion and women who consumed seven or more units. One unit is 8 g or 10 ml of pure alcohol, equivalent to a small glass of wine or half a pint of beer.

Roger Dobson *Abergavenny*

Global ecological disaster predicted in next 50 years

Owen Dyer *London*

Humanity is heading for ecological disaster if instead of foreseeing and preventing environmental degradation we just react to it. This is the conclusion of a United Nations report compiled by 1300 leading scientists from 95 countries.

The *Millennium Ecosystem Assessment Synthesis Report* previews the ecological state of the world in 2050. It lists 24 essential "ecosystem services," such as timber, clean air, and fresh water, and finds that 60% of them are being degraded or used unsustainably.

This degradation obstructs the UN Millennium Development Goals, set in 2000, the most ambitious of which was a halving of the world's population existing on less than a \$1 (£0.53; €0.78) a day or threatened by hunger or lack of clean water. Among the gravest threats to the environment are excessive "nutrient loading" from agricultural fertilisers and the progressive disappearance of biodiversity. The extinction rate of species is already a thousand times higher than the average rate shown by the fossil record and is set to increase 10-fold in the next 50 years.

"Humans are fundamentally and to a significant extent irreversibly changing the diversity of life on earth," says the report, noting that this could harm pharmaceutical research and development.

The scientists warn of possible "accelerating, abrupt, and potentially irreversible changes."

These include the collapse of fish stocks, such as North Sea cod; rapid growth of marine algae, creating oxygen depleted dead zones in the sea; and emergence of disease.

In Africa, growing pressure on water supplies combined with regional climate change has the potential to greatly enlarge the areas in which cholera is a threat. The range of malaria bearing mosquitoes is also likely to increase.

The release of carbon into the atmosphere and the leaching of nitrogen based fertilisers into water pose serious risks for chronic disease, the report finds. Increases in ultraviolet B radiation, ozone and other air pollution, and pollen production are also predicted.

The report gives four sets of predictions for 2050, based on the approach currently adopted to counter ecosystem degradation. "The scenario we are closest to now is the one we call order from strength," said Dr Reid. This describes a world in which international cooperation is lacking, economic competition is fierce, and environmental threats are faced only after they emerge. This is likely to deliver the fastest population growth, the lowest economic growth, and the most environmental damage of the four scenarios, he said. □

The *Millennium Ecosystem Assessment Synthesis Report* is available at www.millenniumassessment.org/en/Products.Synthesis.aspx.

Slash and burn agriculture is destroying this rain forest in Cameroon