*Interactive case report***Postoperative hypoxia in a woman with Down's syndrome: case presentation**

A K Siotia, A Chaudhuri, S I Muzulu, D Harling, R Muthusamy

Northern General Hospital, Sheffield, S5 7AU

A K Siotia
research fellow in
cardiologyNinewells Hospital,
DundeeA Chaudhuri
specialist registrar in
acute medicineRotherham District
General Hospital,
Rotherham, S60
2UDS I Muzulu
consultant
endocrinologistD Harling
consultant
anaesthetistR Muthusamy
consultant
cardiologist

Correspondence to:

A K Siotia
Anjan.Siotia@
sth.nhs.uk

BMJ 2005;330:834

This is the first of a three part case report where we invite readers to take part in considering the diagnosis and management of a case using the rapid response feature on bmj.com. Next week we will report the case progression and in four weeks' time we will report the outcome and summarise the responses

Results of patient's preoperative blood tests

	Value	Normal range
Haemoglobin (g/l)	121	115-165
White blood cells ($\times 10^9/l$)	17.6	4.0-11.0
Neutrophils ($\times 10^9/l$)	14.5	2.0-7.5
Platelets ($\times 10^9/l$)	184	150-400
Sodium (mmol/l)	141	136-145
Potassium (mmol/l)	3.6	3.5-5.0
Urea (mmol/l)	3.9	2.8-7.4
Creatinine ($\mu\text{mol/l}$)	107	55-96
Total protein (g/l)	67	60-80
Albumin (g/l)	25	34-46
Bilirubin ($\mu\text{mol/l}$)	10	3-20
Alanine aminotransferase (IU/l)	12	9-56
Gamma glutamyltransferase (IU/l)	14	7-33
Alkaline phosphatase (IU/l)	132	100-300
Arterial blood gases:		
pH	7.40	7.35-7.45
Carbon dioxide partial pressure (kPa)	6.46	4.5-6.1
Oxygen partial pressure (kPa)	6.26	12-15
Base excess	-4.8	0-2

Miss Webb, a 24 year old woman with Down's syndrome, was referred by her general practitioner to the gynaecology outpatient clinic. She gave a one year history of heavy and irregular periods, general lethargy, and a painful abdominal lump, which examination showed to be an umbilical hernia. She had no serious medical history (including congenital heart disease) and she was not taking any regular medicine. She lived with her mother, who was her main carer.

Attempts at insertion of a levonorgestrel intrauterine system (Mirena) under general anaesthesia to treat her menorrhagia were unsuccessful. She subsequently had a dual operation to repair a painful umbilical hernia and a total abdominal hysterectomy under general anaesthesia. Preoperative blood tests including full blood count, urea and electrolytes concentrations, liver function tests, and coagulation profile all gave normal results.

Questions

- 1 What are the ethical considerations around obtaining Miss Webb's consent for a hysterectomy and what would be best practice here?
- 2 What are the differential diagnoses for the sudden postoperative deterioration in Miss Webb's condition?
- 3 What further investigations would you do?
- 4 What earlier intervention may have prevented this deterioration?

Please respond through bmj.com, remembering that Miss Webb is a real patient and that she and her carers will read your response

Immediately after the operation, she was slow to recover from the anaesthetic. She was hypoxic with poor respiratory effort. She was given intravenous naloxone for suspected opiate induced hypoventilation, but her condition was unchanged. She required respiratory support in the intensive therapy unit. She was taken off the ventilator after 24 hours, but had slow respiratory effort, and despite high flow oxygen her oxygen saturation was only 93%. On examination she was alert with a Glasgow coma score of 15, respiratory rate of 10 breaths/minute, heart rate of 70 beats/minute, regular rhythm, systolic blood pressure of 98 mm Hg, and temperature of 37°C. Examination of the cardiovascular system, chest, abdomen, and nervous system showed no abnormality. The table shows the results of the blood tests. Portable chest radiography (AP view) showed normal lung fields with possibly increased cardiothoracic ratio.

Competing interests: None declared.

Submitting articles to the BMJ

We are now inviting all authors who want to submit a paper to the *BMJ* to do so via the web (<http://submit.bmj.com>).

Benchpress is a website where authors deposit their manuscripts and editors go to read them and record their decisions. Reviewers' details are also held on the system, and when asked to review a paper reviewers will be invited to access the site to see the relevant paper. The system is secure, protected by passwords, so that authors see only their own papers and reviewers see only those they are meant to.

Anyone with an internet connection and a web browser can use the system.

The system provides all our guidance and forms and allows authors to suggest reviewers for their paper. Authors get an immediate acknowledgment that their submission has been received, and they can watch the progress of their manuscript. The record of their submission, including editors' and reviewers' reports, remains on the system for future reference.

The system itself offers extensive help, and the BMJ Online Submission Team will help authors and reviewers if they get stuck.

Benchpress is accessed via <http://submit.bmj.com> or via a link from bmj.com