Published in final edited form as:

Curr Environ Health Rep. 2016 September ; 3(3): 169–177. doi:10.1007/s40572-016-0102-3.

A Framework to Address Challenges in Communicating the Developmental Origins of Health and Disease

Dr. Liana Winett, DrPH, MPH, MCHES,

School of Community Health and OHSU/PSU School of Public Health, Portland State University,
PO Box 751, Portland, OR 97201

Dr. Lawrence Wallack, DrPH, MPH, MS,

School of Community Health and OHSU/PSU School of Public Health, Portland State University,
PO Box 751, Portland, OR 97201

Dr. Dawn Richardson, DrPH, MPH,

School of Community Health and OHSU/PSU School of Public Health, Portland State University,
PO Box 751, Portland, OR 97201

Dr. Janne Boone-Heinonen, PhD, MPH, and

Public Health and Preventive Medicine and OHSU/PSU School of Public Health, Oregon Health
and Science University, 3181 Sam Jackson Park Road, Portland, OR 97239

Dr. Lynne Messer, PhD, MPH

School of Community Health and OHSU/PSU School of Public Health, Portland State University,
PO Box 751, Portland, OR 97201

Abstract

Findings from the field of Developmental Origins of Health and Disease (DOHaD) suggest that some of the most pressing public health problems facing communities today may begin much earlier than previously understood. In particular, this body of work provides evidence that social, physical, chemical, environmental, and behavioral influences in early life play a significant role in establishing vulnerabilities for chronic disease later in life. Further, because this work points to the importance of adverse environmental exposures that cluster in population groups, it suggests that existing opportunities to intervene at a population level may need to refocus their efforts “upstream” to sufficiently combat the fundamental causes of disease. To translate these findings into improved public health, however, the distance between scientific discovery and population application will need to be bridged by conversations across a breadth of disciplines and social roles. And importantly, those involved will likely begin without a shared vocabulary or conceptual starting point. The purpose of this paper is to support and inform the translation of DOHaD findings from the bench to population-level health promotion and disease prevention, by: 1)

*Corresponding Author: Liana Winett, lwinett@pdx.edu, Tel. 503-725-8262, Fax 503-725-5100.

Compliance with Ethics Guidelines

Conflict of Interest

Liana Winett, Lawrence Wallack, Dawn Richardson, Janne Boone-Heinonen, and Lynne Messer declare that they have no conflict of interest.

Human and Animal Rights and Informed Consent

This article does not report any new studies with human or animal subjects performed by any of the authors.

Discussing the unique communication challenges inherent to translation of DOHaD for broad audiences, 2) Introducing the First-hit/Second-hit Framework with an epidemiologic planning matrix as a model for conceptualizing and structuring communication around DOHaD, and 3) Discussing the ways in which patterns of communicating DOHaD findings can expand the range of solutions considered, and encourage discussion of population-level solutions in relation to one another, rather than in isolation.

Keywords

Developmental Origins of Health and Disease (DOHaD); Epigenetics; Message Framing; Mother-blame; Community-blame; First-hit/Second-hit Framework

Introduction

There is burgeoning scientific evidence that social, physical, chemical, environmental, and behavioral influences in early life play a significant role in establishing vulnerabilities for chronic disease later in life [1–6]. Much of this work has been collected in the field now referred to as of the Developmental Origins of Health and Disease (DOHaD) [2–4]. The mechanisms for disease vulnerability this field reveals have been linked to obesity, heart disease, diabetes, and some cancers, in addition to cognitive development, educational performance, and mental health issues [7–14]. These changes result, in part, from adverse exposures to the reproductive environment and developing organ systems during – and even prior to – gestation, as well as during the very early days of childhood [15–19]. A key mechanism by which these influences exert themselves is epigenetics. This has been described as “the biochemical process through which some genes are expressed and others remain silent, and it reinforces and explains the powerful impact that the environment has on human development. Epigenetic effects occur not only through diet, chemical exposure, and high levels of environmental stress, but also through chronic poverty and racism” [In 20, also 21–22]. As such, the research that comprises the field of DOHaD indicates that much chronic disease results less from genetic predisposition than from genetic responses to environmental stressors; this is to say that the “epigenetic process of gene regulation is where nature and nurture come together” [23].

It may be, then, that some of the most pressing public health problems facing communities today begin much earlier than previously understood, and that existing opportunities to intervene at a population level need to refocus “upstream” to sufficiently combat the fundamental causes of disease. To translate DOHaD into improved public health, however, the distance between scientific discovery and population application will need to be bridged by conversations across a breadth of disciplines and social roles. And, it is important to recognize that this breadth means that those involved will likely begin without a shared vocabulary or conceptual starting point.

This is a definitional moment for those scientists, public health professionals, policy makers, community members, and others working to move DOHaD findings from the bench to population-level health promotion and disease prevention. The purpose of this paper is to support the extension of the science into applied arenas by:

1. Providing an introduction to communication framing, in general, and challenges in communicating DOHaD, in particular.
2. Introducing the First-hit/Second-hit framework with a planning matrix as a model for conceptualizing and structuring communication around DOHaD, and as a beginning step toward addressing some of the inherent framing challenges.
3. Discussing the ways in which patterns of communicating DOHaD can expand the range of solutions considered; moreover, how – by cutting across traditional professional and academic boundaries – the framework may encourage discussion of solutions in relation to one another, rather than in isolation.

We conclude by offering recommendations for the application of the communication planning tool to the translation of DOHaD findings for broad audience engagement.

The DOHaD Communication Task

Those seeking to bridge DOHaD research and practice encounter a communication task fraught with difficulty [24]. First, having emerged over the past three decades, this is a relatively new body of science that is undergoing constant clarification and is marked by great complexity. For many, it represents a very different way of conceptualizing the onset of disease risk. Second, the specific vocabulary that animates this science (e.g., “programming,” “wiring,” “embedding,” “imprinting,” “poor control”) and the range of language used to describe the “quality” of a person’s physical constitution can prove problematic when used in non-scientific arenas [24].

Perhaps the most vexing of the communication problems marking this field, and those on which we focus in this paper, relate to the unintended consequences of reporting causal associations. Because DOHaD suggests that disease vulnerability can be rooted in any combination of factors ranging from the individual-level to those established in social, political, economic, and physical environments, there is the risk that in discussing one key aspect in isolation from the total picture we may present a distorted view of the complete range of opportunities to effect change. And as we will discuss, in so doing we also run the risk of inadvertently holding at fault those who are at greatest risk.

The Importance of Communication Frames

When researchers and others communicate about the nature and urgency of DOHaD, they do so through what sociologists and cognitive scientists call “frames”¹ [25–29]. Frames are the ideological structures, grounded in cultural values, that people use to process and understand the external cues they perceive [30]. Frames tell us which features are within a problem’s parameters, and which are outside of it [29, 31]. Because they characterize a problem as being of a particular type (and thus not another), frames also tell us literally and by abstraction what the problem is, why it matters, what can be done about it, and who is responsible [31]. Importantly, cognitive science demonstrates that the frames we use to interpret problems lead us to particular understanding of the range of solutions available, and may also functionally limit our ability to consider possibilities outside of that range [28].

Frames can be invoked by any cue (e.g., words, metaphor, imagery, core arguments, appeals to values) that leads us to filing an issue into a particular mental category [32]. The importance lies in the articulated meaning that accompanies those cues. For example, if we frame a problem as a “public health issue,” we are suggesting that it is important because of its population health impact; that it is amenable to the tools, techniques, and actions of public health professionals; and that there likely will be an emphasis on primary prevention through broad population-level action. If we were to frame that same problem as a “criminal justice issue,” by contrast, we would suggest its importance and urgency are rooted in concern for law and order; that it is amenable to the tools, techniques, and actions of law enforcement professionals; and that there likely will be an emphasis on deterrence and punishment. And, if we were to say that same problem was a “behavioral issue,” we would be suggesting that it is important because of the effects on the person involved (or the effects that person has on others), that the persons responsible are those exhibiting the behavior, and that the likely tools available are those that sculpt individual habits and practices [e.g., 33]. Thus, by acting as the cognitive structures and filters with which we organize and process the information we receive, frames also help us consider the range of possible action going forward.

Two DOHaD Framing Challenges

The Risk of Mother-blame—As we and others have noted, there is a tendency in communicating DOHaD to frame it disproportionately around the biology, behaviors, and life circumstances of the most proximal link in the gestational chain: the mother [24, 34–36]. Common experimental designs evident within the field of DOHaD may contribute to this frame, as they typically compare health outcomes across proximal exposures -- the mother’s constitution, her pregnancy experience, and her actions during or prior to it. When associations are found, the public spotlight tends to focus on these same maternal factors unless and until subsequent research and/or more expansive framing situates initial findings in their broader contexts. Communicating across that divide is part of the challenge.

We have identified three additional potential contributors to the tendency toward mother-blame in DOHaD communications [24]. First, cognitive linguistics explains that people’s minds create unavoidable associations, or “cognitive maps,” between concepts nested within categories. [37–39]. Because “pregnancy” is a category including “fetus + mother,” this cognitive map may foster a predisposition toward conceptualizing gestational problems from the narrow perspective of the maternal environment. Second, much of the language that commonly characterizes discussion of the science (e.g., “her pregnancy,” “the intrauterine environment she provides,” “the mother’s prolonged stress,” “if the baby had been poorly nourished prior to birth”) pinpoints epigenetic phenomena within, or more directly, as being *of* the mother [24, 34, 36]. Third, there is a dominant personal responsibility frame anchoring much of Western narrative, and thus also shaping our mental processing of problems and solutions [40–43]. This readily accessed cognitive map can predispose us to understand that health risks established during gestation are primarily the product of the mother’s choices and behaviors, or at very least a function of her constitution. While we acknowledge that other frames exist that do not hold mothers primarily culpable for the

health of their offspring, cognitive science tells us that extra cognitive work is required for such less dominant frames to rise above the dominant maps and guide our thinking. [39].

The importance of mothers in the lives of their infants and children is without question. Yet to circumscribe the extent of the problem and solutions to their bodies and behaviors excludes the critical influences and constraints of the broader societies in which women are pregnant and children develop and grow. A robust translation and integration of DOHaD findings into applied health promotion/disease prevention work requires highlighting the less readily accessed cognitive maps that connect individuals to the environments in which they exist.

The Risk of Community-blame—At the same time communicators of the science underlying DOHaD run the risk of an inadvertent and generalized blaming of mothers, there is a second potential risk to be aware of that is similarly problematic but perhaps less obvious – and, indeed, may emerge in reaction to the first. This is that in seeking to avoid the threat of mother-blame our response may be to instead focus our attention on the contexts in which women reside, including their neighborhoods and communities. Such refocusing is important, and reflects the public health orientation to population-level understanding and intervention. Yet, if we settle on the community as the point of concern without linking it to the larger societal forces within which its circumstances come to be and are sustained, we can also simultaneously diminish the very communities needing the most support. Thus, communicators of DOHaD must also be alert to the risk for community-blame that can directly result from efforts to overcome mother-blame.

Community-blame can lead to a kind of categorical stigmatization [44–48]. Such “degrading marks” on communities evolve through characterizations casting them as undesirable places without resources, capacity, or inherent value [44, 49]. These portrayals are further perpetuated through media coverage and academic discourse, which variously describe disadvantaged communities as “poverty-stricken,” “crime-ridden,” “vulnerable,” and “needy” [50]. Characterizations such as these can create a double jeopardy for groups already dealing with significant health challenges, as they can send the unintended message that the very communities most in need of social and economic investment are instead hopeless and beyond repair.

Further, focusing on decontextualized community factors may limit our effectiveness in addressing DOHaD-related problems. First, we may miss important root causes of disparate health outcomes -- the “causes of the causes,” or the powerful social forces that result in differential risks borne across social groups [in 51, also 52–53]. Second, by missing the macro dimensions of the problem, we may fail to engage the range of stakeholders required for robust discussion of solutions [54]. Finally, messages that are perceived to tarnish communities may (understandably) invite resistance from members of the very communities we seek to help [54–55].

Community level influences are important in the ecology of DOHaD phenomena and are a critical part of the conversation about solutions. And yet -- as is the case with mother-blame

-- we risk holding community members accountable for the circumstances in which they find themselves, which they neither independently created nor are well-positioned to change.

Orienting DOHaD Communication

We have argued that to avoid misplaced blame, DOHaD phenomena must be articulated within the context of the systems that give rise to them. Here, we present a systems model that we believe may provide a first step toward structuring more comprehensive and inclusive communications. It is important to note that developing frames, and ultimately messages, involves far more than the core perspective around which they are crafted [56]. But, articulating that core perspective is the first step [56–57].

The model we propose draws from two complementary public health frameworks. The first is the First-hit/Second-hit framework, previously articulated in the “Two-hit hypothesis” of cancer [58–59], etiology of schizophrenia, [60–61] automobile injury prevention [62], and recently in explaining processes underlying DOHaD [63–64]. The second is a planning tool originally developed to support a comprehensive public health approach to injury prevention, which we adapt to demonstrate the interrelationships among first-hit/second-hit variables [65]. We explain, below, how we feel each informs the DOHaD communication work ahead. Importantly, we believe that these tools allow us to tell the story of the systems and systems breakdowns that give rise to DOHaD risk, and in so doing, help to avoid mother- and community-blame pitfalls.

First Hit/Second Hit Framework

The First- and Second-hits framework provides a means to explain how lifelong health can be adversely affected by a series of “hits,” or insults, experienced at critical developmental intervals and across the lifespan. These insults can have enduring effects both on an individual’s predisposition to disease, as well as on the disease predispositions of subsequent generations [63–64].

“First hits” in DOHaD are adverse exposures experienced by the mother that induce disease vulnerabilities prior to birth. Examples of first hits studied include insufficient nutrient flow during gestation, high calorie malnutrition, environmental toxicants, and stress hormones [7, 22, 23, 66–70]. For example, inadequate nutritional flow has been associated with enduring disease-predisposing alterations in the heart, kidneys, and liver of the developing fetus [71–72]. These adverse effects persist across the lifespan and translate to increased risk for diabetes, hypertension, and heart disease. Gestational exposures to environmental toxicants or stress hormones also have been associated with similar adverse effects on fetal development and subsequent disease risk [73–78]. Importantly, each of these first hit exposures may be initiated, perpetuated, magnified, or limited by the broader physical and social contexts in which people exist.

“Second hits,” then, are additional exposures encountered by offspring after birth and throughout their lives. Second hits take many forms, but the most extensively researched include nutrition, social stress, and environmental toxicants [73–78]. Second hits can activate or exacerbate the initial biological vulnerabilities developed as a result of first hits.

That is, for those who were primed during gestation for potential onset of disease, second hits may further heighten vulnerabilities to poor health and adult illness.

Importantly, the occurrence of first- and second-hits are not random, but rather cluster along the same social, racial, and economic stratifications that divide communities [79–86]. Therefore, those most likely to experience first hits are also at the highest risk of exposure to second hits. And, because of intergenerational transfer of risk in DOHaD, the “second hits” of the current reproductive age generation can become the “first hits” of the next. For example, due to the effects of racial and/or socioeconomic segregation, children born to pregnant women living in neighborhoods with significant social disadvantage are likely to remain in the same or similar neighborhoods as they age. This sets the stage for not only initial and subsequent vulnerabilities in this first generation of offspring, but as they reach reproductive age, also provides conditions in which the enduring environmental stressors can be mediated through them to establish vulnerabilities in the next generation. It is through this cycle that persistent health disparities can be perpetuated and magnified within communities: this cycle lies at the intersection of biological vulnerability (first hits), modifiability of risk (second hits), and social vulnerability (resources/ability to modify risk).

The First- and Second-hits Framework, by detailing the staged, multi-level, and mutual influences in DOHaD, offers important structure to communicating the science. In particular, it: 1) explicates the critical interconnections among a broad scope of chronological influences that communicators need to be prepared to articulate, and 2) provides a model with which to illustrate the importance of lifelong social, environmental, and physical contexts in the likelihood of disease.

First-hit/Second-hit Planning Matrix

We draw from the Haddon Matrix, an innovative approach developed for injury prevention that applies epidemiological principles to the understanding of non-infectious health outcomes [65, 87–88]. It has subsequently been expanded to a variety of public health issues [e.g., 89–92]. In particular, the matrix aligns the epidemiologic triad of host, agent, and environment with opportunities for prevention in the pre-event, event, and post-event phases of a health phenomenon. To reflect the increasing focus on place-based interventions in public health -- and because a particular communication challenge in speaking to this body of science is adequately incorporating the breadth of context -- we have adapted this tool to further specify the multiple nested physical and social environments in which effects occur. Table 1 illustrates the use of the tool, using examples of the types of DOHaD influences that may populate it.

We believe the matrix is particularly well suited to illustrating and speaking to the causal factors identified through the First- and Second-hits framework because it specifies the chronological and systemic nature of influences. In addition, the matrix provides visual structure to the idea of staged “hits,” and it places the people affected along the continuum of influences so that we can also describe what influences them. Further, by drawing connections to the larger contexts in which risk develops, it allows communicators to speak to any one factor under study without excluding other critical influences. The matrix

provides a structure for planning deeper communication among broad constituencies, by emphasizing interconnections among factors in the cells.

Using the First-hit /Second-hit Framework as Conceptual and Communication Model

In using this matrix (Table 1), scientists, public health professionals, policy makers, and community members, together, can place each causal factor or solution being discussed in relation to the full range of strategies required to meaningfully move the needle on health promotion and disease prevention. For example, maternal nutrition programs can be discussed in relation to food access, affordability, and safety. Prenatal stress management programs can be situated in the context of the need for economic stability, universal living wage, and broad based social opportunity. And, the need for quality, affordable prenatal and pediatric health care can also be described alongside the equally important needs for safe and secure housing that is free of toxic substances. The matrix, when populated with a full range of contributing factors for the aspects of DOHaD being addressed, helps bridge the breadth of causal variables revealed by exploratory science and community prevention, positions these factors in time relative to when people develop vulnerabilities, and specifies the impact of place on the range of solutions available.

This tool also helps us more expansively portray the people involved in DOHaD phenomena, and thus is a first step toward appropriately sharing responsibility. For example, whereas much research frames the mother as the environment as the fetus, the matrix also allows us to demonstrate that society is the environment of the mother. We also can show that she is one of several “hosts” in these processes, simultaneously exposed to and affected by multiple intersecting environments and acting as a conduit through which environments exert influence. Similarly, through the specification of place and intervention options, we can speak of communities in the contexts of the larger social forces in which they operate.

Specifically, the matrix serves as a conceptual, strategic, and communication planning tool for scientists and others wishing to broaden conversations about the etiology of and potential remedies for DOHaD. First, as a **conceptual tool** the matrix incorporates multiple sources of cause and effect (from the individual to societal), a range of contributing factors (from the behavioral to political and economic), and sequential phases in the causal chain (initial and subsequent vulnerabilities). As such, the framework can provide communicators and audiences a robust portrayal of the range of factors that contribute to DOHaD effects.

Second, the matrix can serve as a **strategic planning tool** that provides communicators and audiences a means to map the connections among individual efforts toward a more comprehensive approach. And, by providing a broad portfolio of approaches and points of intervention, it allows audiences to determine in which areas activities are most robust and in which they may be lacking. Thus, it positions communicators and their audiences to evaluate any one approach in relation to others, and creates meaningful connection to other possible solutions.

Third, as a **communication planning tool** the matrix provides a model with which to describe and illustrate the essential and oftentimes reciprocal interconnections between the social determinants of health and their biological impacts. Moreover, it helps people frame

the message around the levels of social effect, seating not only mothers (and others) in their larger contexts, but also placing communities relative to the larger societies that shape them. Finally, because the model helps reach across disciplinary silos and traditional boundaries to emphasize the interconnections among contributions, it invites voices into the conversation that had not previously been included.

Conclusion

Every Institute of Medicine report begins with a quote from Goethe: “Knowing is not enough; we must apply. Willing is not enough; we must do.” These early efforts to translate findings from the field of DOHaD for broad audiences will establish widely shared frames about what this science means and how it can be translated into community benefit. Importantly, these initial frames will also suggest who or what bears responsibility in both the genesis of problems observed and in their resolution.

How we communicate about an issue is rooted in how we understand that issue. The First-hit/Second-hit matrix provides a useful starting point for expanding our thinking about DOHaD and suggesting new ways to talk about it. In particular, we believe that the First-hit/Second-hit framework can help move beyond the traditional variable-by-variable portrayals of the problem to instead bridge the divide between science and community application. By introducing new and broader means of characterizing and speaking to the problem, we can change the questions people ask, and thus also the range of solutions to be considered. Moreover, providing the broader picture will help people speak concretely about potential starting points of intervention, and can help do so through a systems perspective.

This is the beginning of a conversation about how this framework can be adapted to meet DOHaD’s specific communication needs. It is important to note that the matrix is a living instrument, requiring ongoing reevaluation and clarification as circumstances surrounding a health problem and its potential strategies evolve. Maintaining the matrix’s currency will necessitate the continued investment and involvement of a breadth of professional, scientific, and lay constituencies, all who have a recognized stake in this issue [88].

Subsequent steps to developing this approach include empirically determining:

- Which variables should populate each cell in the table for any given problem illuminated by DOHaD, in order to reinforce the continuum of causes, problems, and consequences, and to support the most inclusive public dialogue?
- What specific language, phrasing, verbal “imagery,” and mental models -- all components of issue framing -- are most supportive of audience engagement with these issues and in what ways? Similarly, which may in fact be obstructive?
- How do we balance the short-term need to provide explanations of key findings with the long-term need for conversations about population-level change?

Communicating the field of DOHaD and engaging others in the conversation about how to move forward based on this knowledge will undoubtedly require sensitivity and thoughtfulness. Framing research can provide us with new insights into how policy makers,

public health professionals, and community organizations hear, interpret, and respond to DOHaD findings. Our society has long struggled with a gap between what science tells us and how we structure our programs and policies in response. The potentially paradigm-shifting potential of DOHaD and epigenetic mechanisms provides us with an opportunity to develop a communication approach that is as sophisticated as the science that is changing our understanding of the world.

Acknowledgments

The project described was supported in part by the National Institute of Diabetes and Digestive and Kidney Diseases, award number K01DK102857-01 (JBH).

We would like to thank Ms. Jocelyn Wegman for her thoughtful review of earlier drafts.

References

Papers of particular interest, published recently, have been highlighted as:

• Of importance

•• Of outstanding importance

- Godfrey K, Barker D. Fetal programming and adult health. *Public Health Nutr.* 2001; 4:611–24. [PubMed: 11683554]
- Barker D. The Developmental Origins of Adult Disease. *J Am Coll Nutr.* 2004; 23:588S–95S. [PubMed: 15640511]
- Barker DJ. The origins of the developmental origins theory. *J Int Med.* 2007; 261:412–7.
- Barker D. Developmental Origins of Chronic Disease. *Public Health Nutr.* 2012; 126:185–9.
- Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med.* 2008; 359:61–73. [PubMed: 18596274]
- Wadhwa PD, Buss C, Entringer S, Swanson JM. Developmental origins of health and disease: brief history of the approach and current focus on epigenetic mechanisms. *Semin Reprod Med.* 2009; 27:358–68. [PubMed: 19711246]
- Barker DJ, Thornburg KL. The obstetric origins of health for a lifetime. *Clin Obstet Gynecol.* 2013; 56(3):511–9. [PubMed: 23787713]
- Hanson M. Understanding the origins of diabetes. *JAMA.* 2014; 311:575–6. [PubMed: 24519295]
- Roseboom T, van der Meulen J, Ravelli A, Osmond C, Barker D, Bleker O. Effects of prenatal exposure to the Dutch famine on adult disease in later life: an overview. *Mol Cell Endocrinol.* 2001; 185:93–8. [PubMed: 11738798]
- Barouki R, Gluckman PD, Grandjean P, Hanson M, Heindel JJ. Developmental origins of non-communicable disease: implications for research and public health. *Environ Health.* 2012; 11:42. [PubMed: 22715989]
- Monk C, Georgieff MK, Osterholm EA. Research review: maternal prenatal distress and poor nutrition - mutually influencing risk factors affecting infant neurocognitive development. *J Child Psychol Psychiatry.* 2013; 54:115–30. [PubMed: 23039359]
- Figlio D, Guryan J. The Effects of Poor Neonatal Health on Children's Cognitive Development. *Am Econ Rev.* 2014; 104:3921–55.
- Van den Bergh BR, Mulder EJ, Mennes M, Glover V. Antenatal maternal anxiety and stress and the neurobehavioural development of the fetus and child: links and possible mechanisms. A review. *Neurosci Biobehav Rev.* 2005; 29:237–58. [PubMed: 15811496]
- Bagby SP. Maternal nutrition, low nephron number, and hypertension in later life: pathways of nutritional programming. *J Nutr.* 2007; 137:1066–72. [PubMed: 17374679]

15. Thayer ZM, Kuzawa CW. Biological memories of past environments: epigenetic pathways to health disparities. *Epigenetics*. 2011; 6:798–803. [PubMed: 21597338]
16. Waterland RA, Michels KB. Epigenetic epidemiology of the developmental origins hypothesis. *Annu Rev Nutr*. 2007; 27:363–88. [PubMed: 17465856]
17. Shonkoff JP, Boyce WT, McEwen BS. Neuroscience, molecular biology, and the childhood roots of health disparities: building a new framework for health promotion and disease prevention. *JAMA*. 2009; 301:2252–9. [PubMed: 19491187]
18. Hunter RG. Epigenetic effects of stress and corticosteroids in the brain. *Front Cell Neurosci*. 2012; 6:18. [PubMed: 22529779]
19. Zeisel SH. Epigenetic mechanisms for nutrition determinants of later health outcomes. *Am J Clin Nutr*. 2009; 89:1488S–93S. [PubMed: 19261726]
20. Combs-Orme T. Epigenetics and the Social Work Imperative. *Soc Work*. 2012; 58(1):23–30.
21. Berger SL, Kouzarides T, Shiekhattar R, Shilatifard A. An operational definition of epigenetics. *Genes Dev*. 2009; 23:781–783. [PubMed: 19339683]
22. Kuzawa C, Sweet E. Epigenetics and the embodiment of race: Developmental origins of US Racial Disparities in Cardiovascular Health. *Am J Hum Biol*. 2009; 21:2–15. [PubMed: 18925573]
23. Wallack L, Thornburg K. Developmental origins, epigenetics, and equity: moving upstream. *Matern Child Health J*. 2016; 20:935–940. [PubMed: 27029539]
24. Winett LB, Wulf AB, Wallack L. Framing strategies to avoid mother-blame in communicating the origins of chronic disease. *Am J Public Health*. (forthcoming).
25. Scheufele DA. Framing as a theory of media effects. *J Commun*. 1999; 49(1):103–122.
26. Gitlin, T. *The whole world is watching: mass media in the making and unmaking of the new left*. Berkeley, CA: University of California Press; 1980.
27. Gamson, WA., Modigliani, A. The changing culture of affirmative action. In: Burstein, P., editor. *Equal employment opportunity: labor market distribution and public policy*. Piscataway, NJ: Aldine Transaction Publishers; 1994. p. 373–394.
28. Iyengar, S. *Is anyone responsible? How television frames political issues*. Chicago, IL: University of Chicago Press; 1991.
29. Entman R. Framing: toward clarification of a fractured paradigm. *J Commun*. 1993; 43(4):51–58.
30. Lakoff, G. *Moral politics: what conservatives know that liberals don't*. Chicago, IL: University of Chicago Press; 1999.
31. Ryan, C. *Prime time activism: media strategies for grassroots activism*. Boston, MA: South End Press; 1991.
32. Gamson, WA., Lasch, KE. The political culture of social welfare policy. In: Spiro, SE., Yuchtman-Yar, E., editors. *Evaluating the welfare state: social and political perspectives*. New York, NY: Academic Press; 1983. p. 397–416.
33. Gusfield J. Constructing the ownership of social problems: fun and profit in the welfare state. *Soc Probl*. 1989; 36(5):431–441.
34. Richardson S, Daniels C, Gillman M, Golden J, Kukla R, Kuzawa C, Rich-Edwards J. Don't blame the mothers. *Nature*. 2014; 512:131–132. This paper focuses on the potential for mother-blame in communicating DOHaD science, and calls for targeted communication work to help eliminate that risk. [PubMed: 25119222]
35. Warin M, Moore V, Zivkovic T, Davies M. Telescoping the origins of obesity to women's bodies: How gender inequalities are being squeezed out of Barker's hypothesis. *Ann Human Biol*. 2011; 38(4):453–460. [PubMed: 21696324]
36. Warin M, Zivkovic T, Moore V, Davies M. Mothers as smoking guns: fetal overnutrition and the reproduction of obesity. *Fem Psychol*. 2012; 22(3):360–375.
37. Fillmore C. Frame semantics and the nature of language. *Origins and evolution of language and speech*. *Ann NY Acad Sci*. 1976; 280:20–32.
38. Fillmore, C. *Linguistic Society of Korea. Linguistics in the morning calm*. Seoul: Hanshin Publishing Co; 1982. Frame semantics; p. 111–137.
39. Lakoff, G. *Women, fire, and dangerous things*. Chicago, IL: University of Chicago Press; 1987.
40. Beauchamp DE. Public health as social justice. *Inquiry*. 1976; 13:3–14. [PubMed: 130348]

41. Wallack L, Lawrence R. Talking about public health: developing America's "second language". *Am J Public Health*. 2005; 95(4):56–70.
42. Bernstein, J. All together now: common sense for a fair economy. San Francisco, CA: Berrett-Koehler Publishers; 2006.
43. Haidt, J. The righteous mind: why good people are divided by politics and religion. New York, NY: Pantheon Books; 2012.
44. Goffman, E. Stigma: notes on the management of a spoiled identity. Englewood Cliffs, NJ: Prentice Hall; 1963.
45. Kleinman A, Hall-Clifford R. Stigma: a social, cultural and moral process. *J Epidemiol Community Health*. 2009; 63(6):418–9. [PubMed: 19439576]
46. Weiss MG, Ramakrishna J. Stigma interventions and research for international health. *Lancet*. 2006; 367(9509):536–8. [PubMed: 16473134]
47. Link BG, Phelan JC. Conceptualizing stigma. *Annu Rev Sociol*. 2001; 27:363–85.
48. Wutich A, Ruth A, Brewis A, Boone C. Stigmatized neighborhoods, social bonding, and health. *Med Anthropol Q*. 2014; 28(4):556–77. [PubMed: 25196035]
49. Keene DE, Padilla MB. Spatial stigma and health inequality. *Crit Public Health*. 2014; 24(4):392–404.
50. Ettema JS, Peer L. Good news from a bad neighborhood: toward an alternative to the discourse of urban pathology. *Journalism Mass Comm*. 1996; 73:835–856.
51. Syme SL. The Social-Environment and Health. *Daedalus*. 1994; 123(4):79–86.
52. Frohlich KL, Potvin L. The inequality paradox: the population approach and vulnerable populations. *Am J Public Health*. 2008; 98(2):216–21. [PubMed: 18172133]
53. Rose, G. The strategy of preventive medicine. New York, NY: Oxford University Press; 1992.
54. Chavez, V., Duran, B., Baker, QE., Avila, MM., Wallerstein, N. The Dance of Race and Privilege in CBPR. In: Minkler, MWN., editor. *Community-Based Participatory Research for Health: From Process to Outcomes*. 2. San Francisco, CA: Jossey-Bass; 2008.
55. Wallerstein, NDB., Minkler, M., Foley, K. Developing and maintaining partnerships with communities. In: Israel, BA, Eng, E, Schulz, AJ, Parker, EA., Satcher, D., editors. *Methods in community-based participatory research for health*. San Francisco, CA: Jossey-Bass; 2005.
56. Wallack, L., Woodruff, K., Dorfman, L., Diaz, I. News for a change: an advocate's guide to working with the media. Thousand Oaks, CA: Sage Publications; 1999.
57. Dorfman L, Wallack L, Woodruff K. More than a message: framing public health advocacy to change corporate practices. *Health Educ Behav*. 2005; 32(3):320–336. [PubMed: 15851542]
58. Armitage P, Doll R. The age distribution of cancer and a multi-stage theory of carcinogenesis. *Br J Cancer*. 1954; 8:1–12. [PubMed: 13172380]
59. Knudson AG. Mutation and cancer: statistical study of retinoblastoma. *Proc Natl Acad Sci*. 1971; 68:820–823. [PubMed: 5279523]
60. Maynard TM, Sikich L, Lieberman JA, LaMantia A-S. Neural development, cell-cell signaling, and the "two-hit" hypothesis of schizophrenia. *Schizophr Bull*. 2001; 27(3):457–476. [PubMed: 11596847]
61. Feigenson KA, Kusnecov AW, Silverstein SM. Inflammation and the two-hit hypothesis of schizophrenia. *Neurosci Biobehav Rev*. 2014; 38:72–93. [PubMed: 24247023]
62. Nader, R. Unsafe at any speed: the designed-in dangers of the American automobile. New York, NY: Grossman Publishers; 1965.
- 63•. Messer LC, Boone-Heinonen J, Mponwane L, Wallack L, Thornburg KL. Developmental programming: priming disease susceptibility for subsequent generations. *Curr Epidemiol Rep*. 2015; 2(1):37–51. This paper provides an in-depth discussion of how first and second hit effects may accumulate over multiple generations, manifesting as persistent intergenerational health disparities. [PubMed: 26366336]
- 64•. Boone-Heinonen J, Messer LC, Fortmann SP, Wallack L, Thornburg KL. From fatalism to mitigation: a conceptual framework for mitigating fetal programming of chronic disease by maternal obesity. *Prev Med*. 2015; 81:451–9. This paper provides an in-depth discussion of how

the first and second hits framework can be applied to mitigating fetal priming effects within the domain of public health. [PubMed: 26522092]

65. Haddon W. A logical framework for categorizing highway safety phenomena and activity. *J Trauma*. 1972; 12(3):193–207. [PubMed: 5012817]
66. Thornburg KL, Shannon J, Thuillier P, Turker MS. In utero life and epigenetic predisposition for disease. *Adv Genet*. 2010; 71:57–78. [PubMed: 20933126]
67. Gluckman PD, Hanson MA, Cooper C, Thornburg KL. Effect of in utero and early-life conditions on adult health and disease. *N Engl J Med*. 2008; 359(1):61–73. [PubMed: 18596274]
68. Rauh VA, Horton MK, Miller RL, Whyatt RM, Perera F. Neonatology and the environment: impact of early exposure to airborne environmental toxicants on infant and child neurodevelopment. *Neoreviews*. 2010; 11:363–9. [PubMed: 21566672]
69. Entringer S, Buss C, Swanson JM, Cooper DM, Wing DA, Waffarn F, et al. Fetal programming of body composition, obesity, and metabolic function: the role of intrauterine stress and stress biology. *J Nutr Metab*. 2012; 2012:632548. [PubMed: 22655178]
70. Brunton P. Resetting the dynamic range of hypothalamic-pituitary-adrenal axis stress responses through pregnancy. *J Neuroendocrinol*. 2010; 22(11):1198–1213. [PubMed: 20819122]
71. Thornburg KL, O'Tierney PF, Louey S. Review: the placenta is a programming agent for cardiovascular disease. *Placenta*. 2010; 31(Suppl):S54–59. [PubMed: 20149453]
72. Barker DJ, Thornburg KL. Placental programming of chronic diseases, cancer and lifespan: a review. *Placenta*. 2013; 34(10):841–845. [PubMed: 23916422]
73. Auten RL, Potts EN, Mason SN, Fischer B, Huang Y, Foster WM. Maternal exposure to particulate matter increases postnatal ozone-induced airway hyperreactivity in juvenile mice. *Am J Respir Crit Care Med*. 2009; 180(12):1218–26. [PubMed: 19762564]
74. Bolton JL, Auten RL, Bilbo SD. Prenatal air pollution exposure induces sexually dimorphic fetal programming of metabolic and neuroinflammatory outcomes in adult offspring. *Brain Behav Immun*. 2014; 37:30–44. [PubMed: 24184474]
75. Van den Hove DL, Leibold NK, Strackx E, Martinez-Claros M, Lesch KP, Steinbusch HW, et al. Prenatal stress and subsequent exposure to chronic mild stress in rats; interdependent effects on emotional behavior and the serotonergic system. *Eur Neuropsychopharmacol*. 2014; 24(4):595–607. [PubMed: 24139910]
76. Schopper H, Palme R, Ruf T, Huber S. Effects of prenatal stress on hypothalamic-pituitary-adrenal (HPA) axis function over two generations of guinea pigs (*Cavia aperea f. porcellus*). *Gen Comp Endocrinol*. 2012; 176(1):18–27. [PubMed: 22202601]
77. Entringer S, Kumsta R, Nelson EL, Hellhammer DH, Wadhwa PD, Wust S. Influence of prenatal psychosocial stress on cytokine production in adult women. *Dev Psychobiol*. 2008; 50(6):579–87. [PubMed: 18683180]
78. Tamashiro KL, Terrillion CE, Hyun J, Koenig JI, Moran TH. Prenatal stress or high-fat diet increases susceptibility to diet-induced obesity in rat offspring. *Diabetes*. 2009; 58(5):1116–25. [PubMed: 19188431]
79. Norton JM, Wing S, Lipscomb HJ, Kaufman JS, Marshall SW, Cravey AJ. Race, wealth, and solid waste facilities in North Carolina. *Environ Health Perspect*. 2007; 115(9):1344–50. [PubMed: 17805426]
80. Jones MR, Diez-Roux AV, Hajat A, Kershaw KN, O'Neill MS, Guallar E, et al. Race/ethnicity, residential segregation, and exposure to ambient air pollution: the Multi-Ethnic Study of Atherosclerosis (MESA). *Am J Public Health*. 2014; 104(11):2130–7. [PubMed: 25211756]
81. Laraia B, Messer L, Evenson K, Kaufman JS. Neighborhood factors associated with physical activity and adequacy of weight gain during pregnancy. *J Urban Health*. 2007; 84(6):793–806. [PubMed: 17710552]
82. Miranda M, Messer L, Kroger G. The impact of neighborhood quality on pregnancy outcomes. *Environ Health Perspect*. 2011; 12(3):471–7.
83. Larson N, Story M, Nelson M. Neighborhood environments: disparities in access to healthy foods in the US. *Am J Prev Med*. 2009; 36(1):74–81. [PubMed: 18977112]

84. Moore LV, Diez Roux AV, Evenson KR, McGinn AP, Brines SJ. Availability of recreational resources in minority and low socioeconomic status areas. *Am J Prev Med.* 2008; 34(1):16–22. [PubMed: 18083446]
85. Dubowitz T, Heron M, Bird CE, Lurie N, Finch BK, Basurto-Davila R, et al. Neighborhood socioeconomic status and fruit and vegetable intake among whites, blacks, and Mexican Americans in the United States. *Am J Clin Nutr.* 2008; 87(6):1883–91. [PubMed: 18541581]
86. Lovasi GS, Hutson MA, Guerra M, Neckerman KM. Built environments and obesity in disadvantaged populations. *Epidemiol Rev.* 2009; 31:7–20. [PubMed: 19589839]
87. Runyan C. Using the Haddon Matrix: introducing the third dimension. *Injury Prev.* 1998; 4:302–307.
88. Williams, A. The Haddon Matrix: its contribution to injury prevention and control. In: McClure, R., editor. *Third National Conference on Injury Prevention and Control.* Queensland: Brisbane; 1999.
89. Runyan C. Back to the future: revisiting Haddon's conceptualization of injury epidemiology and prevention. *Epidemiol Rev.* 2003; 25:60–64. [PubMed: 12940231]
90. Barnett DJ, Balicer RD, Lucey DR, Everly GS, Omer SB, Steinhoff MC, Grotto I. A systematic analytic approach to pandemic influenza preparedness planning. *PLoS Med.* 2005; 2(12):1235–1241.
91. Barnett DJ, Balicer RD, Blodgett D, Fewes AL, Parker CL, Links JM. An application of the Haddon Matrix to public health readiness and response planning. *Environ Health Perspect.* 2005; 113(5): 561–566. [PubMed: 15866764]
92. Wall LL. Preventing obstetri fistulas in low-resource countries: Insights from a Haddon Matrix. *Obstet Gynecol Surv.* 2012; 67(2):111–121. [PubMed: 22325301]

Table 1
First-hit/Second-hit Planning Matrix for Communicating Developmental Origins of Health and Disease (DOHaD) With Broad Audiences^{*}

	Host	Agent	Environment (Place) – <i>Examples</i>				Society
	Individual	Nutritional, Physical, and Social Stress	Home	School	Workplace	Community	
Pre-gestation (1 st Hit)	Developing girl through woman of reproductive age	Nutrient flow, toxic exposures, and cortisol levels affecting reproductive environment of future mother	<ul style="list-style-type: none">• Safe, toxicant-free, secure housing	<ul style="list-style-type: none">• Safe, toxicant-free school facilities	<ul style="list-style-type: none">• Safe, toxicant-free workplaces	<ul style="list-style-type: none">• Safety regulations and enforcement	<ul style="list-style-type: none">• Programs and subsidies for community development• Equitable economic and social opportunity• Non discriminatory educational, housing, and employment practices• Accessible quality health care for all• Healthy food systems• Immunization policies• Regulated food pricing, accessibility, marketing• Social norms that value equity and inclusion• Social norms that value women and girls
			<ul style="list-style-type: none">• Economic security	<ul style="list-style-type: none">• Healthy food vending and service policies	<ul style="list-style-type: none">• Living wage	<ul style="list-style-type: none">• Health, safety, and social services	
			<ul style="list-style-type: none">• Healthy food preparation facilities and skills	<ul style="list-style-type: none">• Health, nutrition, and physical education	<ul style="list-style-type: none">• Health insurance	<ul style="list-style-type: none">• Grocery store zoning	
			<ul style="list-style-type: none">• Family support for healthy eating and physical activity		<ul style="list-style-type: none">• Paid sick time	<ul style="list-style-type: none">• Community health centers	
			<ul style="list-style-type: none">• Adherence to immunization schedules		<ul style="list-style-type: none">• Healthy, affordable cafeteria & food storage options	<ul style="list-style-type: none">• Neighborhood associations	
Intrauterine /First 1000 days post-conception (1 st Hit)	Pregnant woman and other family members Developing fetus Male or female child to age 2	Nutrient flow, toxic exposures, and cortisol levels affecting development of offspring's organ systems	<ul style="list-style-type: none">• Safe, toxicant-free, secure housing	<ul style="list-style-type: none">• Safe, toxicant-free child care facilities	<ul style="list-style-type: none">• Safe, toxicant-free workplaces	<ul style="list-style-type: none">• Safety regulations and enforcement	<ul style="list-style-type: none">• Quality prenatal and pediatric health care• Grocery store zoning• Breastfeeding support policies and facilities• Skilled nutritional support during pregnancy• Community-based early childhood recreational programs
			<ul style="list-style-type: none">• Economic security	<ul style="list-style-type: none">• Quality, affordable child care services	<ul style="list-style-type: none">• Living wage	<ul style="list-style-type: none">• Quality prenatal and pediatric health care	
			<ul style="list-style-type: none">• Healthy food preparation facilities and skills	<ul style="list-style-type: none">• Affordable, quality early childhood development programming	<ul style="list-style-type: none">• Health insurance	<ul style="list-style-type: none">• Grocery store zoning	
			<ul style="list-style-type: none">• Breastfeeding support in household	<ul style="list-style-type: none">• Skilled early childhood educators	<ul style="list-style-type: none">• Paid family leave	<ul style="list-style-type: none">• Breastfeeding support policies and facilities	
			<ul style="list-style-type: none">• Social support	<ul style="list-style-type: none">• Healthy food service programs in child care	<ul style="list-style-type: none">• Flexible work time	<ul style="list-style-type: none">• Skilled nutritional support during pregnancy	
Later childhood through adulthood (2 nd Hit)	Developing male or female who has already experienced 1 st hit exposures	Nutrient flow, toxic exposures, and cortisol levels compounding and activating initial	<ul style="list-style-type: none">• Safe, toxicant-free, secure housing	<ul style="list-style-type: none">• Safe, toxicant-free school facilities	<ul style="list-style-type: none">• Safe, toxin-free workplaces	<ul style="list-style-type: none">• Safety regulations and enforcement	<ul style="list-style-type: none">• Health, safety, social services
			<ul style="list-style-type: none">• Economic security	<ul style="list-style-type: none">• Quality PreK-20 education	<ul style="list-style-type: none">• Living wage	<ul style="list-style-type: none">• Health, safety, social services	
					<ul style="list-style-type: none">• Health insurance		

	Host	Agent	Environment (Place) – <i>Examples</i>				Society
	Individual	Nutritional, Physical, and Social Stress	Home	School	Workplace	Community	
			<ul style="list-style-type: none">• Healthy food preparation facilities and skills• Family support for healthy eating and physical activity	<ul style="list-style-type: none">• Health, nutrition, and physical education programs• Safe, accessible recreational facilities• Healthy food vending and service policies	<ul style="list-style-type: none">• Paid sick time• Healthy, affordable cafeteria & food storage options• Worksite health promotion and injury prevention programs	<ul style="list-style-type: none">• Grocery store zoning• Community health centers• Family recreational facilities	

* One of the strengths of the matrix is that it highlights interacting causal factors in the different phases of the health problem. For this reason, single factors may appear in more than one cell. Social factors transcend all DOHaD phases, and are combined into a single column. Factors provided are examples and are not exhaustive.

Matrix Adapted From:
Haddon W. A logical framework for categorizing highway safety phenomena and activity. J Trauma. 1972;12(3):193–207