Sleep alterations in mammals: Did aquatic conditions inhibit rapid eye movement sleep?

Vibha Madan, Sushil K. Jha

School of Life Sciences, Jawaharlal Nehru University, New Delhi-110067, India

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2012

Abstract: Sleep has been studied widely in mammals and to some extent in other vertebrates. Higher vertebrates such as birds and mammals have evolved an inimitable rapid eye movement (REM) sleep state. During REM sleep, postural muscles become atonic and the temperature regulating machinery remains suspended. Although REM sleep is present in almost all the terrestrial mammals, the aquatic mammals have either radically reduced or completely eliminated REM sleep. Further, we found a significant negative correlation between REM sleep and the adaptation of the organism to live on land or in water. The amount of REM sleep is highest in terrestrial mammals, significantly reduced in semi-aquatic mammals and completely absent or negligible in aquatic mammals. The aquatic mammals are obligate swimmers and have to surface at regular intervals for air. Also, these animals live in thermally challenging environments, where the conductive heat loss is approximately ~90 times greater than air. Therefore, they have to be moving most of the time. As an adaptation, they have evolved unihemispheric sleep, during which they can rove as well as rest. A condition that immobilizes muscle activity and suspends the thermoregulatory machinery, as happens during REM sleep, is not suitable for these animals. It is possible that, in accord with Darwin's theory, aquatic mammals might have abolished REM sleep with time. In this review, we discuss the possibility of the intrinsic role of aquatic conditions in the elimination of REM sleep in the aquatic mammals.

Keywords: aquatic mammals; aquatic adaptation; mammalian sleep; phylogenetic evolution

1 Introduction

Two distinct sleep states, non-rapid eye movement (NREM) and rapid eye movement (REM) sleep, are present only in mammals^[1] and birds^[2]. In other vertebrate groups such as fishes, amphibians and reptiles, sleep is unitary in nature^[3–6]. Similar to birds and mammals, fishes^[7], amphibians^[4] and reptiles^[3] periodically undergo sleep-wake states but display spike-like activity in the electroencephalogram (EEG) during sleep and low-amplitude waves

when awake. In birds and mammals, the EEG exhibits a synchronized pattern of high-amplitude slow waves during NREM sleep and a desynchronized pattern of high-frequency waves of low amplitude during wakefulness and REM sleep. These parameters, along with behavioral observation, help to characterize vigilant states in animals^[1,8]. Further, sleep-wakefulness in almost all organisms is regulated by two independent mechanisms: circadian and homeostatic^[9]. Although sleep in invertebrates and lower vertebrates shares several behavioral and functional properties with mammalian and avian sleep, it also has some fundamental differences. For example, sleep in invertebrates and lower vertebrates resembles avian and mammalian sleep in the daily timing of inactivity, periods of increased arousal

Corresponding author: Sushil K. Jha
Tel: +91-11-26704190; Fax: +91-11-26742558
E-mail: sushil_1000@yahoo.com

Article ID: 1673-7067(2012)06-0746-13

Received date: 2011-10-01; Accepted date: 2012-10-16

threshold, and rebound after sleep loss^[10]. However, the different stages of NREM and REM sleep, along with state-specific electrophysiological correlates, have evolved only in mammals and birds. It is believed that the complex nature of sleep in birds and mammals might have evolved gradually from the sleep state of lower organisms^[2].

Since the discovery of REM sleep, a number of studies have been performed to understand how and why it evolved only in birds and mammals. Current knowledge suggests that ancient or basal birds like the ostrich^[11], and mammals such as platypus^[12], opossum^[13] and ferret^[14], manifest a large amount of REM sleep. On the other hand, reptiles (precursors of birds and mammals) exhibit only a single sleep-like state^[15]. REM sleep or a REM sleep-like state has not been persuasively demonstrated in reptiles, suggesting that it has evolved in parallel and independently in mammals and birds. Further, semi-aquatic and aquatic mammals such as fur seals, dolphins and white whales have shed a significant amount of REM sleep and/or have eliminated it completely (Fig. 1). It is not known, however, why REM sleep is present in terrestrial mammals and birds and substantially reduced or eliminated in aquatic mammals. In this review, we primarily discuss the nature of sleep in aquatic and terrestrial mammals and the possible impact of aquatic conditions on the evolution of REM sleep.

2 Discovery of REM sleep

Aserinsky and Kleitman, in 1953, observed for the first time a unique sleep stage in humans (both children and adults) when the eyes rolled in a regular fashion. They noted that the eyes of sleeping subjects were moving at a regular interval during which the EEG was similar to the waking condition^[16]. Since the eyes were moving rapidly while the subject was still asleep, they named this new sleep stage “rapid eye movement (REM) sleep”. They also found that most subjects were dreaming and the respiratory rate was augmented during REM sleep^[16,17]. Later, they also found an increased heart rate during REM sleep^[18]. Following Aserinsky’s work, Dement and Kleitman reported in 1957 that during REM sleep, there were almost

no gross body and limb movements (postural muscle atonia), although numerous jerky limb and digital movements occurred^[19]. In subsequent years, the REM sleep state and its associated phenomena were recorded in several experimental animals. For example, Dement in 1958 identified a sleep state of low-voltage EEG with eye movements in cats^[20]. Jouvet’s group in 1959 found a similar sleep stage occurring after slow-wave sleep, which comprised rapid low-voltage cortical activity with no muscle tone but rapid eye movements with jerky paw and tail movements^[21]. They characterized this stage as ‘paradoxical stage of sleep’ because the EEG resembled that during wakeful-

Fig. 1. Correlation between REM sleep (% total recording time, TRT, in A; % total sleep time, TST, in B) and adaptation to terrestrial, semi-aquatic and aquatic environments in mammals. The percentages of REM sleep relative to total recording time and total sleep time in mammals showed a significant inverse correlation with adaptation to an aquatic medium ($n = 94$). All data points used for analysis are listed in Table 1.

Table 1. Percentage of REM sleep out of total recording time (TRT) and total sleep time (TST) in terrestrial, semi-aquatic and aquatic mammals

	Mammals	REM sleep (% TRT)	REM sleep (% TST)	References
	Terrestrial mammals			
1	Platypus	33.33	56.33	[12]
2	Virginia opossum	30.58	44.97	[74]
3	Latrine opossum	27.21	33.67	[75, 76]
4	Giant armadillo	25.42	33.71	[75, 76]
5	Ferret	24.42	40.41	[14]
6	Southern opossum	23.54	29.12	[13]
7	Large hairy armadillo	18.67	21.94	[41]
8	Big brown bat	16.25	19.79	[77]
9	Nine-banded armadillo	14.83	19.81	[78]
10	Golden hamster	14.63	23.43	[79, 80]
11	Thirteen-lined ground squirrel	14.21	24.71	[80]
12	Arctic ground squirrel	14.08	21.02	[81]
13	Cat	13.42	24.34	[82]
14	Collared lemming	13.29	21.00	[83]
15	Little pocket Mouse	13.12	15.70	[84]
16	Belding's ground squirrel	12.54	19.00	[85]
17	Siberian chipmunk	12.25	24.09	[86]
18	Western european hedgehog	12.00	28.57	[87]
19	Brandt's hedgehog	11.75	27.33	[88]
20	Golden-mantled ground squirrel	11.50	19.01	[85]
21	Kangaroo rat	11.33	17.00	[89]
22	Pig	10.96	29.13	[90]
23	Common tree shrew	10.79	16.40	[91]
24	Norway rat	10.75	19.48	[80]
25	Mountain beaver	10.21	17.01	[87]
26	Red fox	10.00	24.51	[92]
27	Lesser mole rat	9.96	22.53	[93]
28	Common tenrec	9.75	15.00	[87]
29	Northern short-tailed shrew	9.58	15.44	[94]
30	Mongolian jird	9.08	16.60	[95]
31	Star-nosed mole	9.04	21.03	[96]
32	Eastern mole	8.79	24.97	[96]
33	Djungarian hamster	8.63	14.47	[97]
34	House shrew	8.33	15.62	[94]
35	African giant pouched rat	8.33	24.09	[98]
36	Little brown bat	8.29	9.98	[99]
37	Human	7.92	23.75	[98]

To be continued

Continued

38	Blind mole rat	7.92	15.27	[100]
39	Northern night monkey	7.58	10.72	[76]
40	Yellow-bellied marmot	7.54	10.39	[86]
41	Mexican volcano mouse	7.54	10.46	[101]
42	Brushtail possum	7.00	12.28	[102]
43	Chimpanzee	6.75	15.00	[76, 103]
44	Common marmoset	6.67	16.85	[104]
45	Dog	6.46	17.96	[105, 106]
46	Chinchilla	6.46	12.38	[80]
47	Stump-tailed macaque	6.33	15.96	[107]
48	Hispid cotton rat	6.16	13.09	[93]
49	Least shrew	5.83	15.38	[94]
50	Olive baboon	5.79	14.23	[93]
51	Gray mouse lemur	5.71	9.57	[108]
52	Common genet	5.42	21.31	[98]
53	Hamadryas baboon	5.25	12.80	[109]
54	House mouse	5.25	9.58	[80]
55	Pale-throated three-toed sloth	4.95	11.30	[110]
56	Greater short-nosed fruit bat	4.77	7.73	[111]
57	Rabbit	4.54	12.91	[112]
58	Barbary macaque	4.45	9.11	[113]
59	Pig-tailed macaque	4.42	11.04	[114]
60	Guinea pig	4.42	12.28	[115]
61	Bonnet macaque	4.37	11.45	[116]
62	Common squirrel monkey	4.29	11.28	[117]
63	Rhesus macaque	4.08	10.95	[118]
64	Yellow-spotted hyrax	3.79	17.16	[119]
65	Senegal galago	3.63	11.15	[120]
66	Patas monkey	3.58	7.93	[121]
67	Lesser dawn fruit bat	3.57	5.80	[111]
68	Black lemur	3.50	8.70	[122]
69	Horse	3.29	27.43	[39]
70	Yellow baboon	3.25	7.88	[123]
71	Cow	3.13	18.89	[39]
72	Goat	2.95	13.15	[40]
73	Long-nosed potoroo	2.58	5.81	[124]
74	Vervet monkey	2.50	5.80	[121]
75	Sheep	2.37	14.81	[39]
76	Rock hyrax	2.29	11.23	[119]
77	Guinea baboon	2.25	6.01	[109]

To be continued

Continued

78	Degu	2.16	7.51	[125]
79	Tree hyrax	2.08	10.22	[119]
80	Mongoose lemur	1.50	3.03	[122]
Semi-aquatic mammals				
81	Southern sea lion	9.71	39.29	[46]
82	Gray seal	6.25	24.19	[52]
83	Walrus	5.00	8.33	[48]
84	Elephant seal	5.00	11.11	[51]
85	Northern fur seal	4.70	12.46	[54]
86	Cape fur seal	4.58	14.32	[45]
87	Harp seal	3.2	12.83	[50]
88	Sea otter	2.50	6.66	[44]
89	Caspian seal	1.71	11.78	[49]
Aquatic mammals				
90	Amazon manatee	1	3.57	[64]
91	Bottlenose dolphin	0	0	[60]
92	Black sea porpoise	0	0	[61]
93	White whale	0	0	[63]
94	Amazon dolphin	0	0	[62]

Note: Total sleep time in aquatic mammals is quantified by combining all slow-wave stages, such as moderate synchronization with sleep spindles and delta waves appearing individually in either hemisphere during unihemispheric sleep.

ness, whereas behaviorally the animal remained asleep^[21]. In the same year, this sleep stage was also recorded in the rabbit^[22], and in subsequent years in the dog^[23], monkey^[24], rat^[25], mouse^[26] and several other mammals (details in Table 1). The classic signs of REM sleep, such as fast low-voltage EEG, atonia, cessation of thermoregulation, and ponto-geniculo-occipital (PGO) spikes were discovered first in the cat, but tonic features associated with REM sleep were later recorded in most terrestrial mammals (for details see review^[27]).

After the discovery of REM sleep in mammals, attempts were made to find it in non-mammalian species such as birds and reptiles. In 1964, two groups (Klein *et al.*, and Ookawa and Gotoh) reported a periodic recurrence of fast low-voltage EEG waves during behavioral sleep in birds^[28,29], which was later reconfirmed as REM/paradoxical sleep^[30–33]. Interestingly, hippocampal theta activity (4–7

Hz), typical of mammalian REM sleep and equivalent to spontaneous mammalian PGO waves had not been reliably observed in birds^[32]. Although some of the REM sleep-associated tonic features were not reliably recorded in birds, it was confirmed by several reports that REM sleep is present in birds^[34]. In order to trace the phylogenetic correlates of REM sleep in lower vertebrates, attempts were made to characterize this sleep state in reptiles such as crocodiles, alligators, lizards, snakes and tortoises but REM sleep or a REM sleep-like state has never been observed or recorded in reptiles^[7]. Although Huntley and associates and a few others have reported the presence of REM sleep in some reptiles^[35], clear confirmatory evidence is still lacking. Since the discovery of REM sleep, several studies have clearly demonstrated that it is found distinctly in birds and mammals and not in lower vertebrates such as reptiles, amphibians and fishes.

3 REM sleep in the terrestrial, semi-aquatic and aquatic mammals: Comparative studies

3.1 REM sleep in terrestrial mammals Sleep studies have mostly been performed in mammals and REM sleep has been electrophysiologically recorded in ~80 terrestrial species (Table 1). Phylogenetically, mammals fall into three sub-classes: (1) prototheria [egg-laying mammals: the echidna (*Tachyglossus aculeatus*) and the duckbilled platypus (*Ornithorhynchus anaticus*)]; (2) metatheria (marsupials such as kangaroo and opossum); and (3) eutheria (placental mammals). In the mammalian lineage, it is believed that the prototheria might have evolved from reptiles, hence are ancient in origin and a connecting link between reptiles and mammals^[36]. Some features of the echidna such as microchromosomes, filiform sperm, and egg-laying, closely resemble reptiles^[36] and interestingly, REM sleep has not been found reliably in echidna and reptiles^[7,35,37]. It was proposed, therefore, that REM sleep might have originated in recent metatheria and/or eutheria^[37]. Siegel's group, however, later found that REM sleep is present in the duckbilled platypus^[12], thus refuting the hypothesis that REM sleep evolved relatively recently. They observed that the platypus not only exhibits REM sleep, but also displays the highest amount (5.8–8 h/day), more than any other animal studied so far^[12]. Further, based on the firing properties of brainstem neurons in the echidna, the same group reported that echidna sleep may have features of combined REM and NREM sleep. It remains ambiguous whether the echidna has combined aspects of REM and NREM sleep or no REM sleep because only one study has reported some of the typical characteristics of REM sleep^[38].

Ancient mammals spend the most time in REM sleep (platypus: 33.33%; Virginia opossum: 30.58%; lutrine opossum: 27.20%; ferret: 24.41%) while relatively modern mammals spend varied amounts of time, ranging from 13.42% in the cat to 3% in the cow, goat and sheep (Table 1). A decreasing trend was further noted in the mongoose lemur, which spends only 1.5% of its time per day in REM sleep (the lowest amount of REM sleep in any terrestrial

mammal studied so far) (Table 1). These data suggest that REM sleep might have evolved in ancient mammals or their ancestors, but later, during the course of evolution, it remained high in some while it dwindled in others. Further, some mammalian species sleep notably less than others. For example, cow and goat sleep only 4–5 h/day^[39,40], while the large hairy armadillo usually sleeps 20 h/day^[41]. It can also be argued that animals that sleep less would exhibit less REM sleep. Although this could be true, an important interesting fact is that the majority of both long and short sleepers maintain REM sleep at a relatively constant $16.87 \pm 0.009\%$ of the total sleep period. Species that show exceptionally different NREM/REM sleep proportions are the platypus, opossum and ferret, which have a very high proportion of REM sleep (25–56%) and some species of bat, monkey and lemur which exhibit a very low proportion (3–7%) of the total sleep period (details in Table 1). We do not yet have a clear understanding of why some terrestrial mammals have reduced their total sleep time, although some reasons could be energy conservation, survival strategy, ecological need or other evolutionary forces^[42,43].

3.2 REM sleep in semi-aquatic mammals Pinnipeds, semi-aquatic mammals, have adapted their sleep to the aquatic environment. They have a distinctive sleep pattern as they can sleep on land as well as in water. On land, their sleep is very similar to terrestrial mammals displaying both NREM and REM sleep with bihemispheric symmetry in slow-wave generation during NREM sleep. While in water, they exhibit inter-hemispheric asymmetry in the generation of slow-waves during NREM sleep and interestingly, a substantial decrease in REM sleep^[44–48]. Some pinnipeds, such as Caspian seals (*Phoca caspica*)^[49], harp seals (*Pagophilus groenlandica*)^[50] and elephant seals (*Mirounga angustirostris*)^[51], do not exhibit inter-hemispheric asymmetry in slow-wave generation during NREM sleep, suggesting that unihemispheric sleep is typical of fur seals and sea lions but not of all semi-aquatic pinnipeds.

Ridgway *et al.*, in 1975, for the first time studied sleep telemetrically in four adult gray seals (*Halichoerus grypus*)^[52] and found that they can sleep under the water, on the sur-

face, and out of the water. They found that sleep in gray seals is distinct from that in terrestrial mammals in many ways: (1) REM sleep comprises 6.35% of total recording time, and is accompanied by rapid but regular heart and respiratory rates (which are usually irregular during REM sleep in terrestrial mammals); and (2) REM sleep appears before NREM sleep, whereas in other terrestrial mammals it normally follows NREM sleep^[52]. Later, Mukhametov *et al.*, while recording sleep polygraphically in three subadult northern fur seals (*Callorhinus ursinus*), observed a sleep pattern similar to that reported by Ridgway *et al.* in the gray seal^[52]. Mukhametov *et al.* recorded 5.8% REM sleep in the northern fur seal^[53]. Interestingly, they also showed, for the first time, an inter-hemispheric asymmetry in the generation of slow waves during NREM sleep in the northern fur seal^[53]. Later, they further confirmed this unique event as ‘the generation of inter-hemispheric asynchrony of slow wave activity’ in the northern fur seal and suggested that this phenomenon is not specific to aquatic cetaceans^[54]. Nevertheless, in two other pinnipeds (the Caspian seal and the harp seal, which are morphologically very close to the northern fur seal), inter-hemispheric asymmetry in slow waves has not been found^[49,50]. Although the harp seal does not exhibit unilateral sleep, its sleep pattern exhibits a remarkable phenomenon. When harp seals sleep on land, they exhibit ~3.2% REM sleep, but in water they barely go into REM (0.5%)^[50]. Castellini *et al.* compared sleep in northern elephant seal pups under dry and wet conditions and demonstrated that the length of REM sleep episodes significantly decreases under wet conditions^[51]. Similar characteristics have also been recorded in northern fur seals^[55]. When the northern fur seal sleeps on land, it exhibits ~5% REM sleep but when it sleeps in water, the proportion of REM sleep decreases considerably^[55]. In order to find similar REM sleep patterns in other mammals that have adapted to constant living in the aquatic environment, Laymin *et al.* investigated the behavioral sleep pattern in sea otters (*Enhydra lutris*) and found that they exhibit REM sleep-like behavior in water as well as on land^[44], though it is not clear if the proportion decreases significantly when they sleep in water. Interest-

ingly, it was noted that the longest episodes of REM sleep occurred on land^[44]. Further, they studied the sleep patterns in the walrus, the only surviving representative of the family Odobenidae. When on land, the walrus exhibits 6.9% REM sleep, but in water this significantly decreases to 1.1%^[47]. This evidence thus suggests that REM sleep is not expressed at its full length if these animals sleep in water.

The three families of pinnipeds, Odobenidae, Otariidae and Phocidae, comprise 35 species. Among these, the sleep-wakefulness pattern has been studied in only 8, and REM sleep has been reported in all of these. Most of these animals [walrus^[47] (Odobenidae), elephant seal^[51] and harp seal^[50] (Phocidae), and northern fur seal^[55] (Otariidae)] exhibit either less REM sleep or shorter REM episodes in water than on land. In another four pinnipeds: gray seal^[52] and Caspian seal^[49] (Phocidae); cape fur seal^[45] and sea lion^[46] (Otariidae) the amount of REM sleep could not be characterized in both conditions, so it is undetermined whether these animals also exhibit differential REM sleep expression on land and in water. REM sleep was also found in another semi-aquatic mammal, the sea otter, which belongs to the Mustelidae and is adapted for constant living in the aquatic environment. Nevertheless, out of the nine semi-aquatic mammalian species studied so far, the majority (five species) express more REM sleep when on land, but drastically curtail it while sleeping in water. Thus, it seems that aquatic conditions may not be favorable for REM sleep.

3.3 REM sleep in aquatic mammals Aquatic mammals belonging to two families, Cetacea and Sirenia, have a notable sleep pattern: one half of the brain remains active while the other half sleeps, and each half exhibits ~4 h of slow-wave sleep per day^[56]. John Lilly for the first time noted that dolphins sleep with one eye open and the other closed^[57]. He assumed that the dolphin scans its environment all the time by keeping one half of the brain always active^[57]. His finding may have led to the discovery of uni-hemispheric sleep in dolphins^[56].

Mukhametov and his team (in 1975) recorded EEG in the bottle-nose dolphin for the first time and reported occasional asynchronous development of EEG slow waves in both hemispheres^[58]. In addition, they noted that the

dolphin does not show any polygraphic signs of REM sleep^[58], though an earlier study by Shurley *et al.* (in 1969) reported episodes of REM sleep in the pilot whale^[59]. In all subsequent physiological studies, however, it was confirmed that REM sleep is absent from the Cetacea bottle nose dolphin^[60], black sea porpoise^[61], amazon river dolphin^[62] and beluga whale^[63], while in the amazon manatee (Sirenia), only a few episodes of REM sleep were recorded^[64]. Although sleep has been studied extensively in only a few cetaceans and REM sleep is absent in most of them, further studies are needed to confirm whether REM sleep or its associated signs are indubitably absent in Cetacea. Lyamin *et al.* pointed out in their review that the traditional criteria used to identify REM sleep may not be appropriate for cetaceans because if dolphins have REM sleep of short duration, it could remain unnoticed^[56]. Another reason for unidentified REM sleep in cetaceans could be the possibility that they have a modified form of REM sleep^[56]. The latter, however, seems unlikely because the semi-aquatic pinnipeds exhibit the conventional form of REM sleep and its associated signs, living in similar conditions in water and on land. The difference in REM sleep in water and on land is that pinnipeds have significantly less REM sleep in water. It is likely that during the course of divergent evolution, REM sleep may not necessarily be required, and thus was eliminated in all aquatic mammals.

4 Do aquatic conditions disfavor REM sleep continuity?

Sleep in some semi-aquatic mammals is unique because it changes from bihemispheric on land to unihemispheric in water. It is noteworthy that when on land, sleep in fur seals generally resembles that of most terrestrial mammals, i.e. the sleep cycle alternates between NREM and REM sleep and the bilateral EEG exhibits synchronization during NREM sleep and desynchronization during REM sleep. But when the fur seal is in water, the occurrence of REM sleep declines to the extent that there may not be even a single episode^[46]. It is not known how the brain switches from terrestrial bihemispheric to aquatic unihemispheric sleep in semi-aquatic mammals. In fact, it has

been suggested that sleep in fur seals defies the principle of homeostatic regulation, since no rebound of lost REM is seen when it returns to land after staying several weeks in water^[42]. Unihemispheric and bihemispheric sleep have their own specific properties, probably regulated by different mechanisms. A neuronal mechanism activated in animals when they are in water could induce the distinctive unihemispheric aquatic sleep and simultaneously eliminate REM sleep. Hence, one would not expect a REM rebound once the organism regained bihemispheric terrestrial sleep.

The amount of REM sleep in mammals may depend on their degree of adaptation to the aquatic and non-aquatic environments. REM sleep in completely aquatic mammals such as the dolphin, porpoise, white whale and manatee is either completely absent or present in a negligible amount (dolphin and white whale: 0%; manatee: 1%)^[64,65]. Semi-aquatic mammals such as seals exhibit comparatively more REM sleep than aquatic mammals but less than terrestrial mammals. The amount of REM sleep (% of total recording time and % of total sleep time) in 94 mammalian species is listed in Table 1. When we correlated the percentage of REM sleep out of total recording time (Fig. 1A) and also out of total sleep time (Fig. 1B) with adaptation to aquatic, semi-aquatic and terrestrial conditions for these 94 mammals, the correlations were significantly negative. Most terrestrial mammals exhibit a high amount and proportion of REM sleep compared to semi-aquatic mammals, while aquatic mammals either do not have REM sleep or even if they have it, it is very little. This suggests that when animals remain in water, aquatic conditions somehow restrain the genesis of REM sleep.

Studies on the sleep patterns in reptiles that have either adapted to aquatic life or live near water [such as the turtle, black iguanid lizard (*Ctenosaura pectinata*) and green iguanid lizard (*Iguana iguana*)] are limited. It is likely that the unitary form of sleep will be found in such animals. Sleep remains to be examined thoroughly in reptiles that stay away from water. This might provide clues regarding the evolution of the dual nature of sleep. Interestingly, REM sleep can be artificially abolished in terrestrial mammals using specific deprivation methods, such

as disk-over water, multiple or single platform, and wire-mesh grid platform in which water is used as the surrounding medium^[66-69]. In the wire-mesh grid platform method, the animal is kept on a platform with a guard of wire mesh grid and the water level is kept 1 cm below the platform in a tank, so the animal cannot fall into the water and/or make contact with it. Merely keeping an animal in such a condition suppresses REM sleep. However, keeping an animal over water could be stressful and/or fear-inducing, and it is known that stress/fear suppresses REM sleep^[70]. It is not clear what contributes to the elimination of REM sleep in these protocols, stress or the presence of water; but from the examples of sleep patterns in the semi-aquatic pinnipeds, it seems that less REM sleep is expressed in aquatic conditions.

5 Possible advantages of REM sleep elimination in aquatic mammals

Significantly less REM sleep in semi-aquatic and some aquatic mammals and no REM sleep in cetaceans raise unanswered questions, such as whether the evolution of REM sleep in vertebrates was confined to life on land and/or whether adaptation to life in an aquatic environment inhibited the occurrence of REM sleep. It is possible that REM sleep serves a meaningful purpose only in terrestrial animals and hence its evolution is restricted to them. In the majority of terrestrial mammals, basic somatic features of REM sleep such as atonia, REMs and myoclonic twitches occur^[71]. The dramatic atonic condition during REM sleep occurs because the motoneurons, which innervate muscle fibers, are actively inhibited^[72]. The aquatic mammals are obligate swimmers and have evolved unihemispheric sleep, during which they can sleep with one half of the brain and simultaneously remain moving to sustain life. Therefore, a condition that could immobilize them or restrict their movement, as usually happens during REM sleep in terrestrial mammals, cannot be advantageous. In addition, it has been observed in terrestrial mammals that the thermoregulatory mechanism and response to thermal load remains suspended or inhibited during REM sleep^[71], so the responses are similar to those of poikilothermic animals.

The aquatic mammals live in thermally challenging environments and it is known that conductive heat loss is ~90 times greater in water than air. Since all mammals have distinctive thermoregulatory machinery, aquatic mammals do not differ in this from other mammals. If aquatic mammals had REM sleep, they would face the herculean task of managing their body temperature during REM sleep. There could, however, be several other reasons for doing away with REM sleep. Considering all, it seems that REM sleep may not be favorable for life in an aquatic environment and might have subsequently been reduced or eliminated in aquatic mammals.

6 Conclusion

We have limited information to understand the evolution of the dual nature of sleep in higher vertebrates. REM sleep has been characterized in birds and mammals only. Within mammals, however, REM sleep is absent in cetaceans and is drastically reduced in semi-aquatic mammals. Although in-depth sleep studies have only been performed in a limited number of aquatic and semi-aquatic species, the data consistently show less or no REM sleep in these animals. Since aquatic and semi-aquatic mammals live in a challenging environment, the presence of REM sleep could do more harm than good. Therefore, REM sleep might have been reduced or eliminated in the evolution of some mammals. This also indicates a great influence of selection/evolutionary forces on sleep characteristics. It has also been proposed that some aquatic mammals may survive for several weeks without sleep, while terrestrial mammals cannot survive long^[73]. This suggests that the sleep patterns in aquatic and terrestrial conditions are basically distinct. Therefore, we need to develop a detailed understanding of the evolution of sleep and its functions.

Acknowledgements: This review was supported by grants from DBT, DST, UGC-CAS, New Delhi, India.

References:

- [1] Zepelin H, Siegel JM, Tobler I. Mammalian Sleep. In: Kryger MH,

- Roth T, Dement WC (Eds.) Principles and Practice of Sleep Medicine. Philadelphia: WB Saunders Company, 2005: 91–100.
- [2] Rattenborg NC. Evolution of slow-wave sleep and palliopalial connectivity in mammals and birds: a hypothesis. *Brain Res Bull* 2006, 69: 20–29.
- [3] Gonzalez J, Gamundi A, Rial R, Nicolau MC, de Vera L, Pereda E. Nonlinear, fractal, and spectral analysis of the EEG of lizard, *Gallotia galloti*. *Am J Physiol* 1999, 277: R86–93.
- [4] Lazarev SG. Electroencephalographic analysis of wakefulness and the primary form of sleep in the frog, *Rana temporaria*. *Zh Evol Biokhim Fiziol* 1978, 14: 379–384.
- [5] Nitz DA, van Swinderen B, Tononi G, Greenspan RJ. Electrophysiological correlates of rest and activity in *Drosophila melanogaster*. *Curr Biol* 2002, 12: 1934–1940.
- [6] Ramon F, Hernandez-Falcon J, Nguyen B, Bullock TH. Slow wave sleep in crayfish. *Proc Natl Acad Sci U S A* 2004, 101: 11857–11861.
- [7] Hartse KM. Sleep in insects and non-mammalian vertebrates. In: Kryger MH, Roth T, Dement WC (Eds.). Principles and Practice of Sleep Medicine. Philadelphia: WB Saunders Company, 1989: 64–73.
- [8] Roth TC 2nd, Lesku JA, Amlaner CJ, Lima SL. A phylogenetic analysis of the correlates of sleep in birds. *J Sleep Res* 2006, 15: 395–402.
- [9] Cirelli C. The genetic and molecular regulation of sleep: from fruit flies to humans. *Nat Rev Neurosci* 2009, 10: 549–560.
- [10] Cirelli C, Tononi G. Is sleep essential? *PLoS Biol* 2008, 6: e216.
- [11] Lesku JA, Meyer LC, Fuller A, Maloney SK, Dell'Omo G, Vyssotski AL, *et al.* Ostriches sleep like platypuses. *PLoS One* 2011, 6: e23203.
- [12] Siegel JM, Manger PR, Nienhuis R, Fahringer HM, Shalita T, Pettigrew JD. Sleep in the platypus. *Neuroscience* 1999, 91: 391–400.
- [13] van Twyver H, Allison T. Sleep in the opossum *Didelphis marsupialis*. *Electroencephalogr Clin Neurophysiol* 1970, 29: 181–189.
- [14] Jha SK, Coleman T, Frank MG. Sleep and sleep regulation in the ferret (*Mustela putorius furo*). *Behav Brain Res* 2006, 172: 106–113.
- [15] Flanigan WF Jr. Sleep and wakefulness in iguanid lizards, *Ctenosaura pectinata* and *Iguana iguana*. *Brain Behav Evol* 1973, 8: 401–436.
- [16] Aserinsky E, Kleitman N. Regularly occurring periods of eye motility, and concomitant phenomena, during sleep. *Science* 1953, 118: 273–274.
- [17] Aserinsky E, Kleitman N. Regularly occurring periods of eye motility, and concomitant phenomena, during sleep. 1953. *J Neuropsychiatry Clin Neurosci* 2003, 15: 454–455.
- [18] Aserinsky E, Kleitman N. Two types of ocular motility occurring in sleep. *J Appl Physiol* 1955, 8: 1–10.
- [19] Dement W, Kleitman N. Cyclic variations in EEG during sleep and their relation to eye movements, body motility, and dreaming. *Electroencephalogr Clin Neurophysiol* 1957, 9: 673–690.
- [20] Dement W. The occurrence of low voltage, fast, electroencephalogram patterns during behavioral sleep in the cat. *Electroencephalogr Clin Neurophysiol* 1958, 10: 291–296.
- [21] Jouvet M, Michel F, Courjon J. Sur un stade d'activité électrique cérébrale rapide au cours du sommeil physiologique. *C R Soc Biol* 1959, 153: 1024–1028. [Article in French]
- [22] Sawyer CH, Kawakami M. Characteristics of behavioral and electroencephalographic after-reactions to copulation and vaginal stimulation in the female rabbit. *Endocrinology* 1959, 65: 622–630.
- [23] Shimazono Y, Horie T, Yanagisawa Y, Hori N, Chikazawa S, Shozuka K. The correlation of the rhythmic waves of the hippocampus with the behaviors of dogs. *Neurologia* 1960, 2: 82–88.
- [24] Weitzmann ED. A note on the EEG and eye movements during behavioral sleep in monkeys. *Electroencephalogr Clin Neurophysiol* 1961, 17: 376–381.
- [25] Michel F, Klein M, Jouvet M, Valatx JL. Etude polygraphique du sommeil chez le Rat. *C R Soc Biol* 1961, 155: 2389–2392. [Article in French]
- [26] Weiss T, Fifkova E. Sleep cycles in mice. *Physiol Bohemoslov* 1964, 13: 242–245.
- [27] Gottesmann C. The golden age of rapid eye movement sleep discoveries. I. Lucretius--1964. *Prog Neurobiol* 2001, 65: 211–287.
- [28] Ookawa T, Gotoh J. Electroencephalographic study of chickens: Periodic recurrence of low voltage and fast waves during behavioral sleep. *Poultry Sci* 1964, 43: 1603–1604.
- [29] Klein M, Michel F, Jouvet M. Etude polygraphique du sommeil chez les oiseaux. *C R Soc Biol (Paris)* 1964, 158: 99–103. [Article in French]
- [30] Rojas-Ramirez JA, Tauber ES. Paradoxical sleep in two species of avian predator (*Falconiformes*). *Science* 1970, 167: 1754–1755.
- [31] Tradardi V. Sleep in the pigeon. *Arch Ital Biol* 1966, 104: 516–521.
- [32] Van Twyver H, Allison T. A polygraphic and behavioral study of sleep in the pigeon (*Columba livia*). *Exp Neurol* 1972, 35: 138–153.
- [33] Walker JM, Berger RJ. Sleep in the domestic pigeon (*Columba livia*). *Behav Biol* 1972, 7: 195–203.
- [34] Amlaner CJ, Ball NJ. Avian sleep. In: Kryger MH, Roth T, Dement WC (Eds.). Principles and Practice of Sleep Medicine. Philadelphia: W.B. Saunders Company, 1994: 81–94.
- [35] Huntley AC, Friedmann JK, Cohen HB. Sleep in an Iguanid lizard, *Dipsosaurus dorsalis*. *Sleep Res* 1977, 6: 104.
- [36] Goldman AS. Evolution of the mammary gland defense system and the ontogeny of the immune system. *J Mammary Gland Biol Neoplasia* 2002, 7: 277–289.
- [37] Allison T, Van Twyver H, Goff WR. Electrophysiological studies of

- the echidna, *Tachyglossus aculeatus*. I. Waking and sleep. *Arch Ital Biol* 1972, 110: 145–184.
- [38] Nicol SC, Andersen NA, Phillips NH, Berger RJ. The echidna manifests typical characteristics of rapid eye movement sleep. *Neurosci Lett* 2000, 283: 49–52.
- [39] Ruckebusch Y. The relevance of drowsiness in the circadian cycle of farm animals. *Anim Behav* 1972, 20: 637–643.
- [40] Bell FR, Itabashi T. The electroencephalogram of sheep and goats with special reference to rumination. *Physiol Behav* 1973, 11: 503–514.
- [41] Affanni JM, Cervino CO, Marcos HJ. Absence of penile erections during paradoxical sleep. Peculiar penile events during wakefulness and slow wave sleep in the armadillo. *J Sleep Res* 2001, 10: 219–228.
- [42] Siegel JM. Clues to the functions of mammalian sleep. *Nature* 2005, 437: 1264–1271.
- [43] Siegel JM. Do all animals sleep? *Trends Neurosci* 2008, 31: 208–213.
- [44] Lyamin O, Oleksenko AI, Sevostyanov VF, Nazarenko EA, Mukhametov LM. Behavioral sleep in captive sea otters. *Aquatic Mammals* 2000, 26: 132–136.
- [45] Lyamin OI, Chetyrbok IS. Unilateral EEG activation during sleep in the Cape fur seal, *Arctocephalus pusillus*. *Neurosci Lett* 1992, 143: 263–266.
- [46] Lyamin OI, Mukhametov LM, Chetyrbok IS, Vassiliev AV. Sleep and wakefulness in the southern sea lion. *Behav Brain Res* 2002, 128: 129–138.
- [47] Lyamin OI, Kosenko PO, Vyssotski AL, Lapierre JL, Siegel JM, Mukhametov LM. Study of sleep in a walrus. *Dokl Biol Sci* 2012, 444: 188–191.
- [48] Pryaslova JP, Lyamin OI, Siegel JM, Mukhametov LM. Behavioral sleep in the walrus. *Behav Brain Res* 2009, 201: 80–87.
- [49] Mukhametov LM, Supin A, Poliakova IG. Sleep of Caspian seals. *Zh Vyssh Nerv Deiat Im I P Pavlova* 1984, 34: 259–264.
- [50] Lyamin OI. Sleep in the harp seal (*Pagophilus groenlandica*). Comparison of sleep on land and in water. *J Sleep Res* 1993, 2: 170–174.
- [51] Castellini MA, Milsom WK, Berger RJ, Costa DP, Jones DR, Castellini JM, *et al.* Patterns of respiration and heart rate during wakefulness and sleep in elephant seal pups. *Am J Physiol* 1994, 266: R863–869.
- [52] Ridgway SH, Harrison RJ, Joyce PL. Sleep and cardiac rhythm in the gray seal. *Science* 1975, 187: 553–555.
- [53] Mukhametov LM, Lyamin OI, Poliakova IG. Sleep and wakefulness in *Callorhinus ursinus*. *Zh Vyssh Nerv Deiat Im I P Pavlova* 1984, 34: 465–471.
- [54] Mukhametov LM, Lyamin O, Polyakova IG. Interhemispheric asynchrony of the sleep EEG in northern fur seal. *Experientia* 1985, 41: 1034–1035.
- [55] Lyamin O, Mukhametov LM. Organization of sleep in the northern fur seal. In: Sokolov VE, Aristov AA, Lisitzina TU (Eds.). *The Northern Fur Seal. Systematic, Morphology, Ecology, Behavior*. Moscow: Nauka, 1998: 280–302.
- [56] Lyamin OI, Manger PR, Ridgway SH, Mukhametov LM, Siegel JM. Cetacean sleep: an unusual form of mammalian sleep. *Neurosci Biobehav Rev* 2008, 32: 1451–1484.
- [57] Lilly JC. Animals in the aquatic environments: adaptations of mammals to the ocean. In: Dill DB (Ed.). *Handbook of Physiology-Environment*. Washington DC: American Physiology Society, 1964: 741–747.
- [58] Mukhametov LM, Supin A. An EEG study of different behavioral states of freely moving dolphins. *Zh Vyssh Nerv Deiat Im I P Pavlova* 1975, 25: 396–401.
- [59] Shurley JT, Serafetinides EA, Brooks RE, Elsner R, Kenney DW. Sleep in Cetaceans: The pilot whale, *Globicephala scammoni*. *Psychophysiology* (abstract) 1969, 6: 230.
- [60] Mukhametov LM, Oleksenko AI, Poliakova IG. Quantification of ECoG stages of sleep in the bottlenose dolphin. *Neurophysiology* 1988, 20: 532–538.
- [61] Mukhametov LM, Poliakova IG. Electroencephalographic study of sleep in Sea of Azov porpoises. *Zh Vyssh Nerv Deiat Im I P Pavlova* 1981, 31: 333–339.
- [62] Mukhametov LM. Unihemispheric slow-wave sleep in the Amazonian dolphin, *Inia geoffrensis*. *Neurosci Lett* 1987, 79: 128–132.
- [63] Lyamin OI, Mukhametov LM, Siegel JM, Nazarenko EA, Polyakova IG, Shpak OV. Unihemispheric slow wave sleep and the state of the eyes in a white whale. *Behav Brain Res* 2002, 129: 125–129.
- [64] Mukhametov LM, Lyamin OI, Chetyrbok IS, Vassilyev AA, Diaz RP. Sleep in an Amazonian manatee, *Trichechus inunguis*. *Experientia* 1992, 48: 417–419.
- [65] Zepelin H. Mammalian sleep. In: Dement WC (Ed.). *Principles and Practice of Sleep Medicine*. Philadelphia: Saunders Company, 1989: 30–49.
- [66] Rechtschaffen A, Bergmann BM. Sleep deprivation in the rat: an update of the 1989 paper. *Sleep* 2002, 25: 18–24.
- [67] Mallick BN, Siegel JM, Fahringer H. Changes in pontine unit activity with REM sleep deprivation. *Brain Res* 1990, 515: 94–98.
- [68] Newman SM, Paletz EM, Rattenborg NC, Obermeyer WH, Bencs RM. Sleep deprivation in the pigeon using the Disk-Over-Water method. *Physiol Behav* 2008, 93: 50–58.
- [69] Machado RB, Hipolide DC, Benedito-Silva AA, Tufik S. Sleep deprivation induced by the modified multiple platform technique: quantification of sleep loss and recovery. *Brain Res* 2004, 1004: 45–51.
- [70] Jha SK, Brennan FX, Pawlyk AC, Ross RJ, Morrison AR. REM sleep: a sensitive index of fear conditioning in rats. *Eur J Neurosci*

- 2005, 21: 1077–1080.
- [71] Parmeggiani PL. Physiologic regulation in sleep. In: Kryger MH, Roth T, Dement W (Eds.). *Principles and Practice of Sleep Medicine*. Philadelphia: Elsevier Saunders, 2005: 154–168.
- [72] Chase MH, Morales FR. Controls of motoneurons during sleep. In: Kryger MH, Roth T, Dement W (Eds.). *Principles and Practice of Sleep Medicine*. Philadelphia: Elsevier Saunders, 2005: 154–168.
- [73] Lyamin O, Pryaslova J, Lance V, Siegel J. Animal behaviour: continuous activity in cetaceans after birth. *Nature* 2005, 435: 1177.
- [74] Walker JM, Berger RJ. The ontogenesis of sleep states, thermogenesis, and thermoregulation in the Virginia opossum. *Dev Psychobiol* 1980, 13: 443–454.
- [75] Affanni J. Observations on the sleep of some South American marsupials and edentates. *Perspect Brain Sci* 1972, 1: 21–23.
- [76] Elgar MA, Pagel MD, Harvey PH. Sources of variation in mammalian sleep. *Animal Behavior* 1990, 40: 991–995.
- [77] Zepelin H, Rechtschaffen A. Mammalian sleep, longevity, and energy metabolism. *Brain Behav Evol* 1974, 10: 425–470.
- [78] Prudom AE, Klemm WR. Electrographic correlates of sleep behavior in a primitive mammal, the armadillo *Dasypus novemcinctus*. *Physiol Behav* 1973, 10: 275–282.
- [79] Tobler I, Jaggi K. Sleep and EEG spectra in the Syrian hamster (*Mesocricetus auratus*) under baseline conditions and following sleep deprivation. *J Comp Physiol A* 1987, 161: 449–459.
- [80] Van Twyver H. Sleep patterns of five rodent species. *Physiol Behav* 1969, 4: 901–905.
- [81] Chepkasov IE. Daily rhythm of sleep and wakefulness in the arctic ground squirrel *Citellus parryi* during the summer season. *J Evol Biochem Physiol* 1980, 17: 77–80.
- [82] Ursin R. The two stages of slow wave sleep in the cat and their relation to REM sleep. *Brain Res* 1968, 11: 347–356.
- [83] Karmanova IG, Maksimuk FM, Murav'eva LN, Pastukhov YF, Sazonov VS. Specific features of the cycle 'wakefulness-sleep' in the arctic lemming *Dicrostonyx torquatus*. *J Evol Biochem Physiol* 1979, 15: 190–195.
- [84] Walker JM, Walker LE, Harris DV, Berger RJ. Cessation of thermoregulation during REM sleep in the pocket mouse. *Am J Physiol* 1983, 244: R114–118.
- [85] Walker JM, Glotzbach SF, Berger RJ, Heller HC. Sleep and hibernation in ground squirrels (*Citellus spp*): electrophysiological observations. *Am J Physiol* 1977, 233: R213–221.
- [86] Dijk DJ, Daan S. Sleep EEG spectral analysis in a diurnal rodent: *Eutamias sibiricus*. *J Comp Physiol A* 1989, 165: 205–215.
- [87] Snyder F, Bugbee N, Douthitt TC. Telemetric studies of 24-hour sleep-waking patterns in some primitive mammals. *Psychophysiology* 1972, 9: 122.
- [88] Tauber ES, Michel F, Roffwarg HP. Preliminary note on the sleep and waking cycle in the desert hedgehog (*Paraechinus hypomelas*). *Psychophysiology* 1968, 5: 201.
- [89] Sakaguchi S, Glotzbach SF, Heller HC. Influence of hypothalamic and ambient temperatures on sleep in kangaroo rats. *Am J Physiol* 1979, 237: R80–88.
- [90] Robert S, Dallaire A. Polygraphic analysis of the sleep-wake states and the REM sleep periodicity in domesticated pigs (*Sus scrofa*). *Physiol Behav* 1986, 37: 289–293.
- [91] Berger RJ, Walker JM. A polygraphic study of sleep in the tree shrew (*Tupaia glis*). *Brain Behav Evol* 1972, 5: 54–69.
- [92] Dallaire A, Ruckebusch Y. Rest-activity cycle and sleep patterns in captive foxes (*Vulpes vulpes*). *Experientia* 1974, 30: 59–60.
- [93] Elgar MA, Pagel MD, Harvey PH. Sleep in mammals. *Anim Behav* 1988, 36: 1407–1419.
- [94] Allison T, Gerber SD, Breedlove SM, Dryden GL. A behavioral and polygraphic study of sleep in the shrews *Suncus murinus*, *Blarina brevicauda*, and *Cryptotis parva*. *Behav Biol* 1977, 20: 354–366.
- [95] Susic V, Masirevic G. Sleep patterns in the Mongolian gerbil, *Meriones unguiculatus*. *Physiol Behav* 1986, 37: 257–261.
- [96] Allison T, Van Twyver H. Sleep in the moles, *Scalopus aquaticus* and *Condylura cristata*. *Exp Neurol* 1970, 27: 564–578.
- [97] Deboer T, Franken P, Tobler I. Sleep and cortical temperature in the Djungarian hamster under baseline conditions and after sleep deprivation. *J Comp Physiol A* 1994, 174: 145–155.
- [98] Meddis R. The evolution of sleep. In: Mayes A (Ed.). *Sleep Mechanisms and Functions*. Berkshire: Van Nostrand Reinhold, 1983: 57–95.
- [99] Brebbia DR, Pyne ET. Environmental hypothermia and cerebral electrical activity in the little brown bat, *Myotis lucifugus*. *Psychophysiology* 1972, 9: 122.
- [100] Tobler I, Deboer T. Sleep in the blind mole rat *Spalax ehrenbergi*. *Sleep* 2001, 24: 147–154.
- [101] Ayala-Guerrero F, Vargas-Reyna L, Ramos JJ, Mexicano G. Sleep patterns of the volcano mouse (*Neotomodon alstoni alstoni*). *Physiol Behav* 1998, 64: 577–580.
- [102] LoPresti RW, McGinty DJ. Sleep in the phalanger (*Trichosurus vulpecula*): an Australian marsupial. *Psychophysiology* 1970, 7: 304.
- [103] McNew JJ, Howe RC, Adey WR. The sleep cycle and subcortical-cortical EEG relations to the unrestrained chimpanzee. *Electroencephalogr Clin Neurophysiol* 1971, 30: 489–503.
- [104] Crofts HS, Wilson S, Muggleton NG, Nutt DJ, Scott EA, Pearce PC. Investigation of the sleep electrocorticogram of the common marmoset (*Callithrix jacchus*) using radiotelemetry. *Clin Neurophysiol* 2001, 112: 2265–2273.
- [105] Lucas EA, Powell EW, Murphree OD. Baseline sleep-wake patterns in the pointer dog. *Physiol Behav* 1977, 19: 285–291.
- [106] Copley MP, Jennings DP, Mitler MM. A study of continuous forty-eight hour sleep waking recordings in five dogs. *Sleep Res* 1976, 5: 94.

- [107] Leinonen L, Stenberg D. Sleep in *Macaca arctoides* and the effects of prazosin. *Physiol Behav* 1986, 37: 199–202.
- [108] Barre V, Petter-Rousseaux A. Seasonal variations in sleep-wake cycle in *Microcebus murinus*. *Primates* 1988, 29: 53–64.
- [109] Bert J. Similarities and differences in the sleep of two baboons, *Papio hamadryas* and *Papio papio*. *Electroenceph Clin Neurophysiol* 1973, 39: 657–662.
- [110] De Moura Filho AG, Huggins SE, Lines SG. Sleep and waking in the three-toed sloth, *Bradypus tridactylus*. *Comp Biochem Physiol A Comp Physiol* 1983, 76: 345–355.
- [111] Zhao X, Sun H, Tang Z, Flanders J, Zhang S, Ma Y. Characterization of the sleep architecture in two species of fruit bat. *Behav Brain Res* 2010, 208: 497–501.
- [112] Pivik RT, Bylisma FW, Cooper P. Sleep-wakefulness rhythms in the rabbit. *Behav Neural Biol* 1986, 45: 275–286.
- [113] Gonzalez FF, Zaplana J, Ruiz de Elvira C, Delgado JM. Nocturnal and diurnal sleep in *Macaca sylvana*. *Electroencephalogr Clin Neurophysiol* 1979, 46: 13–28.
- [114] Reite ML, Rhodes JM, Kavan E, Adey WR. Normal sleep patterns in macaque monkey. *Arch Neurol* 1965, 12: 133–144.
- [115] Ibuka N. Ontogenesis of circadian sleep-wakefulness rhythms and developmental changes of sleep in the altricial rat and in the precocial guinea pig. *Behav Brain Res* 1984, 11: 185–196.
- [116] Bert J, Pegram V, Balzamo E. Comparison of sleep between two macaques (*Macaca radiata* and *Macaca mulatta*). *Folia Primatol* 1972, 17: 202–208.
- [117] Edgar DM, Dement WC, Fuller CA. Effect of SCN lesions on sleep in squirrel monkeys: evidence for opponent processes in sleep-wake regulation. *J Neurosci* 1993, 13: 1065–1079.
- [118] Balzamo E, Van Beers P, Lagarde D. Scoring of sleep and wakefulness by behavioral analysis from video recordings in rhesus monkeys: comparison with conventional EEG analysis. *Electroencephalogr Clin Neurophysiol* 1998, 106: 206–212.
- [119] Synder F. Sleep-waking patterns of Hyracoidea. *Sleep Res* 1974, 3: 87.
- [120] Bert J, Collomb H, Martino A. The electroencephalogram of sleep of a prosimian. Its place in the organization of sleep in primates. *Electroencephalogr Clin Neurophysiol* 1967, 23: 342–350.
- [121] Bert J, Pegram V. The sleep electroencephalogram in Cercopithecinae: *Erythrocerbus patas* and *Cercopithecus aethiops sabaeus*. *Folia Primatol* (Basel) 1969, 11: 151–159. [Article in French]
- [122] Balzamo E, Vuillon-Caciuttolo G, Petter JJ, Bert J. States of vigilance in two Lemuridae: EEG patterns and organisation obtained using telemetry. *Waking Sleeping* 1978, 2: 237–245.
- [123] Balzamo E. Study of states of vigilance in adult *Papio cynocephalus*. *C R Seances Soc Biol Fil* 1973, 167: 1168–1172.
- [124] Astic L, Saucier D, Megirian D. Sleep circadian rhythm in rat kangaroo (*Potorous apicalis*): Effect of food distribution. *Physiol Behav* 1979, 22: 441–446.
- [125] Mendelson WB. The Octodon degu: a diurnal small mammal for sleep studies. *Sleep Res* 1982, 11: 89.