In brief

Cheap, fixed dose treatment is planned for malaria: The Drugs for Neglected Diseases Initiative (a non-profit drug development organisation) and Sanofi-Aventis have agreed to develop a new, fixed dose combination of artemisinin based compound to treat falciparum malaria in Africa. The target price of the treatment is less than \$1 (£0.53; €0.77) for a three day course.

US doctors rebuff drug watchdog group: The American College of Physicians has refused to allow the non-profit US organisation No Free Lunch, a group devoted to reducing competing interests in medicine, to rent a booth at its annual meeting. The college said the presence of the group would inhibit dialogue between industry exhibitors and doctors.

UK government to sponsor research into gel against HIV: The UK government will provide £24m (\$45m; €35m) to fund a Medical Research Council trial assessing a microbicide gel aimed at preventing HIV infection in women, particularly those who cannot insist on use of a condom. Starting this summer the researchers will follow 12 000 women for 39 months in four African countries: South Africa, Uganda, Tanzania, and Zambia.

Drug safety committees warn of food and drug interactions: The Committee on Safety of Medicines and the Food Standard Agency's committee on toxicity have issued draft guidance for consultation on the interaction between some drugs and certain foods. See www.food.gov.uk/multimedia/pdfs/tox200509.pdf

Cambodia has third death from avian flu: An eight year old girl has died from avian flu in Cambodia—the country's third death from the disease. All three cases were from Kamptot province, bordering Vietnam, where the disease has infected 33 and killed 15 people since late 2004. A World Health Organization spokeswoman said that the child's death may have been caused by contact with asymptomatic ducks rather than sick chickens.

Pfizer is asked to suspend sales of painkiller

Jeanne Lenzer *New York*

In a move that took experts by surprise, regulatory agencies in the United States and Europe have asked the drug maker Pfizer to suspend sales of its cyclooxygenase-2 (COX 2) inhibitor valdecoxib (Bextra).

The US Food and Drug Administration's move comes less than two months after an advisory panel recommended to the FDA that the drug remain on the market (*BMJ* 2005;330:440, 26 Feb).

Pfizer said it "respectfully disagreed" with the regulators' assessments of the risk-benefit ratio of the drug but agreed to "voluntarily" suspend sales.

Although the suspension of valdecoxib was welcomed by many experts, the decision by the FDA to go further than the European Medicines Agency by ordering "black box" warnings on all non-steroidal anti-inflammatory drugs drew criticism from a number of experts.

"To lump everything within a black box dilutes the message about COX 2s and sows confusion among practitioners and consumers," said Dr Garret FitzGerald, professor of cardiology at the University of Pennsylvania. He added, "I think the FDA has oscillated from saying there is no problem with anything to saying there's a problem with everything."

Dr Curt Furberg, a member of the FDA's data safety and risk management advisory committee, agreed with Dr FitzGerald, saying that although he was pleased with the decision to withdraw valdecoxib, important

differences existed among the non-selective non-steroidal anti-inflammatory drugs. He said that in certain studies the drug naproxen performed better than other non-steroidal anti-inflammatory drugs, so it was wrong to lump all these drugs together.

Some of the experts seemed to disagree over why Pfizer had been asked to withdraw valdecoxib from the market. Dr Steven Galson, acting director of the FDA's Center for Drug Evaluation and Research, said that there were concerns over rare but serious skin reactions ("FDA announces strong warnings for pain killers," *New York Times*, 8 April, <http://nytimes.com>).

But Dr Alastair Wood, chairman of the FDA's advisory panel on COX 2 inhibitors, said that it was cardiovascular risk, not skin disease, that caused the panel to have concerns about valdecoxib's safety. □

Wet nursing increases risk of HIV infection among babies

Pat Sidley *Johannesburg*

South African children using public hospitals are exposed to HIV infection because of a lack of infection control and because some babies are breast fed by women who are not their mothers and who are HIV positive, a new study says.

The study was conducted by the country's Human Sciences Research Council, Medical Research Council, and others for the Nelson Mandela Foundation.

It was commissioned after a previous study by the same group in 2002 found an unusually high proportion of children between the ages of 2 and 9 years infected with HIV. The suspicion had been voiced at the time that infection during these years may have been caused by particularly high rates of sexual abuse of children in South Africa. However, the study's authors, Dr Olive Shisana and

colleagues, say that for ethical and legal reasons they did not look into the possibility of sexual abuse among the children in the study. They have instead concentrated on the possibility that HIV infection was contracted in hospitals.

The study was conducted at several public hospitals in the Free State province and looked at dental facilities, maternity wards, and paediatric wards. In all these settings the researchers found that HIV could potentially be contracted.

Nearly 30% of the milk to be fed to babies tested positive for HIV viral RNA. Almost half (47%) of the instruments to be used clinically on children and a quarter of instruments to be used in children's mouths and gums had traces of blood on them.

The new finding of wet nursing as a mode of HIV transmission in South Africa followed from the finding that 1.7% of children in the study were breast fed by women who weren't their mothers. HIV positive children were 17 times more likely than HIV negative children to have been breast fed by a woman other than their mother. □

A South African child with AIDS is comforted by a child psychologist. Some children with AIDS have contracted HIV through wet nursing

HIV Risk Exposure in Children Aged 2-9 Years Served by Public Health Facilities in the Free State, South Africa is available at www.hsac.ac.za