Protective effects of carbenoxolone are associated with attenuation of oxidative stress in ischemic brain injury

Lang Zhang^{1,2#}, Yu-Min Li^{2#}, Yu-Hong Jing^{1,3}, Shao-Yu Wang^{1,4}, Yan-Feng Song¹, Jie Yin¹

¹*Institute of Anatomy, Histology and Embryology, School of Basic Medical Sciences, Lanzhou University, Lanzhou 730000, China*

²*Key Laboratory of Digestive System Tumors of Gansu Province, The Second Hospital of Lanzhou University, Lanzhou 730030, China*

³*Key Laboratory of Preclinical Study for New Drugs of Gansu Province, School of Basic Medical Sciences, Lanzhou University, Lanzhou 730000, China*

⁴*Division of Radiology, Tumor Hospital of Gansu, Lanzhou 730050, China*

#These authors contributed equally to this work.

Corresponding author: Yu-Hong Jing. E-mail: jingyh@lzu.edu.cn

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2013

ABSTRACT

Accumulating evidence has suggested that the gap junction plays an important role in the determination of cerebral ischemia, but the underlying mechanisms remain to be elucidated. In this study, we assessed the effect of a gap-junction blocker, carbenoxolone (CBX), on ischemia/reperfusion-induced brain injury and the possible mechanisms. By using the transient cerebral ischemia model induced by occlusion of the middle cerebral artery for 30 min followed by reperfusion for 24 h, we found that pre-administration of CBX (25 mg/kg, intracerebroventricular injection, 30 min before cerebral ischemic surgery) diminished the infarction size in rats. And this was associated with a decrease of reactive oxygen species generation and inhibition of the activation of astrocytes and microglia. In PC12 cells, H₂O₂ treatment induced more coupling and apoptosis, while CBX partly inhibited the opening of gap junctions and improved the cell viability. These results suggest that cerebral ischemia enhances the opening of gap junctions. Blocking the gap junction with CBX may attenuate the brain injury after cerebral ischemia/reperfusion by partially contributing to amelioration of the oxidative stress and apoptosis.

Keywords: gap junction communication; cerebral ischemia; reactive oxygen species

INTRODUCTION

Gap junctions exist ubiquitously in cell membranes as intercellular channels and are composed of transmembrane proteins called connexins (Cxs), which, when open, permit the intercellular diffusion of ions and small molecules^[1,2]. Gap junctions have numerous functions in the brain, such as the synchronization of electrical activity between neurons and the regulation of metabolic action between astrocytes^[3,4].

Cx43 hemichannels in astrocytes are susceptible to metabolic inhibition or in solutions free of divalent cations, which suggests that gap-junction communication is determined by the metabolic state^[5,6]. Gap junctions remain open after cerebral ischemia^[7], suggesting that gap-junction communication among astrocytes is involved in the pathological process after ischemic insult. The open gap junctions accelerate the spreading depression and induce cell death during ischemia or hypoxia^[8]. On the other hand, however, communication through gap junctions is important in buffering ionic shifts and the exchange of metabolites generated during hypoxia or ischemia. Recent evidence suggests that this could be a protective strategy against

hypoxic or ischemic insults^[9].

Recent studies have suggested that reducing gap-junction coupling with compounds results in decreased neuronal sensitivity to ischemic injury. However, many of the compounds used *in vivo*, such as octanol, heptanol and halothane, have major side-effects^[10–12], and interfere with synaptic transmission^[13,14]. Most of their neuroprotective effects in cerebral ischemia depend not only on blocking gap junctions, but also other actions. In this study, we attempted to explore whether specific gap-junction blockers affect the cellular sensitivity to ischemic injury in rats and whether such blockade diminishes the diffusion of reactive oxygen species (ROS) *in vitro*. Carbenoxolone (CBX) is effective in blocking gap-junction coupling^[15], interferes little with synaptic function *in vivo* and *in vitro*^[16–19], and induces minimal changes in intrinsic neuronal characteristics^[20,21]. In addition, this agent has been widely used as a mineralocorticoid agonist to inhibit 11-beta hydroxysteroid dehydrogenase and block gap-junction communication in various experimental studies^[15,22,23].

Currently, the effect of CBX on ischemic damage remains to be elucidated^[24,25]. Perez *et al.* reported that pretreatment with CBX, *via* a cannula implanted into the hippocampus in one hemisphere, results in decreased apoptosis in a global model of transient ischemia in rats^[26]. Another study suggested that blocking gap-junctional communication with CBX or downregulating Cx43 increases the survival of pyramidal neurons in the ipsilateral hippocampus and improves behavioral scores in the rodent model of middle cerebral artery occlusion (MCAO)^[27]. However, Tamura *et al.* reported that CBX accelerates the initiation and propagation of cortical spreading depression and worsens the outcome of focal cerebral ischemia^[28]. Furthermore, some evidence has shown that CBX cannot pass the blood-brain-barrier very efficiently^[29,30]. Therefore, the current study was performed to investigate *in vivo* the effects of CBX on transient cerebral ischemia through intracerebral ventricular injection, and further explore whether it can decrease ROS production and glial activation. To investigate whether CBX can block the gap-junction coupling associated with ROS diffusion *in vitro*, we mimicked oxidative injury with hydrogen peroxide and evaluated the coupling by the scrape-loading method in PC12 cells.

MATERIAL AND METHODS

Animals

Adult male Sprague-Dawley rats weighing 270–300 g from the Animal Center of Lanzhou University were used. All animals were kept in standard cages at 22–24°C with 40–60% humidity and lights on from 07:00 to 19:00. The protocols were approved by the Experimental Animal Committee of Lanzhou University. All efforts were made to minimize animal suffering and reduce the number of animals used.

Reagents

Carbenoxolone, cresyl violet and diaminobenzidine were from Sigma (St. Louis, MO); the mouse monoclonal antibodies anti-glial fibrillary acidic protein (GFAP), anti-CD11b, and anti- β -actin were from Santa Cruz Biotechnology (Santa Cruz, CA); and rabbit polyclonal anti-Cx43 antibody was from Sigma. Other reagents were from Biotechnology of Shanghai (Shanghai, China) unless specified otherwise.

Cerebral Ischemia

MCAO was established by the intraluminal filament method as described previously^[31]. Briefly, animals were anesthetized by 10% chloral hydrate (360 mg/kg, *i. p.*) and the right common and external carotid arteries were exposed. A nylon thread (4-0) was carefully inserted into the internal carotid artery and advanced towards the origin of the middle cerebral artery, resulting in MCAO. During the process of ischemia, laser Doppler perfusion monitoring was used to measure the blood flow in the trunk of the middle cerebral artery. A blood-flow drop to 60% of the baseline indicated successful blockage of the middle cerebral artery. After 30 min of MCAO, reperfusion was achieved by withdrawing the filament from the artery. After recovery from anesthesia, animals were kept in single cages for 24 h. Rectal temperature was maintained at $37 \pm 0.5^\circ\text{C}$ throughout using an electric blanket. The sham group received only incision of the cervical skin to expose the common carotid artery, without occlusion of blood flow.

Carbenoxolone Administration

The MCAO rats were randomly divided into two groups (12 rats each group). The CBX group received

intracerebroventricular injection of CBX (25 µg/kg, 5 µL) 30 min before MCAO, and the vehicle group received an equal volume of saline. Animals were fixed in a stereotaxic apparatus, and CBX or saline was injected into the right cerebral ventricle at the following coordinates: bregma: AP -0.8 mm, R +1.6 mm (midline) and depth 3.4 mm from dura. Animals were sacrificed after 24 h of reperfusion.

Measurement of ROS

Four rats from each group were randomly selected for ROS measurement. The ipsilateral striatum was dissected out and minced on ice, incubated with 0.25% trypsin at 37°C for 1 h, and washed with Dulbecco's modified Eagle's medium (DMEM) containing 10% FBS to harvest the cellular suspension. Free-radical production was measured by the fluorescence method using 2', 7'-dichlorodihydrofluorescein diacetate (DCFH-DA). DCFH-DA can cross cell membranes and is hydrolyzed by cellular esterases, then is oxidized to the fluorescent 2',7'-dichlorofluorescein (DCF) in the presence of peroxides. Accumulation of DCF was measured using a spectrofluorometer (Shimadzu Corp., Japan).

Measurement of Infarct Size

Coronal sections were cut serially (25 µm thick, at 250-µm intervals) from 4% paraformaldehyde-fixed brain and stained with cresyl violet. Infarct areas were measured and infarct volume was calculated by integration of the infarct area in each section along the rostro-caudal axis.

Immunohistochemistry

For Cx43, GFAP and CD11b immunostaining, sections (at 0.12 to 2.12 mm from bregma) were incubated with 0.3% H₂O₂ for 30 min and then placed in blocking buffer containing 10% normal goat serum and 0.3% Triton X-100 in 0.01 mol/L PBS (pH 7.2) for 30 min at 37°C and incubated with rabbit polyclonal anti-Cx43 (1:400), mouse monoclonal anti-GFAP (1:1 000) or anti-CD11b (1:200) antibody overnight at 4°C. Sections were then incubated with the corresponding biotinylated secondary antibody (1:400, 2 h at 37°C) followed by streptavidin-peroxidase (1:400, 1 h at 37°C). The same treatment omitting the primary antibodies served as a negative control.

Cell Counts

Total cell numbers were counted in a double-blinded

manner. In these experiments, CD11b- and GFAP-positive cells were counted under a 20× objective in the ipsilateral striatum from three sections (at 0.12, 1.12, and 2.12 mm from bregma) from each rat, in four randomly-selected non-overlapping areas (119 × 93 µm²) in each section.

Cell Culture

PC12 cells were cultured in DMEM containing 10% fetal bovine serum and antibiotics (penicillin, 100 IU/mL; streptomycin, 100 µg/mL). The cultures were maintained at 37°C in 95% air-5% CO₂ in a humidified incubator. Cell viability was evaluated with the MTT assay using a microplate reader (Bio-Rad Laboratory, Hercules, CA).

H₂O₂ Treatment

The culture medium was removed and PC12 cells were washed with PBS, then incubated for 20 min in the same medium, followed by incubation for a further 4 h in the presence of 100 mmol/L H₂O₂ at 37°C with or without CBX (dissolved in 0.01 mol/L PBS at a final concentration of 100 or 200 µmol/L).

Scrape-Loading

Gap-junction permeability was determined using the Lucifer Yellow (LY) scrape-loading technique^[32]. Confluent cells in 35-mm dishes were washed with PBS containing Ca²⁺ and Mg²⁺. A scrape line was made using a surgical blade in PBS containing 0.1% LY for 1 min. Cells were then washed again with external solution (in mmol/L: 140 NaCl, 5.5 KCl, 1.8 CaCl₂, 1.0 MgCl₂, 10 HEPES, pH 7.2). LY was allowed to diffuse in the external solution for 5 min. Thereafter, the culture dishes were photographed under a fluorescence microscope. The fluorescent areas were quantified in five consecutive fields using ImageJ (NIH).

Western Blot

Four rats were selected randomly from each group, anesthetized with pentobarbital, and the ipsilateral striatum was collected on ice. Samples were frozen in liquid nitrogen, homogenized, and total protein was extracted using RIPA buffer containing protease inhibitors. Harvested PC12 cells were lysed with RIPA buffer. Proteins (30 µg) were fractionated on 10% SDS-PAGE and transferred onto PVDF membranes. Membranes were blotted with anti-Cx43 (1:1 000) or anti-β-actin (1:5 000) antibody, followed by horseradish peroxidase-conjugated secondary

antibody (1:5 000). The immunoreactive protein bands were visualized by enhanced chemiluminescence.

Acridine Orange/Ethidium Bromide (AO/EB) Fluorescence Staining

Cellular morphological changes were observed by AO/EB staining using fluorescence microscopy. PC12 cells were harvested after different treatments, pooled, pelleted and resuspended in 200 μ L medium. A mixture of the fluorescent dyes (8 μ L) containing 100 μ g/mL AO and 100 μ g/mL EB was added to the cells and mixed gently. A drop of the mixture on a microscope slide was covered with a coverslip. The cells were visualized under a fluorescence microscope using a blue filter. At least 200 cells were counted in random fields. The percentage of apoptotic cells was calculated. The tests were repeated three times.

Statistical Analysis

All data are expressed as mean $\pm$ SEM. Differences between groups were analyzed by one-way ANOVA. For the influence of CBX on the infarct size relative to whole-brain volume (%), the Mann-Whitney U test was used for each evaluation. For other data, unpaired Student's *t* test was used. *P* < 0.05 was considered statistically significant.

RESULTS

CBX Pre-treatment Reduced Infarct Size Following MCAO

To evaluate the effect of CBX on brain damage after cerebral ischemia, we measured the infarct size by cresyl violet staining. CBX treatment significantly reduced the infarct size compared with untreated rats (Fig. 1A, B).

CBX Pre-treatment Reduced Cx43 Expression and ROS Production after Cerebral Ischemia

To examine the changes of gap junctions after cerebral ischemia, we determined the Cx43 expression in the ipsilateral striatum. Immunohistochemistry showed that Cx43 expression was higher in the ipsilateral striatum of ischemic rats than that in control rats. CBX treatment markedly decreased Cx43 expression in ischemic rats (Fig. 2A). We also quantified the Cx43 level in the ipsilateral striatum using Western blot and found similar results (Fig. 2B, C). Because cerebral ischemia can cause the accumulation of ROS, we measured the DCF fluorescence in the ipsilateral striatum and found that it was higher after cerebral ischemia, and the production of ROS was attenuated by CBX pre-treatment (Fig. 2D).

Fig. 1. CBX administration reduced the infarct size following MCAO. A: Representative images of infarcts stained with cresyl violet. Closed outlines denote the infarct foci. B: Infarct area (A_1), contralateral hemisphere area (A_2) and total brain area (A_3) on each coronal slice were directly measured by volume-analysis software, and the volumes were quantified by the formula: $V_{1,2,3} = \Sigma A_{1,2,3} \times 2$. Percentage of infarct volume was calculated as follows: Percentage of infarct volume = $[V_2 - (V_3 - V_2 - V_1)] / V_2 \times 100\%$. **P* < 0.05 (*n* = 6).

Fig. 2. CBX treatment reduced Cx43 expression and ROS production after cerebral ischemia. **A:** Immunohistochemical staining for Cx43 from bregma level 1.12 mm in the ipsilateral striatum. Arrows indicate positive immunoreactive staining. **B** and **C:** Western blots of Cx43 levels in the ipsilateral striatum. **D:** Hydrogen peroxide generation determined by DCF fluorescence. ^{##} $P < 0.01$ compared with control, ^{*} $P < 0.05$ compared with MCAO + vehicle group ($n = 4$).

Fig. 3. CBX treatment inhibited the activation of glial cells after cerebral ischemia. **A:** Representative images of GFAP-positive cells in the ipsilateral striatum. **B:** Positive cell-counts in three sections at 0.12, 1.12 and 2.12 mm from bregma. **C:** Representative images of CD11b-positive cells in these ipsilateral striatum. **D:** Positive cell-counts in three sections at 0.12, 1.12 and 2.12 mm from bregma. ^{##} $P < 0.01$ compared with control, ^{*} $P < 0.05$ compared with MCAO + vehicle group ($n = 6$).

CBX Treatment Inhibited the Activation of Glial Cells after Cerebral Ischemia Previous studies indicated that Cx43 is mainly localized in glia, and ischemia usually causes the activation of astrocytes and microglia, so whether blocking gap junctions affects glial activity after

cerebral ischemia is of interest. Immunohistochemistry showed that the number of GFAP-positive astrocytes induced by ischemia in the ipsilateral striatum was decreased by CBX treatment (Fig. 3A, B). Likewise, CBX-treated rats had fewer CD-11b-positive microglia than the untreated rats after ischemia (Fig. 3C, D).

CBX Inhibited the Gap Junction Coupling and Improved the Viability of PC12 Cells after H_2O_2 Treatment

The above results *in vivo* showed that CBX protected against transient cerebral ischemia partly due to a decrease of ROS and inhibition of Cx43 expression, but whether ROS result in the coupling of gap junctions and CBX reverses it were unknown. Therefore, PC12 cells were treated with H_2O_2 to mimic ROS injury. The scrape loading test showed that diffusion of LY into PC12 cells was increased after H_2O_2 treatment, while 100 $\mu\text{mol/L}$ and 200 $\mu\text{mol/L}$ CBX both clearly inhibited the diffusion of LY (Fig. 4A, B). The cell survival was also improved by treatment with CBX under H_2O_2 stress (Fig. 4C).

CBX Suppressed Cx43 Expression in PC12 Cells after H_2O_2 Treatment

Recent studies have shown that Cx43 is distributed in PC12 cells after induction^[33]. To address whether CBX

Fig. 4. CBX inhibited gap-junction coupling and improved the viability of PC12 cells after H_2O_2 treatment. **A:** Representative images of diffusion of LY into PC12 cells. CBX100, 100 $\mu\text{mol/L}$ CBX; CBX200, 200 $\mu\text{mol/L}$ CBX. Diffusion areas are shown in **B**. **C:** Cell viability was assessed using the MTT method. H_2O_2 treatment decreased viability, but this was reversed by CBX administration. [#] $P < 0.05$, ^{##} $P < 0.01$ compared with control; ^{**} $P < 0.01$ compared with H_2O_2 treatment ($n = 3$).

Fig. 5. CBX suppressed Cx43 expression in PC12 cells after H_2O_2 treatment. **A:** Representative images of Western blots. **B:** Cx43 expression levels. CBX100, 100 $\mu\text{mol/L}$ CBX; CBX200, 200 $\mu\text{mol/L}$ CBX. [#] $P < 0.05$ compared with control, ^{*} $P < 0.05$ compared with H_2O_2 treatment ($n = 3$).

Fig. 6. CBX administration reduced the apoptotic ratio in H₂O₂-treated PC12 cells. A: Representative images of AO/EB staining. Arrows indicate apoptotic cells. B: The apoptotic ratio was calculated according to the following formula: Apoptotic ratio (%) = (number of early apoptotic cells + number of late apoptotic cells) / (number of all cells). #*P* < 0.05, ##*P* < 0.01 compared with control; *P* < 0.01 compared with H₂O₂ treatment (*n* = 3).**

inhibited Cx43 expression in H₂O₂-treated PC12 cells, we determined the Cx43 level using Western blot and showed that it was overexpressed in H₂O₂-treated cells compared with control. Moreover, CBX treatment significantly decreased the overexpression of Cx43 in PC12 cells with H₂O₂ stress (Fig. 5A, B).

CBX Administration Reduced the Apoptosis Ratio of PC12 Cells with H₂O₂ Treatment

In vitro, we found the CBX inhibited the extensive opening of gap junctions induced by H₂O₂ stress. To address whether CBX can protect PC12 cells against ROS injury, the apoptosis induced by H₂O₂ stress was measured by AO/EB staining. We found that the apoptosis ratio of PC12 cells was increased by 100 μmol/L H₂O₂ treatment for 4 h.

In addition, after PC12 cells were incubated with H₂O₂ and CBX together, the apoptosis ratio was clearly decreased (Fig. 6A, B).

DISCUSSION

So far, many studies have documented the roles of gap junctions in cerebral ischemia. But argument continues, partly due to different experimental models, leading to different and even opposing results and conclusions^[34-36]. *In vivo*, most of the studies were designed to down- or up-regulate specific connexins, then investigate the gap-junction effects and underlying mechanisms in cerebral ischemia. But they failed to link the connexin expression to the actual level of opening of gap junctions^[37]. *In vitro*, studies cannot duplicate the ischemic condition, although quantitative assays were performed to evaluate the extent of gap-junction coupling^[38]. In the present study, we found that ischemia-induced overexpression of Cx43 was accompanied by ROS accumulation and glial activation. Furthermore, the gap-junction blocker CBX inhibited Cx43 expression, decreased ROS accumulation, and ameliorated the activation of glia including microglia and astrocytes. Also, the infarct size was diminished with CBX injection into the lateral cerebral ventricle after transient cerebral ischemia/reperfusion. To further explore the relation between ROS and gap-junction communication, PC12 cells were treated with H₂O₂, which induced the extensive opening of gap junctions and overexpression of Cx43. However, the roles of Cx43 in the ischemic process remain controversial. Cx43 heterozygous-null mice and mice with astrocytic Cx43 removed by *cre* expression are more susceptible to infarct expansion after focal ischemia^[37,39,40]. Alternatively, when hypoxic-ischemic brain injury induces an acute inflammatory reaction and reactive astrogliosis associated with peri-lesional upregulation of Cx43, human umbilical cord blood cell transplantation accelerates the regression of inflammatory events, narrows the perilesional astrocytic wall, and leads to the downregulation of Cx43 expression^[41]. A mimetic peptide that blocks Cx43 hemichannels after perfusion into the lateral ventricle reduces seizure activity and improves the survival of oligodendrocytes and neurons after ischemia in sheep^[42].

In vitro, the gap-junction coupling induced by H₂O₂ treatment was partly counteracted by CBX treatment.

Further findings also indicated that the apoptosis induced by H_2O_2 was attenuated by CBX administration. These *in vivo* and *in vitro* results suggested that cerebral ischemia enhances gap-junction coupling associated with the production and accumulation of ROS. And ROS stress enhances the secondary brain injury after cerebral ischemia^[43–45]. Inhibition of gap junctions with CBX decreases the diffusion of ROS, limiting the secondary injury factors in the ischemic core, and protecting cells in the penumbra to attenuate the expansion of the infarct^[46,47]. Other studies also indicate that CBX acts partially as an antioxidant^[26].

Not only do ROS accumulate, but activation of glial cells also occurs after cerebral ischemia. The roles of activated glial cells in brain injury remain controversial; presumably, they have advantages and disadvantages. Glia may play complex roles in the development of neuropathology, for example in triggering secondary injury or promoting self-repair by secreting cytokines^[48–50]. Here, we found that CBX reduced the activation of microglia and astroglia localized in the penumbra after transient cerebral ischemia. However, the underlying mechanisms remain to be clarified. Most evidence indicates that Cx43 is localized in the cell membranes of activated astrocytes and microglia^[51,52]. Accumulation and diffusion of ROS may activate microglia and astrocytes, and these effects may be amplified by gap-junction communication^[53]. In addition, apoptosis signals can diffuse through gap junctions^[54]. Previous studies showed that CBX treatment decreases the numbers of TUNEL-positive neurons in global ischemia^[26] and its administration to ischemic pups prevents caspase-3 activation^[55]. In our experiment, H_2O_2 treatment induced apoptosis in PC12 cells accompanied by gap-junction coupling. And after blocking the gap junctions with CBX, the apoptosis ratio was attenuated under ROS stress. These results confirmed that gap junctions are involved in apoptosis progression in brain injury.

Therapeutic targets have mainly been focused on signals associated with secondary injury, such as inflammation, excitatory amino-acid toxicity, and ROS, after cerebral ischemia. Although gap junctions are only regarded as channels for the diffusion of small molecules, their functional status may affect the efficiency of the above signals and thus participate in the development of

neuropathology after cerebral ischemia. The functional pattern and structural changes in gap junctions under neuropathological conditions remain to be elucidated. Our study only investigated the effect of CBX at the beginning of ischemia, so further studies are needed to determine the protective effect of CBX in long-term ischemia.

In brief, the gap-junction blocker CBX reduced the infarct size after cerebral ischemia, partially by inhibiting the accumulation and diffusion of ROS, suppressing the activation of glial cells, and decreasing apoptosis.

ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (30872731) and Natural Science Foundation of Gansu Province, China (2008GS01435 and 1010RJZA114).

Received date: 2012-07-24; Accepted date: 2012-10-02

REFERENCES

- [1] Simon AM, Goodenough DA. Diverse functions of vertebrate gap junctions. *Trends Cell Biol* 1998, 8: 477–483.
- [2] Bennett MV. Electrical synapses, a personal perspective (or history). *Brain Res Brain Res Rev* 2000, 32: 16–28.
- [3] Bissiere S, Zelikowsky M, Ponnusamy R, Jacobs NS, Blair HT, Fanselow MS. Electrical synapses control hippocampal contributions to fear learning and memory. *Science* 2011, 331: 87–91.
- [4] Giaume C, Koulakoff A, Roux L, Holcman D, Rouach N. Astroglial networks: a step further in neuroglial and gliovascular interactions. *Nat Rev Neurosci* 2010, 11: 87–99.
- [5] Retamal MA, Cortes CJ, Reuss L, Bennett MV, Saez JC. S-nitrosylation and permeation through connexin 43 hemichannels in astrocytes: induction by oxidant stress and reversal by reducing agents. *Proc Natl Acad Sci U S A* 2006, 103: 4475–4480.
- [6] Ye ZC, Oberheim N, Kettenmann H, Ransom BR. Pharmacological "cross-inhibition" of connexin hemichannels and swelling activated anion channels. *Glia* 2009, 57: 258–269.
- [7] Thompson RJ, Zhou N, MacVicar BA. Ischemia opens neuronal gap junction hemichannels. *Science* 2006, 312: 924–927.
- [8] Cotrina ML, Kang J, Lin JH, Bueno E, Hansen TW, He L, *et al.* Astrocytic gap junctions remain open during ischemic conditions. *J Neurosci* 1998, 18: 2520–2537.
- [9] Lin JH, Yang J, Liu S, Takano T, Wang X, Gao Q, *et al.* Connexin mediates gap junction-independent resistance to cellular injury. *J Neurosci* 2003, 23: 430–441.

- [10] Rawanduzy A, Hansen A, Hansen TW, Nedergaard M. Effective reduction of infarct volume by gap junction blockade in a rodent model of stroke. *J Neurosurg* 1997, 87: 916–920.
- [11] Saito R, Graf R, Hubel K, Fujita T, Rosner G, Heiss WD. Reduction of infarct volume by halothane: effect on cerebral blood flow or perifocal spreading depression-like depolarizations. *J Cereb Blood Flow Metab* 1997, 17: 857–864.
- [12] Rami A, Volkmann T, Winckler J. Effective reduction of neuronal death by inhibiting gap junctional intercellular communication in a rodent model of global transient cerebral ischemia. *Exp Neurol* 2001, 170: 297–304.
- [13] Pocock G, Richards CD. Excitatory and inhibitory synaptic mechanisms in anaesthesia. *Br J Anaesth* 1993, 71: 134–147.
- [14] Rorig B, Klaus G, Sutor B. Intracellular acidification reduced gap junction coupling between immature rat neocortical pyramidal neurones. *J Physiol* 1996, 490 (Pt 1): 31–49.
- [15] Davidson JS, Baumgarten IM. Glycyrrhetic acid derivatives: a novel class of inhibitors of gap-junctional intercellular communication. Structure-activity relationships. *J Pharmacol Exp Ther* 1988, 246: 1104–1107.
- [16] Frantseva MV, Kokarotseva L, Naus CG, Carlen PL, MacFabe D, Perez Velazquez JL. Specific gap junctions enhance the neuronal vulnerability to brain traumatic injury. *J Neurosci* 2002, 22: 644–653.
- [17] Gajda Z, Gyengesi E, Hermesz E, Ali KS, Szente M. Involvement of gap junctions in the manifestation and control of the duration of seizures in rats *in vivo*. *Epilepsia* 2003, 44: 1596–1600.
- [18] Ross FM, Gwyn P, Spanswick D, Davies SN. Carbenoxolone depresses spontaneous epileptiform activity in the CA1 region of rat hippocampal slices. *Neuroscience* 2000, 100: 789–796.
- [19] Szente M, Gajda Z, Said Ali K, Hermesz E. Involvement of electrical coupling in the *in vivo* ictal epileptiform activity induced by 4-aminopyridine in the neocortex. *Neuroscience* 2002, 115: 1067–1078.
- [20] Frantseva MV, Kokarotseva L, Perez Velazquez JL. Ischemia-induced brain damage depends on specific gap-junctional coupling. *J Cereb Blood Flow Metab* 2002, 22: 453–462.
- [21] Schmitz D, Schuchmann S, Fisahn A, Draguhn A, Buhl EH, Petrasch-Parwez E, *et al.* Axo-axonal coupling. a novel mechanism for ultrafast neuronal communication. *Neuron* 2001, 31: 831–840.
- [22] Jellinck PH, Monder C, McEwen BS, Sakai RR. Differential inhibition of 11 beta-hydroxysteroid dehydrogenase by carbenoxolone in rat brain regions and peripheral tissues. *J Steroid Biochem Mol Biol* 1993, 46: 209–213.
- [23] Bani-Yaghoub M, Underhill TM, Naus CC. Gap junction blockage interferes with neuronal and astroglial differentiation of mouse P19 embryonal carcinoma cells. *Dev Genet* 1999, 24: 69–81.
- [24] Vakili A, Hosseinzadeh SA, Khorasani MZ. Peripheral administration of carbenoxolone reduces ischemic reperfusion injury in transient model of cerebral ischemia. *J Stroke Cerebrovasc Dis* 2009, 18: 81–85.
- [25] Khorasani MZ, Hosseinzadeh SA, Vakili A. Effect of central microinjection of carbenoxolone in an experimental model of focal cerebral ischemia. *Pak J Pharm Sci* 2009, 22: 349–354.
- [26] Perez Velazquez JL, Kokarotseva L, Sarbaziha R, Jeyapalan Z, Leshchenko Y. Role of gap junctional coupling in astrocytic networks in the determination of global ischaemia-induced oxidative stress and hippocampal damage. *Eur J Neurosci* 2006, 23: 1–10.
- [27] Xie M, Yi C, Luo X, Xu S, Yu Z, Tang Y, *et al.* Glial gap junctional communication involvement in hippocampal damage after middle cerebral artery occlusion. *Ann Neurol* 2011, 70: 121–132.
- [28] Tamura K, Alessandri B, Heimann A, Kempinski O. The effect of a gap-junction blocker, carbenoxolone, on ischemic brain injury and cortical spreading depression. *Neuroscience* 2011, 194: 262–271.
- [29] Leshchenko Y, Likhodii S, Yue W, Burnham WM, Perez Velazquez JL. Carbenoxolone does not cross the blood brain barrier: an HPLC study. *BMC Neurosci* 2006, 7: 3.
- [30] Zhang Y. Determination of furazolidone, carbenoxolone sodium and berberine hydrochloride in wei kang tablets by reversed-phase high performance liquid chromatography (RP-HPLC). *Se Pu* 2002, 20: 350–352.
- [31] Vakili A, Hosseinzadeh F, Sadogh T. Effect of aminoguanidine on post-ischemic brain edema in transient model of focal cerebral ischemia. *Brain Res* 2007, 1170: 97–102.
- [32] el-Fouly MH, Trosko JE, Chang CC. Scrape-loading and dye transfer. A rapid and simple technique to study gap junctional intercellular communication. *Exp Cell Res* 1987, 168: 422–430.
- [33] Nagy JI, Hossain MZ, Hertzberg EL, Marotta CA. Induction of connexin43 and gap junctional communication in PC12 cells overexpressing the carboxy terminal region of amyloid precursor protein. *J Neurosci Res* 1996, 44: 124–132.
- [34] Farahani R, Pina-Benabou MH, Kyrozis A, Siddiq A, Baradas PC, Chiu FC, *et al.* Alterations in metabolism and gap junction expression may determine the role of astrocytes as "good samaritans" or executioners. *Glia* 2005, 50: 351–361.
- [35] Belliveau DJ, Bani-Yaghoub M, McGirr B, Naus CC, Rushlow WJ. Enhanced neurite outgrowth in PC12 cells mediated by connexin hemichannels and ATP. *J Biol Chem* 2006, 281: 20920–20931.
- [36] Kozoriz MG, Bechberger JF, Bechberger GR, Suen MW,

- Moreno AP, Maass K, *et al.* The connexin43 C-terminal region mediates neuroprotection during stroke. *J Neuropathol Exp Neurol* 2010, 69: 196–206.
- [37] Nakase T, Sohl G, Theis M, Willecke K, Naus CC. Increased apoptosis and inflammation after focal brain ischemia in mice lacking connexin43 in astrocytes. *Am J Pathol* 2004, 164: 2067–2075.
- [38] Zvalova D, Cordier J, Mesnil M, Junier MP, Chneiweiss H. p38/SAPK2 controls gap junction closure in astrocytes. *Glia* 2004, 46: 323–333.
- [39] Siushansian R, Bechberger JF, Cechetto DF, Hachinski VC, Naus CC. Connexin43 null mutation increases infarct size after stroke. *J Comp Neurol* 2001, 440: 387–394.
- [40] Nakase T, Fushiki S, Naus CC. Astrocytic gap junctions composed of connexin 43 reduce apoptotic neuronal damage in cerebral ischemia. *Stroke* 2003, 34: 1987–1993.
- [41] Wasielewski B, Jensen A, Roth-Harer A, Dermietzel R, Meier C. Neuroglial activation and Cx43 expression are reduced upon transplantation of human umbilical cord blood cells after perinatal hypoxic-ischemic injury. *Brain Res* 2012, 1487: 39–53.
- [42] Davidson JO, Green CR, Nicholson LF, O'Carroll SJ, Fraser M, Bennet L, *et al.* Connexin hemichannel blockade improves outcomes in a model of fetal ischemia. *Ann Neurol* 2012, 71: 121–132.
- [43] Hugyecz M, Mracsko E, Hertelendy P, Farkas E, Domoki F, Bari F. Hydrogen supplemented air inhalation reduces changes of prooxidant enzyme and gap junction protein levels after transient global cerebral ischemia in the rat hippocampus. *Brain Res* 2011, 1404: 31–38.
- [44] Nakase T, Maeda T, Yoshida Y, Nagata K. Ischemia alters the expression of connexins in the aged human brain. *J Biomed Biotechnol* 2009, 2009: 147946.
- [45] Nakase T, Yoshida Y, Nagata K. Enhanced connexin 43 immunoreactivity in penumbral areas in the human brain following ischemia. *Glia* 2006, 54: 369–375.
- [46] Nodin C, Nilsson M, Blomstrand F. Gap junction blockage limits intercellular spreading of astrocytic apoptosis induced by metabolic depression. *J Neurochem* 2005, 94: 1111–1123.
- [47] Tanaka M, Yamaguchi K, Tatsukawa T, Nishioka C, Nishiyama H, Theis M, *et al.* Lack of Connexin43-mediated bergmann glial gap junctional coupling does not affect cerebellar long-term depression, motor coordination, or eyeblink conditioning. *Front Behav Neurosci* 2008, 2: 1.
- [48] Amantea D, Nappi G, Bernardi G, Bagetta G, Corasaniti MT. Post-ischemic brain damage: pathophysiology and role of inflammatory mediators. *FEBS J* 2009, 276: 13–26.
- [49] Takano T, Oberheim N, Cotrina ML, Nedergaard M. Astrocytes and ischemic injury. *Stroke* 2009, 40: S8–12.
- [50] Turner DA, Adamson DC. Neuronal-astrocyte metabolic interactions: understanding the transition into abnormal astrocytoma metabolism. *J Neuropathol Exp Neurol* 2011, 70: 167–176.
- [51] Garg S, Md Syed M, Kielian T. Staphylococcus aureus-derived peptidoglycan induces Cx43 expression and functional gap junction intercellular communication in microglia. *J Neurochem* 2005, 95: 475–483.
- [52] Retamal MA, Froger N, Palacios-Prado N, Ezan P, Saez PJ, Saez JC, *et al.* Cx43 hemichannels and gap junction channels in astrocytes are regulated oppositely by proinflammatory cytokines released from activated microglia. *J Neurosci* 2007, 27: 13781–13792.
- [53] Takeuchi H, Jin S, Wang J, Zhang G, Kawanokuchi J, Kuno R, *et al.* Tumor necrosis factor- α induces neurotoxicity via glutamate release from hemichannels of activated microglia in an autocrine manner. *J Biol Chem* 2006, 281: 21362–21368.
- [54] Udawatte C, Ripps H. The spread of apoptosis through gap-junctional channels in BHK cells transfected with Cx32. *Apoptosis* 2005, 10: 1019–1029.
- [55] de Pina-Benabou MH, Szostak V, Kyrozis A, Rempe D, Uziel D, Urban-Maldonado M, *et al.* Blockade of gap junctions in vivo provides neuroprotection after perinatal global ischemia. *Stroke* 2005, 36: 2232–2237.