·Correspondence·

Authors' response to "Maternal age as a potential explanation of the role of the L allele of the serotonin transporter gene in anxiety and depression in Asians"

Haixia Long^{1,2}, Bing Liu^{1,2}, Bing Hou^{1,2}, Chao Wang³, Jin Li^{1,2}, Tianzi Jiang^{1,2,3,4}

¹Brainnetome Center, ²National Laboratory of Pattern Recognition, Institute of Automation, Chinese Academy of Sciences, Beijing 100190, China

³Key Laboratory for NeuroInformation of Ministry of Education, School of Life Science and Technology, University of Electronic Science and Technology of China, Chengdu 610054, China

⁴Queensland Brain Institute, University of Queensland, Brisbane, QLD 4072, Australia

Corresponding author: Tianzi Jiang. E-mail: jiangtz@nlpr.ia.ac.cn

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2014

In the letter to the editor, Dr. Comings *et al.* proposed a potential explanation of our findings that the L allele rather than S allele of 5-HTTLPR was associated with higher anxiety levels and reduced amygdala-prefrontal cortex (PFC) connectivity in Han Chinese^[1], which demonstrated an 'allele reversal' in the genetics of the 5-HTTLPR gene in Asians *versus* Caucasians. The authors alleged that this 'allele reversal' might simply result from maternal age and suggested that we test this on our datasets. Unfortunately, we cannot because maternal age information was not included in our sample. However, from the following evidence we can conclude that this explanation cannot be applied to our findings^[1]. First, the main evidence used is the authors' previous study on an association of the DRD1 gene with OCD scores in individuals with Tourette syndrome/ADHD, which is completely different from our study^[1] because the diagnostic status might influence the association analysis between maternal age and platelet serotonin level. So the conclusion from their study cannot be applied to our findings. The authors also used an unpublished study of 5-HTTLPR genotyping on 98 subjects to support their interpretation. But this evidence is unreliable because there is too much uncertainty in such a small sample. For example, in that study it is not known whether the participants are from a healthy population, patients, or a mixture of both, what diseases they have (if any), and whether they are Asian or Caucasian. Second,

there is no clear criterion that defines old maternal age, because the effect of maternal age is considered as a continuum rather than a cut-off. Moreover, the age of women having their first child varies across countries rather than ethnicity. From OECD Family Database (2014, Paris, www.oecd.org/social/family/database) and the work of Matthews and Hamilton^[2], the mean maternal age at the birth of the first child in Japan falls between those in Germany and the USA. And individuals are relatively equally distributed throughout different maternal age groups in a large Chinese cohort: 4324 with a maternal age <25 years, 3550 between 25 and 29 years, and 3433 with a maternal age >29 years^[3]. Thirdly, previous studies in Japan and Korea also showed that the L allele rather than S allele of 5-HTTLPR is associated with depression and higher amygdala activity during an emotional processing task and in the resting-state^[4–9], while the S allele is linked to this effect in Caucasians^[10–16]. More importantly, the allele distribution of 5-HTTLPR variants is significantly different between Asians and Caucasians, with a higher S allele frequency in the former^[13]. Because previous imaging genetics studies on 5-HTTLPR also did not take maternal age into account, based on the same criteria, our current and further studies show that the L allele is associated with reduced amygdala-PFC connectivity and default-mode network connectivity^[1, 17], which is contrary to the findings in Caucasians^[18–20].

In summary, maternal age is not a driving factor for ethnic difference in the effect of 5-HTTLPR based on the current evidence, though it might be a factor that needs to be considered in future studies. Such studies in a large, multi-ethnic sample are needed to investigate the association between 5-HTTLPR, depression/anxiety, neural circuits, and maternal age. In particular, further studies can directly examine the effect of 5-HTTLPR on the anxiety/depression score and neural circuits in Asian and Caucasian populations with matched maternal ages, which can help us to understand whether there is a fundamental Asian-Caucasian difference or just the influence of maternal age. More research will likely provide multiple perspectives.

REFERENCES

- [1] Long H, Liu B, Hou B, Wang C, Li J, Qin W, *et al.* The long rather than the short allele of 5-HTTLPR predisposes Han Chinese to anxiety and reduced connectivity between prefrontal cortex and amygdala. *Neurosci Bull* 2013, 29: 4–15.
- [2] Matthews TJ, Hamilton BE. Delayed childbearing: more women are having their first child later in life. *NCHS Data Brief* 2009: 1–8.
- [3] Luo YL, Cheng YL, Gao XH, Tan SQ, Li JM, Wang W, *et al.* Maternal age, parity and isolated birth defects: a population-based case-control study in Shenzhen, China. *PLoS One* 2013, 8: e81369.
- [4] Kim DK, Lim SW, Lee S, Sohn SE, Kim S, Hahn CG, *et al.* Serotonin transporter gene polymorphism and antidepressant response. *Neuroreport* 2000, 11: 215–219.
- [5] Kim H, Lim SW, Kim S, Kim JW, Chang YH, Carroll BJ, *et al.* Monoamine transporter gene polymorphisms and antidepressant response in Koreans with late-life depression. *JAMA* 2006, 296: 1609–1618.
- [6] Yoshida K, Ito K, Sato K, Takahashi H, Kamata M, Higuchi H, *et al.* Influence of the serotonin transporter gene-linked polymorphic region on the antidepressant response to fluvoxamine in Japanese depressed patients. *Prog Neuropsychopharmacol Biol Psychiatry* 2002, 26: 383–386.
- [7] Zhang K, Xu Q, Xu Y, Yang H, Luo J, Sun Y, *et al.* The combined effects of the 5-HTTLPR and 5-HTT1A genes modulates the relationship between negative life events and major depressive disorder in a Chinese population. *J Affect Disord* 2009, 114: 224–231.
- [8] Lee BT, Ham BJ. Serotonergic genes and amygdala activity in response to negative affective facial stimuli in Korean women. *Genes Brain Behav* 2008, 7: 899–905.
- [9] Li S, Zou Q, Li J, Li J, Wang D, Yan C, *et al.* 5-HTTLPR polymorphism impacts task-evoked and resting-state activities of the amygdala in Han Chinese. *PLoS One* 2012, 7: e36513.
- [10] Osher Y, Hamer D, Benjamin J. Association and linkage of anxiety-related traits with a functional polymorphism of the serotonin transporter gene regulatory region in Israeli sibling pairs. *Mol Psychiatry* 2000, 5: 216–219.
- [11] Caspi A, Sugden K, Moffitt TE, Taylor A, Craig IW, Harrington H, *et al.* Influence of life stress on depression: moderation by a polymorphism in the 5-HTT gene. *Science* 2003, 301: 386–389.
- [12] Hoefgen B, Schulze TG, Ohlraun S, von Widdern O, Hofels S, Gross M, *et al.* The power of sample size and homogenous sampling: association between the 5-HTTLPR serotonin transporter polymorphism and major depressive disorder. *Biol Psychiatry* 2005, 57: 247–251.
- [13] Kiyohara C, Yoshimasu K. Association between major depressive disorder and a functional polymorphism of the 5-hydroxytryptamine (serotonin) transporter gene: a meta-analysis. *Psychiatr Genet* 2010, 20: 49–58.
- [14] Smeraldi E, Zanardi R, Benedetti F, Di Bella D, Perez J, Catalano M. Polymorphism within the promoter of the serotonin transporter gene and antidepressant efficacy of fluvoxamine. *Mol Psychiatry* 1998, 3: 508–511.
- [15] Hariri AR, Mattay VS, Tessitore A, Kolachana B, Fera F, Goldman D, *et al.* Serotonin transporter genetic variation and the response of the human amygdala. *Science* 2002, 297: 400–403.
- [16] Hariri AR, Drabant EM, Munoz KE, Kolachana BS, Mattay VS, Egan MF, *et al.* A susceptibility gene for affective disorders and the response of the human amygdala. *Arch Gen Psychiatry* 2005, 62: 146–152.
- [17] Long H, Liu B, Hou B, Wang C, Kendrick KM, Yu C, *et al.* A potential ethnic difference in the association between 5-HTTLPR polymorphisms and the brain default mode network. *Chin Sci Bull* 2014, 59: 1355–1361.
- [18] Pezawas L, Meyer-Lindenberg A, Drabant EM, Verchinski BA, Munoz KE, Kolachana BS, *et al.* 5-HTTLPR polymorphism impacts human cingulate-amygdala interactions: a genetic susceptibility mechanism for depression. *Nat Neurosci* 2005, 8: 828–834.
- [19] Pacheco J, Beevers CG, Benavides C, McGeary J, Stice E, Schnyer DM. Frontal-limbic white matter pathway associations with the serotonin transporter gene promoter region (5-HTTLPR) polymorphism. *J Neurosci* 2009, 29: 6229–6233.
- [20] Wiggins JL, Bedoyan JK, Peltier SJ, Ashinoff S, Carrasco M, Weng SJ, *et al.* The impact of serotonin transporter (5-HTTLPR) genotype on the development of resting-state functional connectivity in children and adolescents: a preliminary report. *Neuroimage* 2012, 59: 2760–2770.