Managing chronic diseases

Palliative care spectrum starts early on

EDITOR—Palliative care does not start when a patient is dying.¹ A whole spectrum of palliation begins from the day of diagnosis of a potentially terminal disease, such as soft tissue sarcoma.

Palliation, in the sense of needing to absorb what this means for the patient, becomes even then a medical imperative. The “How long have I got, Doc?” question may not be articulated, but it will be there and needs an empathetic response.

Curative treatment is sought, but—with sarcoma—lung metastases signal the end of that phase and the beginning of palliative treatment. The doctor and the patient know the dragon can be disabled but never slain. Palliation is then partnership, working to decide what may be gained at what cost. (This may apply during curative treatment: to take chemotherapy drugs now “just in case,” or to keep them for later?)

Palliative treatment can be long (in sarcoma up to 20 years or more), or it can be brief. Its ending is not solely a medical judgment. Doctors may decide they can do no more, but active patients use online support groups, seeking new treatment options across the world.

Patients may decide to end palliative treatment. Grown weary of painful chemotherapy offering diminishing returns, they may prefer a shorter but better life. Making that choice together is an aspect of palliation.

When death is near, palliative care becomes terminal care. If throughout doctors and patients have aligned their understanding of the disease, and have taken decisions jointly, have explored the holistic meaning of disease in life and death, the palliative spectrum may offer healing if not cure.

If, however, being palliative begins only when the patient is dying doctors cannot then meet patients, where they are most needed, in the ground of their being and their not being.

Peter Cox *sarcoma survivor*
Systems 4, Gidea Park, Essex RM2 6AH
peter.cox@systems4.co.uk

Competing interests: None declared.

1 Murray SA, Boyd K, Sheikh A. Palliative care in chronic illness. *BMJ* 2005;330:611-2. (19 March.)

End of life is more than death

EDITOR—In advanced, non-malignant, chronic illness, the proximity of death and the inadequacy of treatment to prevent it change the patient's circumstances. Accurate prognosis may be impossible and that also influences the circumstances. Although doctor and patient may know death is approaching, planning is needed for a stage of life that could continue for a year or more. For example, two year survival in patients with advanced chronic obstructive pulmonary disease is 50-64%.^{1,2} Murray et al warn against a paralysis of clinical decision making, but the inaccuracy of prognosis in these diseases presents complex challenges.³

Planning for death in advanced chronic disease may seem pessimistic when life still holds rich possibilities. Murray et al seem to use the term end of life as a euphemism for death.³ But the context of their editorial demands a broader understanding. Approaching the end of life has implications distinct from imminent death. These include the need to consider symptom management in preference to disease control, information about advanced disease, advance directives, and choice about place of care. These subjects are the business of palliative care but they need development in advanced disease without entailing for patient and carers the perceived finality of palliative and hospice care.

Much has been learnt about the attributes of a good death.⁴ This learning should be extended to that stage in advanced disease that precedes dying. The seven promises that Lynn et al propose should characterise care at the end of life

and provide a suitable template for the end of life before death becomes imminent.⁵ In the development of our understanding about the end of life in chronic disease we must learn from patients about their needs and how they should be met. These are likely to include talking about the end of life without necessarily preparing for imminent death.

Patrick White *senior lecturer*
Department of General Practice and Primary Care,
Guy's, King's and St Thomas' School of Medicine,
King's College London, London SE11 6SP
patrick.white@kcl.ac.uk

Competing interests: None declared.

- 1 Connors-AF J, Dawson NV, Thomas C, Harrell-FE J, Desbiens N, Fulkerson WJ, et al. Outcomes following acute exacerbation of severe chronic obstructive lung disease. The SUPPORT investigators (study to understand prognoses and preferences for outcomes and risks of treatments). *Am J Respir Crit Care Med* 1996 Oct;154(4 Pt 1):959-67.
- 2 Almagro P, Calbo E, Ochoa de EA, Barreiro B, Quintana S, Heredia JL, et al. Mortality after hospitalization for COPD. *Chest* 2002;121:1441-8.
- 3 Murray SA, Boyd K, Sheikh A. Palliative care in chronic illness. *BMJ* 2005;330:611-2. (19 March.)
- 4 Steinhauser KE, Clipp EC, McNeilly M, Christakis NA, McIntyre LM, Tulsky JA. In search of a good death: observations of patients, families, and providers. *Ann Intern Med* 2002;132:825-32.
- 5 Lynn J, Schuster JL, Kabacoff A. *Improving care for the end of life: a sourcebook for health care managers and clinicians*. Oxford: Oxford University Press, 2000.

Care of children with chronic illness teaches some lessons

EDITOR—Campbell and McGauley share insights from forensic psychiatry about the relationship between doctor and patient in chronic illness.¹ Their observations have relevance to a new and emerging issue in chronic care: the transition of young adults with chronic conditions from child centred (paediatric) to adult oriented health care.

Some young adults and families experience “culture shock” when transferring from paediatric to adult care.² Paediatrics is seen as supportive and compassionate, whereas adult care is perceived as formal, business-like, distant, rigid in therapeutic approach, and less interested in patients' perspectives. Campbell and McGauley's description of “negative” attitudes of the adult system towards patients with “incurable” conditions supports this notion of a difference in the culture of paediatrics and adult medicine. They also give guidance to adult providers who take over care of young adults regarding the importance of building a positive relationship with their patient; eliciting patients' perspectives on the illness; and developing flexible, individualised treatment plans.

Adult providers need to be aware that new young adult patients have ended what

bmj.com

Letters appearing here are an edited selection of rapid responses originally posted on bmj.com

We ask for all letters to the editor to be submitted as rapid responses via bmj.com

For advice see: bmj.com/rapidresponses

may have been a long term psychologically important relationship with paediatric providers and may need to help young adults achieve a sense of closure with previous providers before they can develop a positive relationship with new adult providers.

Campbell and McGauley focused their discussion on the patient as an individual. I suggest that those caring for patients with chronic conditions view the family, rather than the patient, as the unit of care. Relationships between families and doctors in chronic illness are especially important since the patient's "significant others" (spouse, children, parents, extended family) can have a major role in promoting adherence to the therapeutic regimen, monitoring health status, and identifying exacerbations of the chronic condition.

John Reiss *associate professor of paediatrics*
Institute for Child Health Policy, PO Box 100147,
University of Florida, Gainesville, FL 32610-0147, USA
jgr@ichp.ufl.edu

Competing interests: None declared.

- 1 Campbell C, McGauley G. Doctor-patient relationships in chronic illness: insights from forensic psychiatry. *BMJ* 2005;330:667-70. (19 March).
- 2 Reiss JG, Gibson RW, Walker LR. Health care transition: youth, family, and provider perspectives. *Pediatrics* 2005;115:112-20.

Funding follow-up is key to long term specialist care

EDITOR—Gask says that consultants do not manage most patients with chronic diseases even though they should have a key role.¹ She argues that the current NHS organisational structure is largely to blame.

The current service level agreement means that primary care trusts pay hospitals for referrals of new patients and per procedure. No fee has been set for follow-up, and fees for new patients are not varied, to take into account the chronic nature of a condition.

To preserve current levels of service, hospital clinics have to maintain an active discharge policy to stop clinics becoming overwhelmed with long term, non-funded follow-ups.

This can reduce job satisfaction in and patient satisfaction with hospital care. New funding arrangements will need to tackle this problem for secondary as well as primary care to achieve a better service for patients with chronic diseases.

Jon Norman *pain fellow*
South Manchester University Hospitals Trust,
Manchester M23 9LT
j.norman@smuht.nwest.nhs.uk

Competing interests: None declared.

- 1 Gask L. Role of specialists in common chronic diseases. *BMJ* 2005;330:651-3. (19 March).

Condition based payment may improve care of chronic illness in Canada

EDITOR—DiPiero and Sanders identify economic disincentives in payments models and propose fees-for-conditions to encourage innovation consistent with the chronic care model and to reward superior results.¹

In September 2003, chronic care practice enhancement payments were introduced in

British Columbia as part of implementing the chronic care model.² The patient's general practitioner became eligible to receive an annual payment of \$C75 (£32; \$60; €47) for each patient with a confirmed diagnosis of diabetes mellitus or congestive heart failure for whom a flow sheet maintained in the patient's chart documented care consistent with the approved provincial clinical practice guidelines.³ These payments are in addition to the normal fee for service payments for patient care.

Over half the family doctors in the province are now participating, and the payments give them financial flexibility to allow experimentation and innovation in practice redesign. Doctors have accepted payments tied to guidelines and the use of performance measures⁴ but are reluctant to see payments tied to actual results as they are often beyond their control.

Payment reform is a necessary but not sufficient condition to implement the chronic care model. Doctors need support through the process of change. Complementary elements include help with the development of patient registers; developing flow sheets as decision support tools; central storage of flow sheets electronically to enable sharing with other healthcare providers; and support for patient self management.

Our experience shows that small incremental changes to an existing fee for service system are acceptable and effective. Our next step, planned for this year, is to extend the range of chronic disease covered by the payments, and to continue to evaluate and report on the outcome.⁵

Howard Platt *director*
Medical Outcomes Improvement Branch, Clinical
Innovation and Integration Division, Ministry of
Health Services, 2-3, 1515 Blanshard Street,
Victoria, BC, Canada V8W 3C8
howard.platt@gems3.gov.bc.ca

Competing interests: HP is an employee of the British Columbia Ministry of Health Services.

- 1 DiPiero A, Sanders DG. Condition based payment: improving care of chronic illness. *BMJ* 2005;330:654-7. (19 March).
- 2 Ministry of Health Services. Full service family practice incentive program. www.healthservices.gov.bc.ca/cdm/practitioners/fsfpbooklet.pdf (accessed 5 Apr 2005).
- 3 Ministry of Health Services. *Clinical practice guidelines and protocols in British Columbia*. www.healthservices.gov.bc.ca/msp/protoguides/index.html (accessed 5 Apr 2005).
- 4 Ministry of Health Services. *People with diabetes and proportion receiving recommended services by age and gender, British Columbia, 2002/2003. Medical services plan*. www.healthservices.gov.bc.ca/cdm/research/diabetes_rec_servs_agegender_02-03.pdf (accessed 5 Apr 2005).
- 5 Ministry of Health Services. *Chronic disease management update June 2004*. www.healthservices.gov.bc.ca/cdm/research/updatejun04.pdf (accessed 5 Apr 2005).

Combination of inhaler treatment in India has shown good results

EDITOR—The long acting anticholinergic drug tiotropium and the long acting β -2 agonist formoterol have individually been shown to be effective in the management of chronic obstructive pulmonary disease (COPD).¹ Faster and prolonged action in case of formoterol and effective prolonged action of tiotropium make the two drugs with different profiles attractive alternative

options, and combining them seems justified for patients with stable COPD.^{2,3}

In India the combination of drugs is being used to good clinical benefit. People do not have to rely on multiple inhalers, which improves compliance with treatment. This is especially important for the vast illiterate population of India, for whom using inhalators to manage respiratory disorders is associated with taboos.

Most villagers believe that prescription of inhalers spells doom for the patient and that inhalers are prescribed only when the end is near. Doctors are finding it difficult to convince patients to use inhalers. Many doctors use the analogy "eyedrops for eyes, ear-drops for ears, topical ointments for skin disorders, and inhalers for respiratory tract disorders" to make the point of using inhaler treatment to deliver the drug directly to the site of action rather than through a circuitous route.

Patients often fall prey to quacks and unlicensed practitioners who dispense potentially harmful drugs, and they report with serious systemic side effects such as Cushing's syndrome.⁴ Having combination therapy available helps ensure that the correct form of treatment is dispensed to patients in a convenient and easy manner. Public health organisations have to work hard to raise awareness about the inhalers, and a convenient combination certainly helps ensure better compliance.

Parvaiz A Koul *ACP fellow*
North Shore University Hospital, Manhasset,
NY 11030, USA
parvaizk@rediffmail.com

Competing interests: None declared.

- 1 Cooper CB, Tashkin DP. Recent developments in inhaled therapy in stable chronic obstructive pulmonary disease. *BMJ* 2005;330:640-4. (19 March).
- 2 Cazzola M, Marco FD, Santus P, Boveri B, Verga M, Matera MG, et al. The pharmacodynamic effects of single inhaled doses of formoterol, tiotropium and their combination in patients with COPD. *Pulm Pharmacol Ther* 2004;17:35-9.
- 3 Tennant RC, Erin EM, Barnes PJ, Hansel TT. Long-acting beta 2-adrenoceptor agonists or tiotropium bromide for patients with COPD: is combination therapy justified? *Curr Opin Pharmacol* 2003;3:270-6.
- 4 Koul PA. Cushing's syndrome following herbal therapies for rheumatoid arthritis and bronchial asthma in India [electronic response to Doust J, Del Mar D. Why do doctors use treatments that do not work? *BMJ* 2004;328:474-5]. *bmj.com* 2004. <http://bmj.com/cgi/eletters/328/7438/474#51792> (accessed 14 Apr 2005).

bmj.com

Summary of webchat

EDITOR—The theme issue on managing chronic diseases deliberately focused on patients' needs, unlike the previous two on the subject, and several issues emerged in the accompanying webchat.¹

Was the focus on particular diseases the latest thinking in managing them? Patients' needs and generic self management skills might be more important than the disease they had to live with. However, the ability to use population perspectives and outreach programmes is helpful, as is including a public health perspective into training. Participants were unclear whether these approaches were mutually exclusive, but focusing on the population or the disease rather than the patient might be less threatening.

The global movement against chronic diseases has emphasised risk factors, making

health promotion and education and the role of the family important. On the one hand, this approach may stigmatise people with diseases such as HIV infection, given that people will always develop or contract chronic diseases—however well they avoid risk factors. On the other hand, powerful tools help to prevent morbidity once people are ill, which means that a two pronged approach—before and after chronic illness—is needed.

What is the role of “mutual aid groups” in helping people manage a chronic disease? Utilising patients’ generic skills, such groups often cater for several diseases. They sometimes start with professional leadership but all mature with service users taking responsibility. Observed outcomes include improved self efficacy, social networks, and communication skills, but only gradually. Some groups with users who have different diseases focus on a common problem because peer groups can help each other without having to focus on a particular disease. This approach ties in with the difficulty any programme or regular health service has in managing patients who have several chronic conditions simultaneously.

How useful are guidelines? Many evidence based guidelines for clinical care and public health interventions tackle one problem at a time, and in this way drive research protocols. Resources may not be available to plan differently. Awareness of this issue has been raised in journals, which could therefore influence research agendas. Indeed, the *BMJ* should promote broader research into people with multiple comorbidities to add valuable insights into service performance and enable policy makers to plan adequate specialist and primary care responses to the most common conditions.

Participants recommended that the next theme issue should emphasise comorbidities, and authors should be given feedback when important comorbidities for the condition they are studying have not been appropriately considered. And the problems caused by chronic diseases coexisting with infectious diseases in developing countries also warrant more column inches.

Birte Twisselmann *technical editor*
BMJ

Competing interests: None declared.

1 Webchat. Managing chronic diseases. 24 March 2005. <http://bmj.bmjournals.com/cgi/content/full/330/7493/DC1> (accessed 14 Apr 2005).

Time to consider employment conditions for carers

EDITOR—White reports the UK government’s plan to overhaul social care services.¹ The employment conditions of carers seem always to be omitted from discussion about service provision. Conditions of employment (if they exist) seem not to apply for those who work for care agencies.

Employees lack status and support. The job of a carer is perceived as little more than a modified form of normal housework. But

how many housewives are responsible for the organisation and basic cleanliness of 10 or 20 homes? The carer has to learn the likes and dislikes of the client and the geography of the client’s home, often also acting as a psychotherapist. People with pain can be fractious and difficult, and elderly people may not be paragons of charm and cheerfulness. A carer’s job can sometimes be thankless.

Many carers are unstinting in the amount of extra care they provide—shopping, for example, not just for a client’s basic necessities but also for presents for a client’s friend or relative. It might all be part of the job, but financial compensation for the time spent doing this is rare.

Carers are often called on at short notice to make extra visits—usually to cover for someone else—invariably at weekends. Not many people with families wish to work at weekends. Carers therefore often have little control over the amount of time they work, and many care agencies do not pay more for these unsocial hours. Carers must be among some of the lowest paid people in the country. Why does society expect cut price honesty, integrity, and selflessness?

The government proposals are to be welcomed, but implementation will fail unless the workforce is considered.

Cynthia M Lewis *retired doctor*
Hillcrest, Whatstandwell, Nr Matlock, Derbyshire
DE4 5HS
theamayl@aol.com

Competing interests: None declared.

1 White C. Government launches plan to overhaul social care services. *BMJ* 2005;330:687. (26 March.)

Insulin resistance and depression

Difference in patient groups could have been explained by other factors

EDITOR—The overlap between mood disorders and diabetes is an important area of research because a fuller understanding of the pathophysiology of insulin resistance and how this impacts on mood could lead to new insights and treatments for depression.¹ Timonen et al reported a positive correlation between insulin resistance and severity of depressive symptoms that was more obvious in patients with impaired glucose tolerance than patients with type 2 diabetes mellitus.² However, this finding is tentative and may need to be viewed with caution.

The difference between controls and patients with impaired glucose tolerance on Beck depression inventory scores, although significant, was small and is of debatable relevance: controls had a median score of 5, and patients had a median score of only 6.² A score of less than 10 on the Beck depression inventory is within the normal range, and a score of 19 is considered the threshold for moderate depression.³ Although the authors tried to control for factors such as body mass index, smoking, alcohol, exercise, and sex, a difference of only one point on

the Beck depression inventory (in subjects aged between 61 and 63) could easily be explained by other important variables, including any current or previous psychiatric illness, comorbid physical illness, current drug treatment, personality factors, or even subtle cognitive impairment.

Daniel J Smith *specialist registrar in psychiatry*
Southern General Hospital, Glasgow G51 4TF
daniel.smith@glacomen.scot.nhs.uk

Competing interests: None declared.

- 1 Musselman DL, Betan E, Larsen H, Phillips LS. Relationship of depression to diabetes types 1 and 2: epidemiology, biology, and treatment. *Biol Psychiatry* 2003;54:317-29.
- 2 Timonen M, Laakso M, Jokelainen J, Rajala U, Meyer-Rochow VB, Keinänen-Kiukkaanniemi S. Insulin resistance and depression: cross sectional study. *BMJ* 2005;330:17-8. (1 January.)
- 3 Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561-71.

Authors’ reply

EDITOR—The median Beck depression inventory score was 6 for both impaired glucose tolerance and type 2 diabetes, whereas a score of <10 is considered the threshold for non-depression.¹ We emphasise, however, that although people with type 2 diabetes have an increased prevalence of depression, most of them remain non-depressed.

We subjected our study population (n=491), in which previously diagnosed diabetes was excluded, to some additional analyses. We dichotomised the score on the Beck depression inventory into at least mild depression (score ≥10) and non-depression (score <10).¹ We used the World Health Organization’s cut-off point to define insulin resistance as the lowest fourth of the qualitative insulin sensitivity check index (QUICKI) of the general population.² Along with our earlier confounders, we also used information on comorbid physical illnesses, as well as cognitive impairment, defined by verbal fluency test,³ as confounding variables in a logistic regression analysis. Insulin resistance increased the risk for depression twofold after adjusting for confounders (odds ratio 2.04, 95% confidence interval 1.06 to 3.90).

These analyses did not alter the main findings of our paper. However, as in any large population sample, only some of the potential confounding variables could be considered. None the less, we believe our results are reliable and supplement earlier reports that implied a positive association between insulin resistance and depression at population level in the age bracket 61-63, before any clinical diagnosis of type 2 diabetes.⁴

Markku Timonen *acting professor*
mtimon@sun3.oulu.fi

Ulla Rajala *research fellow*
Department of Public Health Science and General Practice, University of Oulu, Box 5000, FIN-90014 Oulu, Finland

Mauri Laakso *senior lecturer*
Jari Jokelainen *biostatistician*
Unit of General Practice, Oulu University Hospital, 90029 OYS, Finland

Competing interests: None declared.

- 1 Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561-71.
- 2 Tenerz A, Norhammar A, Silveira A, Hamsten A, Nilsson G, Ryden L, Malmberg K. Diabetes, insulin resistance, and the metabolic syndrome in patients with acute myocardial infarction without previously known diabetes. *Diabetes Care* 2003;26:2770-6.
- 3 Borkowski JG, Benton AL, Spreen O. Word fluency and brain damage. *Neuropsychologia* 1967;5:135-40.
- 4 Ramasubbu R. Insulin resistance: a metabolic link between depressive disorder and atherosclerotic vascular diseases. *Med Hypotheses* 2002;59:537-51.

Details of the two other authors are on bmj.com

Delayed hypersensitivity due to epidural block with ropivacaine

Report raises several issues

EDITOR—True allergy to amide type local anaesthetic drugs is extremely rare, and, because of the serious implications for the patient (for example, dentistry without pain control), it is one that must be made with great care and by excluding all other possible causes. In testing their patient's sensitivity with a preservative free solution from an ampoule without a latex rubber stopper, Ban and Hattori were obviously aware of the need to avoid those two common confounders, and the case that they present seems definitive, but several aspects of their report concern me.¹

Firstly, their patient seems to have received a daily dose of epidural ropivacaine of 96-192 mg for a period of two weeks. This is considerably in excess of the approved maximum duration of continuous administration of this drug (72 hours), but this is not mentioned. Nor is there any discussion of the possible role of cumulative local anaesthetic toxicity in the generation of the hypersensitivity, or of the possibility that "plasticisers" from the containers might have played a part, given the very large volume of solution that the patient must have received in a two week period.

Secondly, no detail is given of the technical aspects of the "intradermal" test.

Thirdly, the authors say that ropivacaine may cause allergic reactions characterised by signs such as urticaria, angioneurotic oedema, but I know of no published reports of such features. Can they provide definitive references, please?

John A W Wildsmith *professor of anaesthesia*
University Department of Anaesthesia, Ninewells Hospital and Medical School, Dundee DD1 9SY
j.a.w.wildsmith@dundee.ac.uk

Competing interests: JAWW has received research funding from AstraZeneca, although he has had no contact with the company about this report.

- 1 Ban M, Hattori M. Delayed hypersensitivity due to epidural block with ropivacaine. *BMJ* 2005;330:229. (29 January.)

Authors' reply

EDITOR—Continuous epidural analgesia is well known to shorten the duration of pain associated with herpes zoster. Usually 0.5% bupivacaine is infused for about two weeks. Continuous epidural infusion of 0.2% ropivacaine 4-10 ml/h for several days for postoperative analgesia is widely approved.

Two weeks' administration of ropivacaine is not excessive.

We did not perform the challenge test. It is difficult to conclude whether the reaction to ropivacaine in our case is dose dependent.

We tested not only 0.2% ropivacaine but also 1% lignocaine without adrenaline (epinephrine) as control. The skin injection of 1% lignocaine induced no reaction. Both anaesthetics were given from plastic ampoules without a rubber stopper. We did not think that plasticisers from the containers caused allergic reaction in our patient. The opinion of Wildsmith is worthy of consideration. Pharmaceutical companies should pay attention to the material of plastic containers.

As mentioned above, intradermal tests of 0.2% ropivacaine and 1% lignocaine were performed. We did not disinfect the skin and injected 0.1 ml of each drug. Only 0.2% ropivacaine resulted in erythema 8-72 hours after the injection. Two healthy men showed negative intradermal tests with 0.2% ropivacaine at the same time.

Ropivacaine may cause allergic reactions characterised by urticaria, angioneurotic oedema, tachycardia, or vomiting. Anaphylactoid reactions to amide-type local anaesthetics such as lignocaine, mepivacaine, prilocaine, bupivacaine, and etidocaine are not uncommon.

We informed AstraZeneca of our case on the 10 February 2004.

We think that continuous epidural block with ropivacaine will be widely performed because of its low cardiotoxicity.

Masanori Ban *chief, dermatology section*
masanoriban@k5.dion.ne.jp

Masahito Hattori *chief, anaesthesiology section*
Hashima City Hospital, 3-246 Shinseicho, Hashima City 501-6206, Japan

Competing interests: None declared.

Will the lead clinician please stand up?

That'll be the general practitioner

EDITOR—In his personal view Gannon shows that a lead clinician with clear ownership of patients' cases is crucial in ensuring continuity of care.¹ I know who the lead clinician is—the general practitioner. Secondary care now seems so refined and pigeonholed that there are no remaining generalists, excepting accident and emergency and geriatricians, left in hospitals. I have been asked by a cardiologist in the left ventricular function clinic to refer a patient to the general cardiology clinic because the echocardiogram is fine but the patient might have angina.

I recognise Gannon's scenario and could describe several similar instances every year, thankfully not all with such sad ends.

The general practitioner is well placed to provide the overview, but the key problem hampering this role is the quality of communication he or she receives from secondary care. This falls into two main categories: prompt, but illegible and incomplete; and late, but typed and comprehensive.

If general practitioners are to be the lead clinicians, then give them prompt, legible, succinct information. That would be a start.

Maurice Conlon *general practitioner principal*
Ridgacre House Surgery, South Birmingham Primary Care Trust, Birmingham B32 2TJ
maurice.conlon@southbirminghampct.nhs.uk
Competing interests: MC knows Craig Gannon personally.

- 1 Gannon C. Will the lead clinician please stand up? *BMJ* 2005;330:737. (26 March.)

We should be promoting total quality care

EDITOR—Godlee's comment about the fragmented care described by Gannon in his personal view prompts us to provide additional evidence from our study at a district general hospital.^{1,2} As part of a research project we persuaded clinical teams to examine the quality of care in their own units.

We found evidence of adverse events or critical incidents in 60/154 medical cases and 27/134 surgical cases. Most of the issues were comparatively minor, commonplace problems that might be found in any hospital ward—for example, venflon cellulitis, over-infusion of intravenous fluids—but in at least 12 medical cases and 16 surgical cases we found evidence of a lack of integrated care, in some cases with serious consequences.

Gannon thinks that we need to revitalise the role of the lead clinician.² Our findings support his contention. However, is that possible and would it be enough? Over the past 20 years, hospital doctors have striven to become specialists rather than consultants. Twenty years ago Sir Christopher Booth, former director of medicine at the Royal Postgraduate Medical School, warned of the effects of technology on the practice of medicine,³ and today we live with such predictions.

In our study we tried to assess causation. Inevitably, superficial appraisals are subjective, but at least half of the problems seemed to arise from a "failure of overall care." The specialists of today do not have the time to be lead clinicians for all the patients they see. So should we not be training all members of the clinical team to take overlapping responsibility—developed as "total quality care" in the Japanese car industry and subsequently applied to medical care?⁴ And should we not nominate individual secondary care doctors to relate to individual patients to ensure the adequacy of integrated care?

Graham Neale *visiting professor*
g.neale@imperial.ac.uk

Sisse Olsen *research fellow*
Clinical Safety Research Unit, Academic Department of Surgery, Imperial College, St Mary's Hospital, London W2 1NY

Competing interests: None declared.

- 1 Godlee F. Heading where exactly [Editor's choice]? *BMJ* 2005;330:0-g. (26 March.)
- 2 Gannon C. Will the lead clinician please stand up? *BMJ* 2005;330:737. (26 March.)
- 3 Booth CC. What has technology done to gastroenterology? *Gut* 1985;26:1088-94.
- 4 Milakovich ME. Creating a total quality care environment. *Health Care Manage Rev* 1991;16:9-20.