Identification of autophagy signaling network that contributes to stroke in the ischemic rodent brain *via* gene expression

Kun Liang¹, Lei Zhu¹, Jinyun Tan¹, Weihao Shi¹, Qing He¹, Bo Yu²

¹Department of Vascular Surgery, Huashan Hospital, Fudan University, Shanghai 200040, China

²Department of Vascular Surgery, Shanghai Pudong Hospital, Shanghai 201399, China

Corresponding author: Bo Yu. E-mail: yubo120@hotmail.com

© Shanghai Institutes for Biological Sciences, CAS and Springer-Verlag Berlin Heidelberg 2015

ABSTRACT

Autophagy plays a vital role in cerebral ischemia and may be a potential target for developing novel therapy for stroke. In this study, we constructed an autophagy-related pathway network by analyzing the genes related to autophagy and ischemic stroke, and the risk genes were screened. Two autophagy-related modules were significantly up-regulated and clustered to influence cerebral ischemia. Besides, three key modular genes (NFKB1, RELA, and STAT3) were revealed. With 5-fold cross validation, the ROC curves of NFKB1, RELA, and STAT3 were 0.8256, 0.8462, and 0.8923. They formed a complex module and competitively mediated the activation of autophagy in cerebral ischemia. In conclusion, a module containing NFKB1, RELA, and STAT3 mediates autophagy, serving as a potential biomarker for the diagnosis and therapy of ischemic stroke.

Keywords: cerebral ischemia; autophagy; functional module; stroke; gene expression profile

INTRODUCTION

Ischemic stroke is one of the major causes of death and disability^[1–5]. It is generally considered to be a heterogeneous and multifactorial disorder and caused by both conventional environmental risk factors and genetic factors^[6, 7]. It can cause acute neuronal death *via* oxygen and nutrient depletion, and result in disruption of

the blood-brain barrier^[8]. Acute ischemic stroke resulting from intracranial vessel occlusion is associated with high morbidity and mortality^[9]. Therefore, it is pivotal to explore the pathogenesis of ischemic stroke. Moreover, since most ischemic strokes (~80%) occur in the territory of middle cerebral artery (MCA), many animal stroke models of middle cerebral artery occlusion (MCAO) have been developed for studies of ischemic stroke^[10].

Recent research has shown that ischemic insult activates autophagy, and an autophagic mechanism may contribute to ischemic neuronal injury^[11]. Autophagy can be induced by pattern-recognition receptors and stresses such as nutrient depletion, closed head injury, or focal cerebral ischemia. Cerebral ischemia-induced microglial autophagy contributes to ischemic neuronal inflammation and injury. Autophagy is a process for the intracellular bulk degradation of cellular constituents that has multiple effects on immunity^[12–14]. Blocking autophagy in epithelial cells enhances host cell death and finally leads to tissue destruction and inflammation^[15]. Moreover, autophagy eliminates the abnormal protein aggregates and the damaged organelles in neurons after transient cerebral ischemia^[16]. Besides, ischemic stroke increases autophagosomes and activates the autophagy-related pathways (ARPs)^[17]. After neonatal hypoxia-ischemia, autophagy increases in neurons, indicating that over-activation of autophagic pathways is a potential protective mechanism in the early stage of brain injury^[18]. It has been demonstrated that protective autophagy is induced and further promotes the neuroprotective effect on ischemic stroke by regulating mitogen-activated protein kinase

(MAPK) signals^[19]. It has also been reported that global ischemia increases the autophagosomes *via* decreasing autophagosome degradation^[16]. The activation of autophagic and lysosomal pathways has been implicated in neuronal injury in a rat model of permanent focal cerebral ischemia^[11]. Furthermore, by regulating the TSC2-mTOR-S6K1 signaling pathway, autophagy is induced and further promotes neuronal survival during cerebral ischemia^[20].

Based on the above findings, autophagy plays a vital role in cerebral ischemia, and may be a potential target for developing novel therapies for stroke. Thus, it is essential to explore the mechanism of autophagy in cerebral ischemia. In the current study, by using gene-expression and network information, we predicted autophagy-related genes that are highly correlated with ischemic stroke.

MATERIALS AND METHODS

Screening for Differentially-expressed Genes in Ischemic Brain

The National Center for Biotechnology Information (NCBI) GEO database^[21] (<http://www.ncbi.nlm.nih.gov/geo/>) was screened to retrieve the gene expression profiles of the ischemic brain. Through searching for “ischemic brain” in the NCBI GEO database, 3007 results in 158 series were obtained from the DataSets database. Subsequently, we selected “series” for Entry type and “Expression profiling by array” for Study type. Consequently, 54 candidate series of *Mus musculus* and 40 candidate series of *Rattus norvegicus* were acquired. Then, several criteria were applied to screen the candidate series: (1) the tissue must be brain; (2) the study was not related to the reaction of ischemic brain; (3) eliminating series exposed to drug research; (4) eliminating series dealing with hypoxia-ischemia/reperfusion; (5) eliminating samples subjected to MCAO for >24 h. Finally, three series (GSE38037, GSE32529, and GSE58720) were screened for further study, and the samples with MCAO were the test groups. In GSE38037, there were eight samples (4 normal and 4 test samples)^[22]. In GSE32529, there were 224 samples including 6 experimental conditions such as LPS treatment + ischemic challenge, and CpG treatment + ischemic challenge^[23, 24]. In this study, the samples of “brain-unhandled” (i.e. non-treated) were selected as the normal

control, and the brain-ischemia challenged (3 h and 24 h) samples were used as test groups. Finally, 6 normal and 8 test groups were obtained. In GSE58720, the sham and MCAO groups were measured at 24 h after reperfusion. Here, we considered the sham samples as the control group and the MCAO samples as the test groups. Finally, 3 test samples and 3 control groups were selected in GSE58720. The gene expression of each sample was estimated and normalized. Each probeset ID was mapped to gene symbol according to the corresponding platform. If multiple probesets were mapped to the same gene, the expression value for the gene was summarized as the arithmetic mean of the values of multiple probesets (on the log2 scale).

Gene Expression Significance Analysis

With the “MetaDE” package of R project, we merged the three expression profiles and screened the differentially-expressed genes (DEGs) by false discovery rate (FDR) <0.01. Then, Database for annotation, Visualization and Integrated Discovery 6.7 (DAVID, <http://david.abcc.ncifcrf.gov/home.jsp>) software was used to explore the functions of the DEGs, including biological processes and pathways. $P < 0.05$ was set as the threshold used for enrichment analysis of the GO_BP and Kyoto Encyclopedia of Genes and Genomes (KEGG) pathways.

Construction of Autophagy-Related Pathway Network

There were 7 ARPs in KEGG (<http://www.kegg.jp/kegg/pathway.html>): Regulation of autophagy, Lysosome, FoxO signaling pathway, AMPK signaling pathway, Prion diseases, Shigellosis, and mTOR signaling pathways. All the autophagy-related pathways and pathway genes were extracted. To systematically analyze the role of autophagy genes in cerebral ischemia, the STRING v10 database (<http://string-db.org/>) was used to obtain a protein-protein interaction network of *Mus*. By integrating the protein interaction information of DEGs and autophagy-related pathway-gene information, we constructed an ARP^[13] complex network. Four topological properties (degree, average shortest path length, closeness centrality, and clustering coefficient of the ARP network) were analyzed by the “network analyzer” plugin in Cytoscape software (www.cytoscape.org/).

Functional Module Analysis

The “MINE” plugin of Cytoscape was used to identify functional modules in the ARP networks, using a cutoff value for the connectivity degree of nodes (proteins in the network >2). Then, we set the module as a unit and achieved the module expression in different samples by calculating the average expression value of genes that were in the module.

$$M_{ik} = \frac{\sum_{j=1}^n S_{jk}}{n} \quad (1)$$

where M_{ik} is the expression value of module i in sample k . n is the number of genes in module i . S_{jk} is the expression value of module gene j in sample k .

For the “module expression profile”, we further utilized the “limma” package of R project to screen the differentially-expressed modules between the disease and normal groups. Modules with $P < 0.05$ were considered to be significantly expressed.

Identification of Potential Risk Genes

The important autophagy genes in ischemic stroke were screened based on the network topological properties in the ARP network, using Neighborhood scoring (NS) and Interconnectivity (ICN) optimization algorithms. The NS algorithm was based on the distribution of genes in the network and the expression level to screen the candidate genes^[25]. Based on the NS, we weighted every node in the network by combining the fold change of the candidate node and its neighborhoods. If the node was differentially expressed and was linked directly to multiple nodes that were also DEGs, then we considered that the node and its neighbors were correlated. Thus, we inferred that the node was a potential target or diagnosis marker. We followed formula (2) to score the node.

$$Score(i) = \frac{1}{2} * FC_i + \frac{1}{2} * \frac{\sum_{j=1}^{N_i} FC_j}{|N_i|} \quad (2)$$

in which i is the candidate gene i in the network. FC_i represents the fold change of gene i between disease samples and normal samples. N_i is the number of neighbors for node i . j is the neighbor node of candidate gene i . Considering the properties of node self, if node i or its neighbors were not differentially expressed, then the $Score(i)$ was 0.

The ICN^[26] mainly calculates the correlation between the node and all the DEGs, so we followed formula (3) to weight the genes.

$$Score(i) = \frac{1}{n} * \sum_{j=1}^n \frac{2 + N_{i,j}}{\sqrt{N_i * N_j}} \quad (3)$$

where n is the number of all DEGs. N_i is the number of interacting genes for gene i , and N_j is the number of interacting genes for gene j . $N_{i,j}$ is the number of genes shared by gene i and DEG j . In this algorithm, single nodes were rejected. All the genes were ranked in descending order, and a node with a higher score was considered to be more important.

The NS focuses on the expression variation of the node itself and the affected nodes. If more genes are affected and the fold change is more evident, then the gene is more important than other genes in disease development. The ICN focuses on the correlation between the candidate gene and DEGs. If the degree of correlation of a gene with all DEGs is higher, then that gene would be considered more important than others.

RESULTS

Differentially-Expressed Genes

By combining the three expression profiles (GSE38037, GSE32529 and GSE58720), we acquired 15 ischemic samples and 13 normal samples. With the “limma” package of R project, we finally acquired 337 common significant DEGs ($P < 0.05$ and FDR < 0.01) among GSE38037, GSE32529, and GSE58720 (Fig. 1A; Table S1). After functional enrichment, the common DEGs were involved in the Ribosome, Toll-like receptor signaling pathway, MAPK signaling pathway, and the Chemokine signaling pathway, among others (Fig. 1B). Besides, these DEGs played important roles in such processes as vasculature development, blood vessel development, and the regulation of angiogenesis (Fig. 1C).

ARP Network Analysis

Four hundred and ninety-two genes related to the autophagy-associated pathway were acquired from the KEGG database. By integrating the DEGs and autophagy-related genes, an ARP network was constructed that included 1356 nodes and 1983 edges (Fig. 2). We calculated the closeness centrality, average clustering

Fig. 1. Differentially-expressed genes (DEGs) acquired from different studies and function analysis of common DEGs. A, relationships of DEGs among GSE38037, GSE32529, and GSE58720; B, significantly enriched pathways of common DEGs; C, significant biological processes regulated by common DEGs. The larger $-\log(P \text{ value})$, the more important the function.

coefficient, degree distribution, and average shortest path length (Fig. 3). The degree of nodes in the ARP network followed a power-law distribution, and had small-world network characteristics such as a large average clustering coefficient and a short average shortest path length.

Module Analysis in ARP Network

The functional module expression was built for 15 functional modules (Fig. 4A), and two differentially-expressed modules were identified including modules 4 and 10 (Fig. 4B). Using the five-fold cross-validation test, we acquired the ROC curves of all modules (Table 1). Modules 4 and 10 had the highest classification accuracy, with ROC curves >0.85 .

Key Autophagy Genes

We ranked the genes acquired by NS and ICN in

descending order and screened the top 50 genes. Subsequently, we analyzed these 50 genes and obtained five common genes: NFKB1, RELA, STAT3, JAK2, and SHC1. Genes in the same module had similar or the same biological functions to affect the development of cerebral ischemia. Thus, it was likely that the three genes STAT3, NFKB1, and RELA in modules 4 and 10 affect autophagy in the development of ischemic stroke. Here, we focused on the analysis of STAT3, NFKB1 and RELA. At the gene-expression level, the three genes were up-regulated in the disease groups compared with the normal groups (Fig. 4). Five-fold cross validation showed that the ROC curves of all the risk genes were >0.80 (Fig. 4C). Therefore, these genes may be potential risk genes in the development of ischemic stroke, which was highly correlated with the

Fig. 2. The autophagy-related pathway network. Red circles, autophagy-related DEGs; blue circles, autophagy-related genes that were not differentially expressed between disease and normal groups; yellow squares, autophagy-related pathways.

Fig. 3. Topological properties of the ARP network. Closeness centrality, average clustering coefficient, degree distribution, and average shortest path length.

Table 1. Properties of all modules in the ARP network

Modules	Nodes	Edges	P Value	ROC curve	Modular genes
module_1	5	8	0.0313	0.7282	TGFBR2, ACVRL1, CDH5, ENG, TGFBR1
module_2	6	9	0.0019	0.7641	INPP5D, CD22, SOS1, CBL, SHC1, GRB2
module_3	5	7	0.2016	0.5538	PRNP, HSPA5, APP, KCNMA1, PRP
module_4	5	7	0.0006	0.8769	IL6ST, GHR, SHC1, JAK2, STAT3
module_5	4	5	0.0851	0.6718	CCND1, CDKN1B, CDK4, P27
module_6	4	5	0.1274	0.7641	TNFSF6, FASL, PSTPIP1, PTPN12
module_7	4	5	0.1055	0.6872	PSMA1, PPP2R1A, PSMA3, PSMB2
module_8	4	5	0.1799	0.6256	AP2B1, DNAJC6, AP2A2, YWHAZ
module_9	4	5	0.1373	0.6718	LMNA, WDFY2, AKT, FOXO1
module_10	3	3	0.0053	0.8513	HDAC1, RELA, NFKB1
module_11	3	3	0.0044	0.7128	BLNK, GRB2, SH3KBP1
module_12	3	3	0.2375	0.6154	RICTOR, MAPKAP1, FRAP
module_13	3	3	0.1318	0.6769	AP1B1, CALM3, YWHAZ
module_14	3	3	0.2311	0.6564	CCNA2, ABL1, CDK2
module_15	3	3	0.0599	0.7231	CLTA, KCNMA1, YWHAZ

Fig. 4. Modules in the ARP network. A, heat map analysis of module expression. Green to red represents low to high expression. B, differentially-expressed modules in the ARP network. C, expression levels and ROC curves of NFKB1, RELA, STAT3, JAK2, and SHC1.

autophagy related pathways in the ARP network.

DISCUSSION

Autophagy is involved in neuronal death following cerebral ischemia, and plays a pivotal role in the course of ischemic stroke^[27-29]. Moreover, autophagic pathways such as the mTOR pathway are closely related to the ischemic brain^[30]. However, the regulatory network has not been fully explained. In this study, we used the gene expression profile of ischemic stroke from the GEO to explore possible pathogenesis associated with autophagy. By constructing a gene regulation network, we mined highly-correlated modules, and screened autophagy risk genes. Fifteen modules were obtained from the new network, among which modules 4 and 10 were significantly dysregulated, with ROC values >0.8. These two modules contained 5 DEGs, including RELA, NFKB1, and STAT3, and these three genes are closely associated with the autophagic pathway.

The autophagy-related modules were significantly up-regulated and clustered to influence cerebral ischemia. These modules are involved in several pathways including the FOXO signaling pathway and MAPK signaling pathway. The FOXO pathway was associated with most of the modules, and positive regulation of modules would promote the activation of FOXO pathways. Indeed, activation of FOXO pathways is neuroprotective against transient global cerebral ischemia^[31, 32]. The MAPK signaling pathway plays a significant part in cerebral ischemia^[33-35]. MAPK is an upstream regulator of mTORC1 and autophagy can be induced *via* the MAPK-mTOR signaling pathway^[19]. In fact, the mTOR signaling pathway that induces autophagy has neuroprotective effects on ischemic stroke^[19]. This protective autophagy is favorable in the treatment of stroke and avoids unfavorable side-effects^[19]. In addition, our results indicated that genes in two modules interacted with each other and commonly stimulated the PI3K-Akt signaling pathway. Thus, our results verified previous studies showing that autophagy-related modules may play vital roles in cerebral ischemia.

Among all the modular genes in the autophagy-related pathway, three risk genes (NFKB1, RELA, and STAT3) that were differentially expressed in cerebral ischemia were extracted. NFKB1 (also known as NFKB or p50) and RELA form a complex in the NF- κ B signaling

pathway. Commonly, they form a dimer. I κ B α , an inhibitor of NF- κ B, combines with the dimer, sequestering this complex outside the nucleolus. Once I κ B α is degraded, the dimer would be translocated into the nucleolus, initiating transcription^[36] and regulating the expression of numerous genes involved in immunity, inflammation, proliferation, and apoptosis^[37, 38]. Several studies have indicated that dysregulation of these three factors is significantly involved in brain diseases, while suppression of the NF- κ B signaling pathway contributes to neuroprotection against cerebral ischemia^[39-41]. NF- κ B is activated after cerebral ischemia and can potentiate ischemic injury, activating many genes involved in the pathogenesis of cerebral ischemia, such as TNF- α , IL-1 β , IL-6, ICAM-1, iNOS, and COX-2^[42]. Results from immunohistological assay and western blot for NF- κ B (p65) in the ischemic penumbra of rats showed that flurbiprofen inhibits NF- κ B to protect against cerebral ischemia/reperfusion injury^[40]. Brain tissue damage has also been correlated with the dysregulated NFKB1^[43], blockade of which protects the brain against ischemic damage by regulating inflammatory responses^[44, 45]. NFKB1 combines with RELA to form the NFKB1/RELA complex that is a key molecule in the progression of ischemic stroke^[46]. Notably, NFKB1/RELA induces pro-apoptotic transcription in acute brain ischemia and its activation is an important event in ischemic neuronal injury^[47]. Blocking NF- κ B activation or knocking out the p50 subunit of NF- κ B can protect against infarct injury or can result in the development of a smaller infarct volume^[48]. Moreover, I κ B α is an inhibitor of NF- κ B, and its dysregulation prevents brain damage after stroke. The activation of signal transducer and activator of transcription-3 (STAT3), a pivotal part of the JAK-STAT signaling pathway, is reportedly induced by cerebral ischemia^[49-51]. Increased expression of the transcripts of IL-6 and JAK2, which are the essential components of STAT3 activation, is also related to the pathogenesis of cerebral ischemic damage^[51]. All three risk genes were differentially expressed in cerebral ischemia. It is likely that their dysregulation modulates the development of cerebral ischemia and targeting the three genes would reduce post-ischemic brain injury.

Recent findings have demonstrated that NF- κ B complex-related factors (NFKB1 and RELA) also affect the autophagy pathway in cerebral ischemia^[52]. Previous studies have shown that RELA and NFKB1 modulate

canonical autophagy and their activation mediates the repression of autophagy^[53–55]. RELA has been reported to modulate an increase of autophagy by BECN1^[56]. However, in mice with focal cerebral ischemia and NFKB1 knockout, there are significantly more Beclin-1/TUNEL-positive cells than in wild type mice^[52], indicating that cerebral ischemia-induced autophagy-like injury is regulated by the NF- κ B pathway. It has also been found that a high-fat diet is cardioprotective against ischemia-reperfusion injury involving NF- κ B dependent enhancement in autophagy and decreased apoptosis^[57]. Moreover, variations within NFKB1 and I κ B α can potentially influence the function of NF- κ B^[58] and in turn the autophagy process. As autophagy participates in the neuronal death and functional loss induced ischemia/reperfusion injury^[59–61], genes involved in autophagy would be associated with cerebral ischemia. Both RELA and NFKB1 were up-regulated in cerebral ischemia according to our research and further activated the NF- κ B pathway. It has been validated that the NF- κ B pathway regulates the autophagy-like injury induced by cerebral ischemia^[52]. Besides, inhibition of NF- κ B *in vivo* prevents cerebral ischemic injury^[62]. I κ B α interacts with RELA and NFKB1 to form a complex and competitively regulates autophagy in the development of cerebral ischemia. Autophagy induced by the complex in cerebral ischemia would protect against neuronal injury.

In conclusion, Autophagy is a basic catabolic progress for cell survival under stress. Using a computational bioinformatics approach, we obtained two autophagy-related modules. In accordance with previous reports, our result also verified that the FOXO and MAPK signaling pathways are important in cerebral ischemia. Besides, NF- κ B complex-related factors (NFKB1 and RELA) and STAT3 play a key role in inducing autophagy in the development of cerebral ischemia, and are potential targets of therapy.

ELECTRONIC SUPPLEMENTARY MATERIAL

Supplementary material is available in the online version of this article at <http://dx.doi.org/10.1007/s12264-015-1547-3>.

ACKNOWLEDGEMENTS

This work was supported by a Shanghai Leading Talents Project (54[2012]) and a Pudong New Area Leading Talents Project of Health System (PWR12011-02).

Received date: 2015-04-30; Accepted date: 2015-07-09

REFERENCES

- [1] Kishore A, Vail A, Majid A, Dawson J, Lees KR, Tyrrell PJ, *et al.* Detection of atrial fibrillation after ischemic stroke or transient ischemic attack: a systematic review and meta-analysis. *Stroke* 2014, 45: 520–526.
- [2] Chen L, Luo S, Yan L, Zhao W. A systematic review of closure versus medical therapy for preventing recurrent stroke in patients with patent foramen ovale and cryptogenic stroke or transient ischemic attack. *J Neurol Sci* 2014, 337: 3–7.
- [3] Saada F, Antonios N. Existence of ipsilateral hemiparesis in ischemic and hemorrhagic stroke: two case reports and review of the literature. *Eur Neurol* 2014, 71: 25–31.
- [4] Hong KS, Saver JL, Kang DW, Bae HJ, Yu KH, Koo J, *et al.* Years of optimum health lost due to complications after acute ischemic stroke: disability-adjusted life-years analysis. *Stroke* 2010, 41: 1758–1765.
- [5] Lees KR, Bluhmki E, von Kummer R, Brodt TG, Toni D, Grotta JC, *et al.* Time to treatment with intravenous alteplase and outcome in stroke: an updated pooled analysis of ECASS, ATLANTIS, NINDS, and EPITHET trials. *Lancet* 2010, 375: 1695–1703.
- [6] Rosamond W, Flegal K, Furie K, Go A, Greenlund K, Haase N, *et al.* Heart disease and stroke statistics--2008 update: a report from the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. *Circulation* 2008, 117: e25–146.
- [7] Dichgans M. Genetics of ischaemic stroke. *Lancet Neurol* 2007, 6: 149–161.
- [8] Reischl S, Li L, Walkinshaw G, Flippin LA, Marti HH, Kunze R. Inhibition of HIF prolyl-4-hydroxylases by FG-4497 reduces brain tissue injury and edema formation during ischemic stroke. *PLoS One* 2014, 9: e84767.
- [9] Walcott BP, Boehm KM, Stapleton CJ, Mehta BP, Nahed BV, Ogilvy CS. Retrievable stent thrombectomy in the treatment of acute ischemic stroke: analysis of a revolutionizing treatment technique. *J Clin Neurosci* 2013, 20: 1346–1349.
- [10] Chiang T, Messing RO, Chou WH. Mouse model of middle cerebral artery occlusion. *J Vis Exp* 2011, 13: 2761.
- [11] Wen YD, Sheng R, Zhang LS, Han R, Zhang X, Zhang XD, *et al.* Neuronal injury in rat model of permanent focal cerebral ischemia is associated with activation of autophagic and lysosomal pathways. *Autophagy* 2008, 4: 762–769.
- [12] Heymann D. Autophagy: A protective mechanism in response to stress and inflammation. *Curr Opin Investig Drugs* 2006, 7: 443–450.
- [13] White E, Karp C, Strohecker AM, Guo Y, Mathew R. Role of autophagy in suppression of inflammation and cancer. *Curr*

- Opin Cell Biol 2010, 22: 212–217.
- [14] Luciani A, Vilella VR, Esposito S, Brunetti-Pierri N, Medina D, Settembre C, *et al.* Defective CFTR induces aggresome formation and lung inflammation in cystic fibrosis through ROS-mediated autophagy inhibition. *Nat Cell Biol* 2010, 12: 863–875.
- [15] Chang SY, Lee SN, Yang JY, Kim DW, Yoon JH, Ko HJ, *et al.* Autophagy controls an intrinsic host defense to bacteria by promoting epithelial cell survival: a murine model. *PLoS One* 2013, 8: e81095.
- [16] Liu C, Gao Y, Barrett J, Hu B. Autophagy and protein aggregation after brain ischemia. *J Neurochem* 2010, 115: 68–78.
- [17] Yang Y, Gao K, Hu Z, Li W, Davies H, Ling S, *et al.* Autophagy upregulation and apoptosis downregulation in DAHP and triptolide treated cerebral ischemia. *Mediators Inflamm* 2015, 2015: 120198.
- [18] Carloni S, Buonocore G, Balduini W. Protective role of autophagy in neonatal hypoxia-ischemia induced brain injury. *Neurobiol Dis* 2008, 32: 329–339.
- [19] Wang PR, Wang JS, Zhang C, Song XF, Tian N, Kong LY. Huang-Lian-Jie-Du-Decotion induced protective autophagy against the injury of cerebral ischemia/reperfusion via MAPK-mTOR signaling pathway. *J Ethnopharmacol* 2013, 149: 270–280.
- [20] Wang P, Guan YF, Du H, Zhai QW, Su DF, Miao CY. Induction of autophagy contributes to the neuroprotection of nicotinamide phosphoribosyltransferase in cerebral ischemia. *Autophagy* 2012, 8: 77–87.
- [21] Barrett T, Troup DB, Wilhite SE, Ledoux P, Rudnev D, Evangelista C, *et al.* NCBI GEO: mining tens of millions of expression profiles--database and tools update. *Nucleic Acids Res* 2007, 35: D760–765.
- [22] Wang IM, Zhang B, Yang X, Zhu J, Stepaniants S, Zhang C, *et al.* Systems analysis of eleven rodent disease models reveals an inflammatome signature and key drivers. *Mol Syst Biol* 2012, 8: 594.
- [23] Stevens SL, Leung PY, Vartanian KB, Gopalan B, Yang T, Simon RP, *et al.* Multiple preconditioning paradigms converge on interferon regulatory factor-dependent signaling to promote tolerance to ischemic brain injury. *J Neurosci* 2011, 31: 8456–8463.
- [24] Vartanian KB, Stevens SL, Marsh BJ, Williams-Karnesky R, Lessov NS, Stenzel-Poore MP. LPS preconditioning redirects TLR signaling following stroke: TRIF-IRF3 plays a seminal role in mediating tolerance to ischemic injury. *J Neuroinflammation* 2011, 8: 140.
- [25] Nitsch D, Goncalves JP, Ojeda F, de Moor B, Moreau Y. Candidate gene prioritization by network analysis of differential expression using machine learning approaches. *BMC Bioinformatics* 2010, 11: 460.
- [26] Hsu CL, Huang YH, Hsu CT, Yang UC. Prioritizing disease candidate genes by a gene interconnectedness-based approach. *BMC Genomics* 2011, 12 Suppl 3: S25.
- [27] Harris H, Rubinshtein DC. Control of autophagy as a therapy for neurodegenerative disease. *Nat Rev Neurol* 2012, 8: 108–117.
- [28] Gustafsson AB, Gottlieb RA. Recycle or die: the role of autophagy in cardioprotection. *J Mol Cell Cardiol* 2008, 44: 654–661.
- [29] Jiang M, Liu K, Luo J, Dong Z. Autophagy is a renoprotective mechanism during in vitro hypoxia and in vivo ischemia-reperfusion injury. *Am J Pathol* 2010, 176: 1181–1192.
- [30] Baek SH, Noh AR, Kim KA, Akram M, Shin YJ, Kim ES, *et al.* Modulation of mitochondrial function and autophagy mediates carnosine neuroprotection against ischemic brain damage. *Stroke* 2014, 45: 2438–2443.
- [31] Zhan L, Li D, Liang D, Wu B, Zhu P, Wang Y, *et al.* Activation of Akt/FoxO and inactivation of MEK/ERK pathways contribute to induction of neuroprotection against transient global cerebral ischemia by delayed hypoxic preconditioning in adult rats. *Neuropharmacology* 2012, 63: 873–882.
- [32] Zhan L, Wang T, Li W, Xu ZC, Sun W, Xu E. Activation of Akt/FoxO signaling pathway contributes to induction of neuroprotection against transient global cerebral ischemia by hypoxic pre-conditioning in adult rats. *J Neurochem* 2010, 114: 897–908.
- [33] Liu XQ, Sheng R, Qin ZH. The neuroprotective mechanism of brain ischemic preconditioning. *Acta Pharmacologica Sinica* 2009, 30: 1071–1080.
- [34] Mehta SL, Manhas N, Raghubir R. Molecular targets in cerebral ischemia for developing novel therapeutics. *Brain Res Rev* 2007, 54: 34–66.
- [35] Nozaki K, Nishimura M, Hashimoto N. Mitogen-activated protein kinases and cerebral ischemia. *Mol Neurobiol* 2001, 23: 1–19.
- [36] Hayden MS, Ghosh S. Signaling to NF-kappaB. *Genes Dev* 2004, 18: 2195–2224.
- [37] Glavac D, Ravnik-Glavac M, O'Brien SJ, Dean M. Polymorphisms in the 3' untranslated region of the I kappa B/MAD-3 (NFKBI) gene located on chromosome 14. *Hum Genet* 1994, 93: 694–696.
- [38] Nakshatri H, Bhat-Nakshatri P, Martin DA, Goulet RJ, Jr., Sledge GW, Jr. Constitutive activation of NF-kappaB during progression of breast cancer to hormone-independent growth. *Mol Cell Biol* 1997, 17: 3629–3639.
- [39] Luo Y, Yang YP, Liu J, Li WH, Yang J, Sui X, *et al.* Neuroprotective effects of madecassoside against focal cerebral ischemia reperfusion injury in rats. *Brain Res* 2014, 1565: 37–47.

- [40] Sun BZ, Chen L, Wu Q, Wang HL, Wei XB, Xiang YX, *et al.* Suppression of inflammatory response by flurbiprofen following focal cerebral ischemia involves the NF-kappaB signaling pathway. *Int J Clin Exp Med* 2014, 7: 3087–3095.
- [41] Freemerman AJ, Gallegos A, Powis G. Nuclear factor kappaB transactivation is increased but is not involved in the proliferative effects of thioredoxin overexpression in MCF-7 breast cancer cells. *Cancer Res* 1999, 59: 4090–4094.
- [42] Seegers H, Grillon E, Trioullier Y, Vath A, Verna JM, Blum D. Nuclear factor-kappa B activation in permanent intraluminal focal cerebral ischemia in the rat. *Neurosci Lett* 2000, 288: 241–245.
- [43] Zhang N, Cheng GY, Liu XZ, Zhang FJ. Expression of Bcl-2 and NF-kappaB in brain tissue after acute renal ischemia-reperfusion in rats. *Asian Pac J Trop Med* 2014, 7: 386–389.
- [44] Zhang S, Zis O, Ly PT, Wu Y, Zhang S, Zhang M, *et al.* Down-regulation of MIF by NFkappaB under hypoxia accelerated neuronal loss during stroke. *FASEB J* 2014, 28: 4394–4407.
- [45] Wang L, Li Z, Zhang X, Wang S, Zhu C, Miao J, *et al.* Protective effect of shikonin in experimental ischemic stroke: attenuated TLR4, p-p38MAPK, NF-kappaB, TNF-alpha and MMP-9 expression, up-regulated claudin-5 expression, ameliorated BBB permeability. *Neurochem Res* 2014, 39: 97–106.
- [46] Lanzillotta A, Sarnico I, Ingrassia R, Boroni F, Branca C, Benarese M, *et al.* The acetylation of RelA in Lys310 dictates the NF-kappaB-dependent response in post-ischemic injury. *Cell Death Dis* 2010, 1: e96.
- [47] Inta I, Paxian S, Maegele I, Zhang W, Pizzi M, Spano P, *et al.* Bim and Noxa are candidates to mediate the deleterious effect of the NF-kappa B subunit RelA in cerebral ischemia. *J Neurosci* 2006, 26: 12896–12903.
- [48] Yi JH, Park SW, Kapadia R, Vemuganti R. Role of transcription factors in mediating post-ischemic cerebral inflammation and brain damage. *Neurochem Int* 2007, 50: 1014–1027.
- [49] Li HC, Zhang GY. Activation of STAT3 induced by cerebral ischemia in rat hippocampus and its possible mechanisms. *Sheng Li Xue Bao* 2003, 55: 311–316.
- [50] Li HC, Zhang GY. Inhibitory effect of genistein on activation of STAT3 induced by brain ischemia/reperfusion in rat hippocampus. *Acta Pharmacol Sin* 2003, 24: 1131–1136.
- [51] Samraj AK, Muller AH, Grell AS, Edvinsson L. Role of unphosphorylated transcription factor STAT3 in late cerebral ischemia after subarachnoid hemorrhage. *J Cereb Blood Flow Metab* 2014, 34: 759–763.
- [52] Li WL, Yu SP, Chen D, Yu SS, Jiang YJ, Genetta T, *et al.* The regulatory role of NF-kappaB in autophagy-like cell death after focal cerebral ischemia in mice. *Neuroscience* 2013, 244: 16–30.
- [53] Criollo A, Chereau F, Malik SA, Niso-Santano M, Marino G, Galluzzi L, *et al.* Autophagy is required for the activation of NFkappaB. *Cell Cycle* 2012, 11: 194–199.
- [54] Djavaheri-Mergny M, Amelotti M, Mathieu J, Besancon F, Bauvy C, Codogno P. Regulation of autophagy by NFkappaB transcription factor and reactive oxygen species. *Autophagy* 2007, 3: 390–392.
- [55] Copetti T, Demarchi F, Schneider C. p65/RelA binds and activates the beclin 1 promoter. *Autophagy* 2009, 5: 858–859.
- [56] Copetti T, Bertoli C, Dalla E, Demarchi F, Schneider C. p65/RelA modulates BECN1 transcription and autophagy. *Mol Cell Biol* 2009, 29: 2594–2608.
- [57] Haar L, Ren X, Liu Y, Koch SE, Goines J, Tranter M, *et al.* Acute consumption of a high-fat diet prior to ischemia-reperfusion results in cardioprotection through NF-kappaB-dependent regulation of autophagic pathways. *Am J Physiol Heart Circ Physiol* 2014, 307: H1705–1713.
- [58] Curran JE, Weinstein SR, Griffiths LR. Polymorphic variants of NFKB1 and its inhibitory protein NFKBIA, and their involvement in sporadic breast cancer. *Cancer Lett* 2002, 188: 103–107.
- [59] Shao ZQ, Liu ZJ. Neuroinflammation and neuronal autophagic death were suppressed via Rosiglitazone treatment: New evidence on neuroprotection in a rat model of global cerebral ischemia. *J Neurol Sci* 2015, 349: 65–71.
- [60] Ravikumar B, Sarkar S, Davies JE, Futter M, Garcia-Arencibia M, Green-Thompson ZW, *et al.* Regulation of mammalian autophagy in physiology and pathophysiology. *Physiol Rev* 2010, 90: 1383–1435.
- [61] Chen W, Sun Y, Liu K, Sun X. Autophagy: a double-edged sword for neuronal survival after cerebral ischemia. *Neural Regen Res* 2014, 9: 1210–1216.
- [62] Wang R, Liang S, Yue H, Chen L. Using a novel in vivo model to study the function of nuclear factor kappa B in cerebral ischemic injury. *Med Sci Monit* 2012, 18: BR461–467.