Published in final edited form as:

Trends Cell Biol. 2017 September ; 27(9): 621–622. doi:10.1016/j.tcb.2017.07.002.

Nuclear Pore Complexes: a Scaffold Regulating Developmental Transcription?

Atsushi Satomura and Jason H. Brickner*

Department of Molecular Biosciences, Northwestern University, Evanston, IL, USA

Abstract

Nuclear Pore Complexes (NPCs) play a conserved, but poorly understood, role in transcriptional regulation. Recently in *Developmental Cell*, Raices *et al.* argues that tissue-specific nuclear pore proteins act as scaffolds that recruit the transcription factor Mef2C to the NPC, promoting transcription of NPC-associated genes during muscle development.

Keywords

Nuclear Pore Complex; Transcription; Differentiation; Nuclear architecture

The Nuclear Pore Complex (NPC) is a large, selective channel of ~110 MDa, composed of 30 different nuclear pore proteins (Nups). NPCs act as a highly selective, bidirectional transporter between the cytoplasm and nucleoplasm. In addition to this role, NPCs also physically interact with chromatin, impacting the positioning of genes within the nucleus and influencing transcriptional regulation and chromatin structure. In budding yeast, hundreds of active genes interact with the NPC [1]. This interaction leads to positioning of genes to the nuclear periphery and, in several cases, has been shown to promote stronger expression [2, 3]. Likewise, in *Drosophila* and mammals, thousands of genes interact with Nups and these interactions have both positive and negative effects on transcription [4]. However, in flies and mammals, Nup interactions occur both at the NPC and in the nucleoplasm, with soluble Nups. It has been unclear if these two types of interactions are functionally equivalent and if the yeast system reflects the spatial organization of transcription in metazoans.

In mammalian cells, the composition of NPCs varies among different cell types and tissues, raising the possibility that different tissues utilize functionally distinct forms of the NPC to facilitate differentiation. For example, Nup210, an integral membrane Nup, is expressed only in a subset of differentiated tissues such as neurons and muscle cells [5]. Knockdown of Nup210 downregulates several genes and prevents myogenic differentiation. Nup210-

*Correspondence: j-brickner@northwestern.edu (J.H. Brickner).

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

dependent expression is independent of nucleo-cytoplasmic transport through NPCs [5]. This suggests that NPCs directly modulate gene expression to promote differentiation.

Using Zebrafish muscle differentiation and mouse C2C12 myoblast cell line as models, Raices *et al.* explored how Nup210 promotes muscle differentiation [6]. Knockdown of Nup210 in Zebrafish led to shortened muscle fibers and progressive accumulation of actin at the myoseptum due to a defect in myofiber maturation and muscle growth during development. These phenotypes correlated with changes in gene expression, including defects in activation of muscle-specific transcriptional programs. The same phenotypes are associated with loss of a transcription factor necessary for muscle development, Mef2C. The transcriptional changes were highly correlated with the binding of Nup210 and co-immunoprecipitation and *in vivo* assays revealed a physical interaction between Mef2C and Nup210, through the adaptor protein Trip6, at the NPC (Figure 1A). This finding suggests that Nup210 is required for Mef2C-mediated transcriptional activation. DNA-FISH in C2C12 cells showed that Nup210/Mef2C-regulated genes localize at the nuclear periphery, suggesting that transcriptional activation is occurring at the NPC. Thus, Nup210 recruitment of Mef2C to the NPC promotes expression of Mef2C target genes and proper muscle development.

Although many active genes localize at the nuclear periphery and physically interact with the NPC in budding yeast, repressed genes also interact with the NPC and, in other eukaryotes, positioning at the nuclear periphery frequently correlates with transcriptional silencing. Certainly, interactions with the nuclear lamina are associated with transcriptional silencing and heterochromatin [7]. However, the work by D'Angelo *et al.* highlights an important role for NPCs in promoting gene expression during vertebrate development. Likewise, some super-enhancers that play critical roles in development interact with the NPC [8]. Therefore, the nuclear periphery is a complex environment and physical interactions between genes and the NPC can both promote or inhibit transcription. Furthermore, the molecular outputs of NPC-gene interactions appear to have been conserved from yeast to human and likely represent a fundamental mechanism of regulation of gene expression.

Nup210 may anchor a wide variety of genes and transcription factors in different cell types, because Nup210 is also involved in neural differentiation (Fig. 1B; ref. 5). However, it is unclear how the target genes get to the NPC; in quiescent myoblasts that do not express Nup210, Nup210 target genes localize at the nuclear periphery. This suggests that these genes are positioned at the nuclear periphery either through another NPC interaction or through a non-NPC interaction. In yeast, transcription factor binding is both necessary and sufficient for gene localization to nuclear periphery and many different transcription factors mediate the repositioning of different genes [2, 9]. Because Nup210 is required for proper binding of Mef2C, this suggests that Nup210 functions as a scaffold to facilitate Mef2C binding to NPC-associated genes. Perhaps Nup210 target genes are positioned at the NPC by transcription factors other than Mef2C. If so, then transcription factors could control either gene positioning, gene expression or both. This would be consistent with work in yeast showing that enhancers that control gene positioning are not always the same enhancers that

control transcription [2]. This suggests that there may be adaptive advantages to controlling these two phenomena separately.

Some exciting questions remain to be answered. Mef2C localizes both at nuclear periphery and in the nucleoplasm, indicating that some genes regulated by Mef2C are expressed independently on Nup210. What are the mechanistic differences between the transcription occurring at the NPC and in the nucleoplasm? Global expression analysis showed that loss of Nup210 or Mef2C resulted in both upregulation and downregulation of mRNA levels. Similar results have been obtained with other Nups [10]. What explains these two distinct classes? Are genes both activated and repressed by interactions with the NPC? Along these lines, are all Nup210-containing NPCs functionally equivalent or are there functional differences between them? Finally, do cell type-specific Nups function as determinants of cell identity or simply as generic cofactors for transcription factors? If the former, what determines the specificity of their function? The answers to these questions are bound to alter our perspective on the molecular and cellular drivers of developmental gene expression.

References

1. Casolari JM, et al. Genome-wide localization of the nuclear transport machinery couples transcriptional status and nuclear organization. *Cell*. 2004; 117(4):427–39. [PubMed: 15137937]
2. Ahmed S, et al. DNA zip codes control an ancient mechanism for gene targeting to the nuclear periphery. *Nat Cell Biol*. 2010; 12(2):111–8. [PubMed: 20098417]
3. Taddei A, et al. Nuclear pore association confers optimal expression levels for an inducible yeast gene. *Nature*. 2006; 441(7094):774–8. [PubMed: 16760983]
4. Sood V, Brickner JH. Nuclear pore interactions with the genome. *Curr Opin Genet Dev*. 2014; 25:43–9. [PubMed: 24480294]
5. D'Angelo MA, et al. A change in nuclear pore complex composition regulates cell differentiation. *Dev Cell*. 2012; 22(2):446–58. [PubMed: 22264802]
6. Raices M, et al. Nuclear Pores Regulate Muscle Development and Maintenance by Assembling a Localized Mef2C Complex. *Dev Cell*. 2017; 41(5):540–554. e7. [PubMed: 28586646]
7. Pombo A, Dillon N. Three-dimensional genome architecture: players and mechanisms. *Nat Rev Mol Cell Biol*. 2015; 16(4):245–57. [PubMed: 25757416]
8. Ibarra A, et al. Nucleoporin-mediated regulation of cell identity genes. *Genes Dev*. 2016; 30(20):2253–2258. [PubMed: 27807035]
9. Randise-Hinchliff C, et al. Strategies to regulate transcription factor-mediated gene positioning and interchromosomal clustering at the nuclear periphery. *J Cell Biol*. 2016
10. Liang Y, et al. Dynamic association of NUP98 with the human genome. *PLoS Genet*. 2013; 9(2):e1003308. [PubMed: 23468646]

Figure 1. The NPC as a regulatory scaffold for developmental gene expression

A. Model for Nup210 regulation of Mef2C. In myoblasts, expression of Nup210 recruits Mef2C through interaction with the adaptor protein Trip6. Mef2C binds to enhancers associated with genes critical for muscle development, activating their expression. B. Hypothetical Nup210 regulation of neural genes. Nup210 is required for differentiation of neural progenitor cells and, akin to its role in muscle development, could function as a scaffold to recruit transcription factors to NPC-associated genes involved in neural differentiation.